

NOTICE WARNING CONCERNING COPYRIGHT RESTRICTIONS:
The copyright law of the United States (title 17, U.S. Code) governs the making of photocopies or other reproductions of copyrighted material. Any copying of this document without permission of its author may be prohibited by law.

ON A SECOND ORDER NONLINEAR
OSCILLATION PROBLEM

C. V. Coffman

and

J. S. W. Wong

Report 69-2

January, 1969

University Libraries
Carnegie Mellon University
Pittsburgh PA 15213-3890

ON A SECOND ORDER NONLINEAR
OSCILLATION PROBLEM

C. V. Coffman* and J. S. W. Wong

1. INTRODUCTION.

We are here concerned with the oscillatory behavior of solutions of the following second order nonlinear differential equation:

$$(1) \quad y'' + yF(y^2, x) = 0,$$

where the function $F(t, x)$ is continuous and non-negative for $t \in [0, \infty)$, $x \in (0, \infty)$.

It will be tacitly assumed here that every locally defined solution of (1) is continuously extendable throughout the entire non-negative real axis. This will be the case if for example one requires that for fixed t , $F(t, x)$ satisfies a uniform Lipschitz condition in some neighborhood of every $x \in [0, \infty)$. (See Hastings [3], and Coffman and Ullrich [2].) Actually, this tacit assumption can easily be removed, see the remarks at the end of the paper. A nontrivial solution of (1) is said to be non-oscillatory if for every $a > 0$, the number of its zeros in $[a, \infty)$ is finite, and it is said to be oscillatory otherwise. Different from the linear equation, when $F(t, x)$ is independent of t , the nonlinear equation may possess solutions of either kind. In view of this,

*The research of the first-named author was supported by NSF GP-7662.

MAR 21 '68

HUNT LIBRARY
CARNEGIE-MELLON UNIVERSITY

one is led to consider the following types of oscillation and non-oscillation conditions; namely, those which guarantee all solutions of (1) oscillate and its converse, i.e. the existence of one non-oscillatory solution, and those which guarantee all solutions of (1) do not oscillate and its converse, i.e. the existence of one oscillatory solution. The first type of oscillation and non-oscillation conditions have been the centre of considerable amount of research and there are a number of results available for equation (1) or similar equations. For an expository account on this subject, we refer the reader to Wong [12], where other references may be found. An excellent discussion on the nature of oscillatory and non-oscillatory solutions may also be found in the papers by Moore and Nehari [9], and Nehari [10]. The second type of oscillation and non-oscillation conditions have received little attention only until recently. The prototype of equation (1) is the following generalized Emden-Fowler equation:

$$(2) \quad y'' + p(x)y^\gamma = 0,$$

where $\gamma \geq 1$ is the quotient of two odd integers. For equation (2), Jasny [4] and Kurzweil [7] have established the following result on the existence of one oscillatory solution:

THEOREM A. Let $\gamma > 1$. If $x^{\frac{\gamma+3}{2}} p(x)$ is non-decreasing in x , then equation (2) has an oscillatory solution.

Jasny and Kurzweil's result is complemented by the following theorem of Kiguradze [5]:

THEOREM B. If $x^{\frac{\gamma+3}{2} + \epsilon} p(x)$ is non-increasing in x , for some $\epsilon > 0$ then equation (2) is non-oscillatory, i.e. all solutions are non-oscillatory.

Very recently, Nehari [11] obtained a result which is an improvement of the result of Kiguradze [5], namely,

THEOREM C. If $(x \log x)^{\frac{\gamma+3}{2}} p(x)$ is non-increasing in x for sufficiently large x , then equation (2) is non-oscillatory.

In [11], Nehari also initiated the study of obtaining similar results for the more general equation (1). Part of the difficulty of such an extension lies in the fact that the proofs of Theorems A, B, and C depend heavily on the form of the function $F(t,x)$ in that it is separable as a product of functions of t and x . Using some ingenious differential identities and inequalities, Nehari [11] obtained the following non-oscillation result for equation (1) corresponding to Theorem B for equation (2).

THEOREM D. Let $G(t,x)$ be defined by

$$(3) \quad G(t,x) = \int_0^t F(s,x) ds.$$

If (i) for fixed x , $F(t,x)$ is a non-decreasing function of t and (ii) for some positive ϵ and all positive α , $xG(\alpha x^{1+\epsilon}, x)$

is non-increasing for $x \in [a, \infty)$ for some $a > 0$, then equation (1) is non-oscillatory.

Questions were open as to whether (a) a similar result for equation (1) corresponding to Theorem A for equation (2) holds, and (b) Theorem D may be improved to include Theorem C as a special case. The purpose of the present work is to answer these questions in the affirmative, thus completing the extension of Theorems A and C to the more general equation (1). In fact, we prove a little more than that stated above, and we refer the reader to the last section where a discussion of these details will be given.

2. OSCILLATION THEOREM.

The desired extension of Theorem A to the nonlinear equation (1) is the following result:

THEOREM 1. Let $F(t, x)$ and $G(t, x)$ be given as above.

Suppose that (i) there exist constants $x_0, M, c > 0$ and $K \geq 1$ such that

$$(4) \quad x^2 F(\alpha x, x) \geq \frac{1}{4} + c, \quad x \geq x_0, \quad \alpha \geq M,$$

and

$$(5) \quad G(t, x) \leq KtF(t, x), \quad x \geq x_0, \quad t \leq Mx,$$

and (ii) for every $\alpha > 0$ the function $xG(\alpha x, x)$ is non-decreasing in x for $x \geq x_0$. Then equation (1) has an oscillatory solution.

The proof will be carried out in a series of three lemmas each of which may be of interest in itself. In each of these, we

assume without explicit mention, that the hypotheses of the above theorem hold.

Lemma 1. Let $y(x)$ be a non-oscillatory solution of (1).

Then

$$(6) \quad \liminf_{x \rightarrow \infty} x^2 F(y^2(x), x) \leq \frac{1}{4}.$$

Proof. Suppose that (6) fails, then there must exist a $\delta > 0$ and an $x_1 \geq x_0$ such that for $x \geq x_1$

$$x^2 F(y^2(x), x) \geq \frac{1+\delta}{4}.$$

But then $y(x)$ satisfies a linear equation:

$$(7) \quad y'' + p(x)y = 0,$$

where $p(x) \geq \frac{1+\delta}{4x^2}$, for $x \geq x_1$. Since equation (7) is oscillatory when $p(x) = \frac{1+\delta}{4x^2}$, the given solution $y(x)$ must also be oscillatory by the Sturm Comparison Theorem. This contradicts the assumption that y is non-oscillatory. Thus (6) must hold.

Lemma 2. Let $y(x)$ be a non-oscillatory solution of (1).

Then there exists a positive constant B , independent of the initial values of $y(x)$ and $y'(x)$ such that

$$(8) \quad \limsup_{x \rightarrow \infty} \left| x^{1/2} y'(x) - \frac{1}{2} x^{-1/2} y(x) \right| \leq B.$$

Proof. Let $\phi(x) = x^{1/2} y'(x) - \frac{1}{2} x^{-1/2} y(x)$. We note first that, since $y(x)$ is a solution of (1), we have,

$$(9) \quad \frac{d\phi(x)}{dx} = x^{-3/2} y(x) \left(\frac{1}{4} - x^2 F(y^2(x), x) \right).$$

Now if we rewrite (1) as

$$y'' + \frac{1}{4x^2} y = \frac{y}{4x^2} \left(\frac{1}{4} - x^2 F(y^2(x), x) \right),$$

and apply the variation of constants formula, we obtain

$$(10) \quad y(x) = x^{1/2} \left(a + b \log \frac{x}{x_1} \right) + x^{1/2} \int_{x_1}^x s^{-3/2} \log \frac{x}{s} \left(\frac{1}{4} - s^2 F(y^2(s), s) \right) y(s) ds,$$

where $a = x_1^{-1/2} y(x_1)$ and $b = \phi(x_1)$. Assume that y is positive for $x \geq x_1$ and that

$$(11) \quad y(x) < (Mx)^{1/2},$$

when $x = x_1$. By continuity then (11) holds on some interval $[x_1, x_2)$, and on that interval we obtain, using (11) in (10),

$$y(x) \leq x^{1/2} \left(a + b \log \frac{x}{x_1} \right) + \frac{1}{4} M^{1/2} x^{1/2} \int_{x_1}^x s^{-1} \log \frac{x}{s} ds,$$

or, using $a = x_1^{-1/2} y(x_1) < M^{1/2}$,

$$(12) \quad y(x) \leq x^{1/2} \left(M^{1/2} + b \log \frac{x}{x_1} + \frac{1}{8} M^{1/2} \left(\log \frac{x}{x_1} \right)^2 \right),$$

for $x_1 \leq x < x_2$. If the logarithmic polynomial

$$P(\xi) = M^{1/2} + b \log \xi + \frac{1}{8} M^{1/2} (\log \xi)^2$$

satisfies

$$M^{1/2} > P(\xi) > 0, \quad \text{for } 1 < \xi < A,$$

then, by (12), the inequality (11) will hold also when $x = x_2$, provided $x_2 < Ax_1$. This last fact can be used to show that the

validity of (12) extends to $[x_1, Ax_1]$. Now let $A > 1$ be given, there exists then a $B > 0$, such that if $-b > B$ the logarithmic polynomial $P(\xi)$ defined above satisfies

$$M^{1/2} > P(\xi) > 0 \text{ for } 1 \leq \xi < A', \quad P(A') = 0,$$

where $A' = A'(b) < A$; moreover B depends only on A and M . The foregoing remarks show that if y is, as above, positive for $x \geq x_1$ and satisfies (11) at $x = x_1$ then $\phi(x_1) > -B$. Since, in view of (4), the right hand side of (9) is negative for $y(x) \geq (Mx)^{1/2}$, it follows from (6) and (4) that $\phi(x) > -B$ for all $x \geq x_1$. Repeating the same argument, but "going backwards" we see that if x_0 is the last zero of the eventually positive solution y , then for $x_1 > Ax_0$, and when (11) holds at $x = x_1$ we must have $\phi(x_1) < B$. Using (4), (6), and (9) as before we conclude that $\phi(x) < B$ for all large x . If $y(x) < 0$ for all large x , then the above argument applied to the solution $-y(x)$ gives (8). This completes the proof.

Lemma 3. If $y(x)$ is a solution of (1), then the function $\Psi(x)$ defined by

$$\Psi(x) = x(y'(x))^2 + G(y^2(x), x) - y(x)y'(x)$$

is non-decreasing in x .

Proof. We give a proof for the case where $F(t, x)$ is of class C^1 . For the general case, the result will follow by a standard argument involving approximation of $F(t, x)$ by C^1 functions. By a straightforward computation we have the identity

$$\frac{d}{dx} \Psi(x) = (2xy'(x) - y(x)) [y''(x) + y(x)F(y^2(x), x)] + Q(y^2, x),$$

where $Q(t, x) = G(t, x) + tF(t, x) + xG_x(t, x)$. Note that the function $Q(t, x)$ satisfies

$$Q(t, x) = \frac{\partial}{\partial \sigma} (\sigma G(\frac{t\sigma}{x}, \sigma)) \Big|_{\sigma = x},$$

for arbitrary $t, x > 0$. Thus, in view of the nondecreasing character of $xG(\alpha x, x)$ for $\alpha > 0$, we have $Q(t, x) \geq 0$, and the desired conclusion follows.

Proof of Theorem 1. Let $y(x)$ be a solution of (1), vanishing at $x_1 \geq x_0$, and let x_2 be some value of $x \geq x_1$ for which

$$(13) \quad x_2^2 F(y^2(x_2), x_2) \leq \frac{1}{4} + c,$$

so that in particular from (4) we have

$$(14) \quad y^2(x_2) \leq Mx_2.$$

Suppose that $y(x_2) > 0$ and $y'(x_2) > 0$, then, by Lemma 3, it follows

$$(15) \quad x_2 [y'^2(x_2) + G(y^2(x_2), x_2)] \geq C_0,$$

where the constant $C_0 = x_1 y'^2(x_1)$. By (5), (13), and (14), we have

$$(16) \quad x_2 G(y^2(x_2), x_2) \leq Kx_2 y^2(x_2) F(y^2(x_2), x_2) \leq KM(\frac{1}{4} + c).$$

Suppose that $y(x)$ is any solution of (1) defined by the initial values $y(x_1) = 0$ and $x_1 y'^2(x_1) \geq B^2$, where $B^2 > 2B^2$

+ $\frac{M}{2}$ + $(\frac{1}{4} + c)KM$, then, it follows from (15) and (16) that for arbitrary large x_2 for which (13) is satisfied,

$$\begin{aligned} \phi(x_2) &= x_2^{1/2} y'(x_2) - \frac{1}{2} x_2^{-1/2} y(x_2) \\ &\geq x_2^{1/2} y'(x_2) - \frac{1}{2} M^{1/2} \\ &\geq (B^2 - KM(\frac{1}{4} + c))^{1/2} - \frac{1}{2} M^{1/2} > B. \end{aligned}$$

If such a solution $y(x)$ is non-oscillatory then the above clearly contradicts (8); hence $y(x)$ is an oscillatory solution of (1).

3. NONOSCILLATION THEOREM.

We shall now prove a result which is the desired counterpart of Theorem C for equation (2). The following result is also a generalization of Theorem D.

THEOREM 2. Let $F(t,x)$ and $G(t,x)$ be given as before. Suppose that there exists a constant $x_0 > 3$, such that for $x \geq x_0$, we have (i) $F(t_2,x) \leq F(t_1,x)$, $t_2 \leq t_1$, and (ii) for every $\alpha > 0$, the function $x \log x G(\alpha x \log x, x)$ is non-increasing in x . Then equation (1) is non-oscillatory.

Before we prove Theorem 2, we need two preliminary lemmas which also seem to have independent interest in themselves. In the following, we restrict the independent variable x to the half open interval $[x_0, \infty)$ without further mention.

LEMMA 4. Let a_1 be a zero of $y(x)$ satisfying (1), and $a_2 > a_1$ such that $y(a_2)y'(a_2) = 0$. Then,

$$(17) \quad a_2 \log a_2 y'^2(a_2) - a_1 \log a_1 y'^2(a_1) \leq - \int_{a_1}^{a_2} \frac{y(x)y'(x)}{x} dx.$$

PROOF. Denote $Q(t,x) = G(t,x) + tF(t,x) + xG_x(t,x)$ as before and observe that

$$(18) \quad (\log x)Q(t,x) + G(t,x) + tF(t,x) \\ = \frac{\partial}{\partial \sigma} \left(\sigma \log \sigma G\left(\frac{t\sigma \log \sigma}{x \log x}, \sigma\right) \right) \Bigg|_{\sigma = x},$$

for arbitrary $t, x > 0$. Thus in view of the non-decreasing character of $(x \log x)G(\alpha x \log x, x)$ for $\alpha > 0$, the left hand side of (18) is non-positive. Let $y(x)$ be a non-trivial solution of (1), and define $\Psi(x)$ by

$$\Psi(x) = x(y'^2(x) + G(y^2(x), x)) - y(x)y'(x)$$

as in Lemma 3. An easy computation shows

$$(19) \quad \frac{d}{dx}(\log x \Psi(x)) = \log x Q(y^2(x), x) + G(y^2(x), x) \\ + y'^2(x) - \frac{y(x)y'(x)}{x}.$$

Integrating (19) between a_1 and a_2 and using the relation

$$\int_{a_1}^{a_2} y'^2(x) dx = \int_{a_1}^{a_2} y^2(x) F(y^2(x), x) dx,$$

we find

$$\int_{a_1}^{a_2} \frac{d}{dx} (\log x \Psi(x)) dx = \int_{a_1}^{a_2} [\log x Q(y^2, x) + G(y^2, x) + y^2 F(y^2, x)] dx \\ - \int_{a_1}^{a_2} \frac{y(x)y'(x)}{x} dx,$$

from which (17) readily follows.

Lemma 5. Let a be any zero of y(x) satisfying (1) and b be the first zero of y'(x) to the right of a. Then there exists a constant B_0 , depending only on the first zero a_0 of y(x) such that

$$(20) \quad \int_a^b y'^2(x) dx \leq B_0.$$

Proof. With no loss of generality, we may assume $y(x) > 0$ on (a, b) . For $x \in (a, b)$, we have

$$((x - a)y' - y)' = (x - a)y'' = -(x - a)yF(y^2, x) < 0$$

hence $y'(x)(x - a) \leq y(x)$. Using this estimate, we easily verify the following inequalities:

$$(21) \quad \int_a^b y'^2(x) dx = \int_a^b \frac{x-a+a}{x} y'^2(x) dx \\ = \int_a^b \frac{y(x)y'(x)}{x} dx + a \int_a^b \frac{y'^2(x)}{x} dx,$$

$$(22) \quad \int_a^b \frac{y'^2(x)}{x} dx = \int_a^b \frac{x-a+a}{x^2} y'^2(x) dx \\ \leq \int_a^b \frac{y(x)y'(x)}{x^2} dx + a \int_a^b \frac{y'^2(x)}{x^2} dx \\ \leq \frac{1}{a} \int_a^b \frac{y(x)y'(x)}{x} dx + a \int_a^b \frac{y'^2(x)}{x^2} dx.$$

Combining (21) and (22), and using the fact that $y'(x) < y'(a)$ for $x \in (a, b)$, we obtain

$$(23) \quad \int_a^b y'^2(x) dx \leq 2 \int_a^b \frac{y(x)y'(x)}{x} dx + ay'^2(a).$$

Using (17) in (23), we have

$$(24) \quad \int_a^b y'^2(x) dx \leq 2a_0 \log a_0 y'^2(a_0) + \frac{a_0 \log a_0 y'^2(a_0)}{\log a} \\ \leq 3a_0 \log a_0 y'^2(a_0) = B_0.$$

Proof of Theorem 2. Assume that the assertion is false, hence, there exists an oscillatory solution $y(x)$ of (1). Let $a_0, a_1, a_2, \dots, a_n, \dots$ be consecutive zeros of $y(x)$ and $b_0, b_1, b_2, \dots, b_n, \dots$ be zeros of $y'(x)$ with $a_{k-1} < b_{k-1} < a_k$, $k = 1, 2, 3, \dots$. Define a sequence of positive numbers $\{\mu_k\}$ by

$$(25) \quad \mu_k = \int_{a_k}^{b_k} y'^2(x) dx = \int_{a_k}^{b_k} y^2(x) F(y^2(x), x) dx.$$

For $x \in [a_k, b_k]$, we have by Schwarz's inequality

$$(26) \quad y^2(x) \leq (x - a_k) \int_{a_k}^x y'^2(x) dx \leq \mu_k x.$$

Using (26) in (25) and the non-decreasing character of $F(t, x)$ with respect to t , we have

$$(27) \quad 1 \leq \int_{a_k}^{b_k} xF(\mu_k x, x) dx \leq \int_{a_k}^{b_k} xF(B_0 x, x) dx,$$

where B_0 is given by (20). Again by assumption (i), we know

that $G(t, x)$ is convex in t . Hence,

$$(28) \quad G(\beta x \log x, x) - G(\alpha x, x) \geq (\beta \log x - \alpha) x F(\alpha x, x),$$

provided $\beta \log x - \alpha \geq 0$. Let $\beta = \alpha(\log x_0)^{-1}$, then (28)

may be rewritten as

$$(29) \quad G\left(\frac{\alpha x \log x}{\log x_0}, x\right) \geq \left(\alpha \frac{\log x}{\log x_0} - \alpha\right) x F(\alpha x, x).$$

Choose $x_1 \geq x_0$ so that $\log x \geq 2 \log x_0$ for $x \geq x_1$. Now

(29) becomes for $x \geq x_1$

$$(30) \quad G\left(\frac{\alpha x \log x}{\log x_0}, x\right) \geq \frac{\alpha \log x}{2 \log x_0} x F(\alpha x, x).$$

Pick k_0 so that $a_k \geq x_1$ for $k \geq k_0$. Using (30) in (27),

we observe

$$1 \leq \int_{a_k}^{b_k} x F(B_0 x, x) dx \leq \int_{a_k}^{b_k} \frac{2 \log x_0}{B_0 \log x} G(B' x \log x, x) dx$$

$$(31) \quad = \frac{2}{B'} \int_{a_k}^{b_k} \frac{1}{x(\log x)^2} x \log x G(B' x \log x, x) dx,$$

where $B' = B(\log x_0)^{-1}$. Denote $G_0 = a_{k_0} \log a_{k_0} G(B' a_{k_0} \log a_{k_0}, a_{k_0})$.

Using assumption (ii), we obtain from (31) the following estimate

valid for $k \geq k_0$,

$$(32) \quad 1 \leq \frac{2G_0}{B_1} \int_{a_k}^{b_k} \frac{dx}{x(\log x)^2} \leq \frac{2G_0}{B' \log a_k}.$$

Letting $k \rightarrow \infty$ in (32), we obtain the desired contradiction.

4. DISCUSSION.

Theorem 1 generalizes the result of Jasný and Urzweil, i.e. Theorem A, in several directions. First, as remarked before, the function $F(t,x)$ is not necessarily of the form of a product of two functions each of t and x alone. Next, the usual assumption on $F(t,x)$ that it be non-decreasing in t is weakened to condition (5) which admits a much larger class of functions. Finally, we would like to point out that in contrast to Theorem A which is strictly a nonlinear result, our Theorem 1 also covers the linear case as well. Consider equation (2) with $\gamma = 1$. Assumptions (i) and (ii) thus reduce to: there exist constants $x_0, c > 0$ such that

$$(33) \quad x^2 p(x) \geq \frac{1}{4} + c, \quad x \geq x_0$$

and that $x^2 p(x)$ is non-decreasing in x . We note that the second condition may be waived in general. Consider a function $\bar{p}(x)$ defined for each $x \geq x_0$ by $x^2 \bar{p}(x) = \inf_{x \leq s < \infty} s^2 p(s) \geq \frac{1}{4} + c$. Clearly $p(x) \geq \bar{p}(x)$ and $x^2 \bar{p}(x)$ is non-decreasing in x .

Applying our result to the linear equation $y'' + p(x)y = 0$ and making use of the Sturm Comparison Theorem, we obtain that the linear equation $y'' + \bar{p}(x)y = 0$ is oscillatory. In this way, Theorem 1 recovers the well-known result of Kneser [6] for the linear case, although it should be mentioned that in the proof presented here we have already used this linear result of Kneser, Cf. Lemma 1. However, alternative proof may be devised so as to

avoid the explicit use of Kneser's result. Such a proof necessarily involves longer arguments.

It is a simple matter to see that Theorem 2 includes the two results of Nehari [11] as special cases. In contrast to Theorem 1 whose proof is significantly different from that of Theorem A, here we make use of some of the techniques developed in [11] in establishing Theorems C and D. We remark that in both Theorems B and C, Kiguradze [5] and Nehari [11] stated their results only for equation (2) with $\gamma > 1$, (in fact they assume $\gamma = 2n - 1$, where n is a positive integer > 1). It is clear from Theorem 2 or a simple application of Sturm's Comparison Theorem that Theorems B and C remain valid for equation (2) in case $\gamma = 1$. Indeed, the proofs of Kiguradze [7] and Nehari [11] made no explicit use of the fact that $\gamma > 1$.

It should be mentioned that the proofs of both Theorems 1 and 2 can be made more transparent by transforming into an auxiliary equation. For example in Theorem 1, let $t = \log x$ and $x^{-1/2}y(x) = u(t)$, then equation (1) is transformed into

$$(34) \quad \frac{d^2u}{dt^2} + u(e^{2t}F(e^t u^2, e^t) - \frac{1}{4}) = 0.$$

Clearly equation (1) possesses an oscillatory solution if and only if equation (34) does. Since this approach does not shorten the presentation, we choose to use the present analysis in order to conform with Nehari's work. A similar comment applies to Theorem 2. Here we use the change of variables: $t = x \log x$ and $(x \log x)^{1/2}y(x) = u(t)$.

Previously it was tacitly assumed that every local solution of (1) can be extended to $[0, \infty)$. Actually this assumption can be omitted without essential modification of the statements or proofs of Theorems 1 and 2, provided that we understand by a solution of (1) a C^2 solution defined on a right maximal interval of existence $[a_1, \omega)$, $a_1 \geq 0$, $\omega \leq +\infty$. We shall call such a solution oscillatory if it has infinitely many zeros in $[x_1, \omega)$; if we adopt this definition then every solution y having a finite right maximal interval of existence is oscillatory, see for example [2]. Moreover, if $a_1 < a_2 < \dots < \omega < \infty$, are the zeros of the solution y then $\lim_{n \rightarrow \infty} |y'(a_n)| = \infty$, and also $\limsup_{x \rightarrow \omega} |y(x)| = \infty$.

For Theorem 2 the assumption on the continuability of solutions is superfluous, since for a given solution $y(x)$, $y'(x)$ must remain bounded on its maximal interval of existence because of (17). In fact, Nehari [11] has already noted that every non-oscillatory solution is ipso facto continuable throughout (a, ∞) . We also note that the continuability assumption is superfluous in Theorem A, although the proofs of Jasny [4] and Kurzweil [7] rely on Atkinson's erroneous statement [1;p.643] that all solutions of (2) can be extended indefinitely when p is continuous and positive. A similar comment applies to the result of Kiguradze. We note that in these cases the monotonicity of $x^\beta p(x)$, $\beta > 0$, implies that $p(x)$ is locally of bounded variation, and hence the continuability of solutions follows from a result in [2].

We would like to point out that results in this paper may be stated in term of the more general equation:

$$y'' + f(y,x) = 0,$$

where $f(y,x)$ satisfies a set of conditions similar to those we impose on the function $F(t,x)$. Such a generalization does not seem to add much to the present knowledge of the problem, we content ourselves with just a mention of such a possibility.

Finally, we note that in contrast to results which guarantee oscillation of all solution or the existence of a non-oscillatory solution, our conditions on the growth of the function $G(t,x)$ are not in terms of convergence or divergence of a certain integral. In this sense, our results are not totally satisfactory. For example, Atkinson [1] has established a necessary and sufficient condition for equation (2), when $\gamma > 1$, to be oscillatory, namely

$$\int_0^{\infty} x p(x) dx = \infty.$$

(Extension of Atkinson's result to more general equations similar to (1) have been given by Macki and Wong [8].) It would be tempting to conjecture that a necessary and sufficient condition for equation (2) to be non-oscillatory is

$$\int_0^{\infty} x^{\gamma+1/2} p(x) dx < \infty.$$

Thus far, the validity of such a conjecture remains unsettled.

REFERENCES

- [1] F. V. Atkinson, 'On second order nonlinear oscillation', Pacific J. Math., 5 (1955), 643-647.
- [2] C. V. Coffman and D. F. Ullrich, 'On the continuation of solutions of a certain nonlinear differential equation', Monatsh Math., 71 (1967), 385-392.
- [3] S. P. Hastings, 'Boundary value problems in one differential equation with a discontinuity', J. Diff. Eqn., 1 (1965), 346-369.
- [4] M. Jasný, 'On the existence of an oscillatory solution of the nonlinear differential equation of the second order $y'' + f(x)y^{2n-1} = 0$, $f(x) > 0$ ', Časopis Pěstování Matematiky, 85 (1960), 78-83. (Russian) MR 26, No. 408.
- [5] I. T. Kiguradze, 'On conditions for oscillation of solutions of the equation $u'' + a(t)|u|^n \operatorname{sgn} u = 0$ ', Časopis Pěstování Matematiky, 87 (1962), 492-495. (Russian) MR 31, No. 6026.
- [6] A. Kneser, 'Untersuchungen über die reellen Nullstellen der Integrale linearer Differential Gleichungen', Math. Annalen, 42 (1893), 409-435.
- [7] J. Kurzweil, 'A note on oscillatory solution of equation $y'' + f(x)y^{2n-1} = 0$ ', Časopis Pěstování Matematiky, 85 (1960), 357-358. (Russian) MR 23, No. A3322.

- [8] J. W. Macki and J. S. W. Wong, 'Oscillation of solutions to second order nonlinear differential equations', Pacific J. Math., 24 (1968), 111-117.
- [9] R. A. Moore and Z. Nehari, 'Nonoscillation theorems for a class of nonlinear differential equation', Trans. Amer. Math. Soc., 93 (1959), 30-52.
- [10] Z. Nehari, 'On a class of nonlinear second order differential equations', Trans. Amer. Math. Soc., 95 (1960), 101-123.
- [11] Z. Nehari, 'A nonlinear oscillation problem', J. Differential Eqn., to appear; also Carnegie-Mellon University Report, 67-36.
- [12] J. S. W. Wong, 'On second order nonlinear oscillation', Funkcial. Ekvac., to appear; also MRC technical report, no. 836.

Carnegie-Mellon University
Pittsburgh, Pa. 15213