

**NOTICE WARNING CONCERNING COPYRIGHT RESTRICTIONS:**  
The copyright law of the United States (title 17, U.S. Code) governs the making of photocopies or other reproductions of copyrighted material. Any copying of this document without permission of its author may be prohibited by law.

ON THE THERMODYNAMICS OF CHEMICALLY  
REACTING FLUID MIXTURES

by

Morton E. Gurtin

Research Report 71-23

April, 1971

/nlc  
16 April 1971

University Libraries  
Carnegie Mellon University  
Pittsburg, PA 15213-3890

HUNT LIBRARY  
CARNEGIE-MELLON UNIVERSITY

JUN 3 '71

ABSTRACT

On the Thermodynamics of Chemically Reacting Fluid

Mixtures

by

Morton E. Gurtin

This paper develops a thermodynamical theory of chemically reacting, but inviscid, fluid mixtures. Restrictions placed on the constitutive equations by the second law are found. In particular, it is shown that the stress-diffusion matrix is symmetric.

# ON THE THERMODYNAMICS OF CHEMICALLY REACTING FLUID

## MIXTURES

by

Morton E. Gurtin

### Introduction.

After a long history of special theories of diffusion and chemical reactions, Truesdell [1957] established the basic thermomechanical balance laws for mixtures,<sup>1</sup> and Bowen,<sup>2</sup> Müller [1968], and Truesdell [1968] proposed workable forms for the second law. Using this framework together with the Coleman-Noll [1963] interpretation of the second law, Müller [1968]<sup>3</sup> developed a theory of non-reacting fluid mixtures,<sup>4</sup> and Bowen [1969] established a general theory for mixtures of chemically reacting elastic materials.<sup>5</sup>

In this paper, I discuss chemically reacting, but inviscid, fluid mixtures. Among the new results I establish

---

<sup>1</sup>See also Nachbar, Williams, and Penner [1959], Kelly [1964], and Green and Naghdi [1969].

<sup>2</sup>See Truesdell [1969] (Footnote on p. 88).

<sup>3</sup>Thermodynamic theories were put forth before 1968 by Eringen and Ingram [1965], Green and Naghdi [1965,1967], Crochet and Naghdi [1966,1967], Green and Steel [1966], Mills [1966,1967], Bowen [1967], Ingram and Eringen [1967], and Atkin [1967]. As Müller [1968] and Truesdell [1969] have pointed out, all of the above theories suffer serious defects.

<sup>4</sup>See also Dunwoody and Müller [1968], Green and Naghdi [1968], Doria [1969], Graine, Green, and Naghdi [1970], and Dunwoody [1970].

<sup>5</sup>See also Bowen and Wiese [1969], Green and Naghdi [1969], Bowen and Garcia [1970].

are the following:

(i) The stress-diffusion matrix is symmetric. The stress diffusion matrix is that matrix whose  $(\alpha, \beta)$  entry is the derivative of the stress vector of constituent  $\alpha$  with respect to the velocity of constituent  $\beta$ .

(ii) When the diffusion velocity  $\tilde{u}_\alpha$  of constituent  $\alpha$  is small, the diffusive energy flux is approximated by its classical counterpart  $\sum_{\alpha} \mu_\alpha \rho_\alpha \tilde{u}_\alpha$ , where  $\mu_\alpha$  is the chemical potential and  $\rho_\alpha$  the density of  $\alpha$ .

(iii) The elasticity matrix corresponding to a strong equilibrium state is symmetric. This matrix plays a crucial role in the linearized theory and in wave propagation studies.

(iv) When the diffusion velocities and the gradients of density and temperature are small, and when the underlying state is one of strong equilibrium, to within second order terms the heat flux depends only on the temperature gradient and the diffusion velocities, and the mass supply (due to chemical reactions) depends only on the chemical potentials.

For convenience, I omit smoothness hypotheses; it will be clear from the context what these ought to be.

1. Preliminary Definitions. Terminology.

Throughout this paper  $\mathbb{R}$ ,  $\mathbb{R}^+$ , and  $\mathcal{V}$  denote, respectively, the reals, the strictly positive reals, and the vector space associated with three-dimensional euclidean space. Given a (second-order) tensor  $\underline{S}$ ,  $\underline{S}^T$  designates its transpose and  $\text{sym}\underline{S} = \frac{1}{2}(\underline{S} + \underline{S}^T)$  its symmetric part. We denote the tensor product of two vectors  $\underline{u}$  and  $\underline{v}$  by  $\underline{u} \otimes \underline{v}$ .

The mixtures we study will have  $N$  constituents; for convenience, we identify the set of constituents with the set  $\{1, \dots, N\}$ . Further, the letters  $\alpha$  and  $\beta$  will always denote constituents, and we write  $\sum_{\alpha} f_{\alpha}$  and  $\sum_{\alpha} f_{\alpha}$  for  $\sum_{\alpha=1}^N f_{\alpha}$ .

We use the following notation:

$\rho_{\alpha}$  is the mass density of constituent  $\alpha$ ,

$\underline{v}_{\alpha}$  is the velocity of  $\alpha$ .

In addition, we write

$$\rho = \sum \rho_{\alpha}, \quad c_{\alpha} = \frac{\rho_{\alpha}}{\rho}, \quad (1.1)$$

$$\underline{v} = \sum c_{\alpha} \underline{v}_{\alpha}, \quad \underline{u}_{\alpha} = \underline{v}_{\alpha} - \underline{v};$$

$\rho$  is the total mass density,  $c_{\alpha}$  the concentration of  $\alpha$ ,  $\underline{v}$  the velocity of the mixture, and  $\underline{u}_{\alpha}$  the diffusion velocity of  $\alpha$ . Clearly,

$$\sum c_{\alpha} = 1, \quad \sum c_{\alpha} \underline{u}_{\alpha} = 0. \quad (1.2)$$

Since we are dealing with a fluid mixture, it is more convenient to work with the spatial description of any given motion. Thus a motion is specified by prescribing  $\underline{v}_\alpha(\underline{x}, t)$  and  $\rho_\alpha(\underline{x}, t)$  as functions of spatial position  $\underline{x} \in \mathcal{R}$  and time  $t$ ; here  $\mathcal{R}$  denotes a fixed region of space occupied by the mixture during some time interval.

We write  $\text{grad}$  and  $\text{div}$  for the spatial gradient and spatial divergence, i.e., the gradient and divergence with respect to  $\underline{x}$  holding  $t$  fixed. Given a scalar field  $f(\underline{x}, t)$  and a vector field  $\underline{f}(\underline{x}, t)$ , we define the material time derivatives  $\dot{f}(\underline{x}, t)$  and  $\dot{\underline{f}}(\underline{x}, t)$  through

$$\dot{f} = \frac{\partial f}{\partial t} + (\text{grad}f) \cdot \underline{v}, \quad \dot{\underline{f}} = \frac{\partial \underline{f}}{\partial t} + (\text{grad}\underline{f}) \underline{v}; \quad (1.3)$$

we then have the identity

$$\text{grad}\dot{f} = \overline{\text{grad}\dot{f}} + (\text{grad}\underline{v})^T \text{grad}f. \quad (1.4)$$

## 2. Basic Laws.

For the basic laws of our theory we postulate the following:

### balance of mass for each constituent

$$\dot{\rho}_\alpha + \rho_\alpha \operatorname{div}_{\sim\alpha} v_{\sim\alpha} + u_{\sim\alpha} \cdot \operatorname{grad} \rho_\alpha = \rho_\alpha c_\alpha^+, \quad (2.1)$$

### balance of momentum for each constituent

$$\rho_\alpha \dot{v}_{\sim\alpha} + \rho_\alpha (\operatorname{grad}_{\sim\alpha} v_{\sim\alpha}) u_{\sim\alpha} = \operatorname{div}_{\sim\alpha} T_{\sim\alpha} + \rho_\alpha b_{\sim\alpha} + \rho_\alpha t_\alpha^+, \quad (2.2)$$

### balance of energy for the mixture<sup>1</sup>

$$\rho \dot{e} = -\operatorname{div}(q + j) + \sum (\overset{\sim}{T}_\alpha - \rho_\alpha u_{\sim\alpha} \otimes u_{\sim\alpha}) \cdot \operatorname{grad}_{\sim\alpha} v_{\sim\alpha} + \sum \rho_\alpha b_{\sim\alpha} \cdot u_{\sim\alpha} + \rho r, \quad (2.3)$$

### growth of entropy for the mixture

$$\rho \dot{s} \geq -\operatorname{div}\left(\frac{q}{\theta}\right) + \frac{\rho r}{\theta}. \quad (2.4)$$

<sup>1</sup>In Müller's [1968] theory the term  $j$  is not present in the energy equation and the term  $q/\theta$  in the entropy inequality is replaced by an arbitrary entropy flux  $\underline{\xi}$ . To see that there is no contradiction between Müller's theory and the one presented here, let  $q'$  denote the heat flux in the former. If we define a new "heat flux"  $q$  by  $q = \theta \underline{\xi}$  and a diffusive energy flux  $j$  by  $j = q' - q$ , then Müller's equations (2.16), (3.4) reduce to my (2.3), (2.4). The difference is simply a matter of taste. Indeed, within the context of the classical theory of mixtures I would view the quantity  $\sum \mu_\alpha \rho_\alpha u_{\sim\alpha}$ , where  $\mu_\alpha$  is the chemical potential of  $\alpha$ , as a flux energy (as does Eckart [1940]), while Müller would view  $-(\sum \mu_\alpha \rho_\alpha u_{\sim\alpha})/\theta$  as a flux of entropy (as do Meixner and Reik [1959] and de Groot and Mazur [1962]). I take the point of view that the "heat flux" is that vector field which when divided by the temperature gives the entropy flux.

The theory presented here falls within Bowen's [1969] framework provided we take  $j$  to be a certain specified linear function of the diffusion velocities.



Here

- $\underline{T}_{\alpha}$  is the stress tensor for  $\alpha$ ,  
 $\underline{b}_{\alpha}$  is the body force for  $\alpha$ ,  
 $\underline{l}_{\alpha}^{+}$  is the momentum supply for  $\alpha$ ,  
 $e$  is the internal energy,  
 $\underline{q}$  is the heat flux,  
 $\underline{j}$  is the diffusive energy flux,  
 $r$  is the heat supply,  
 $s$  is the entropy,  
 $\theta$  is the temperature.

We also postulate that

$$\Sigma c_{\alpha}^{+} = 0, \quad \Sigma (c_{\alpha\sim\alpha}^{+} l_{\alpha}^{+} + c_{\alpha\sim\alpha}^{+} u_{\alpha}) = 0; \quad (2.5)$$

then (1.1), (2.1), and (2.2) imply the laws of mass and momentum balance for the mixture:

$$\begin{aligned} \dot{\rho} + \rho \operatorname{div} \underline{v} &= 0, \\ \rho \dot{\underline{v}} &= \operatorname{div} (\underline{T} - \Sigma \rho_{\alpha\sim\alpha} u_{\alpha} \otimes u_{\alpha}) + \Sigma \rho_{\alpha\sim\alpha} \underline{b}_{\alpha}, \end{aligned} \quad (2.6)$$

where

$$\underline{T} = \Sigma \underline{T}_{\alpha} \quad (2.7)$$

is (the inner part of) the total stress.

If we use (2.2) to eliminate  $\underline{b}_{\alpha}$  in (2.3), we find, with the aid of (1.1), (1.2), (2.1), and (2.6)<sub>1</sub>,

$$\rho \dot{e}_I = -\text{div}(\underline{q} + \underline{k}) + \sum_{\alpha} \underline{T}_{\alpha} \cdot \text{grad} \underline{v}_{\alpha} - \sum_{\alpha} \underline{n}_{\alpha} \cdot \underline{u}_{\alpha} + \rho r, \quad (2.8)$$

where

$$e_I = e - \frac{1}{2} \sum_{\alpha} c_{\alpha} u_{\alpha}^2 \quad (2.9)$$

is the inner part of the internal energy and

$$\underline{k} = \underline{j} + \sum_{\alpha} (\underline{T}_{\alpha}^T - \frac{1}{2} \rho_{\alpha} u_{\alpha}^2 \underline{1}) \underline{u}_{\alpha}, \quad \underline{n}_{\alpha} = \rho_{\alpha} \underline{l}_{\alpha}^+ + \frac{1}{2} \rho c_{\alpha}^+ \underline{u}_{\alpha}. \quad (2.10)$$

If we define (the inner part of) the free energy by

$$\psi = e_I - \theta s, \quad (2.11)$$

then (2.4) and (2.8) yield the reduced dissipation inequality:

$$\rho(\dot{\psi} + s\dot{\theta}) - \sum_{\alpha} \underline{T}_{\alpha} \cdot \text{grad} \underline{v}_{\alpha} + \text{div} \underline{k} + \sum_{\alpha} \underline{n}_{\alpha} \cdot \underline{u}_{\alpha} + \frac{1}{\theta} \underline{q} \cdot \text{grad} \theta \leq 0. \quad (2.12)$$

### 3. Constitutive Assumptions.

We consider a mixture defined by constitutive equations in which:

$$\psi, \underline{T}_\alpha, s, \underline{q}, \underline{j}, \underline{l}_\alpha^+, c_\alpha^+ \quad \text{are functions of} \quad (\underline{\rho}, \theta, \text{grad} \underline{\rho}, \text{grad} \theta, \underline{\tilde{v}}), \quad (3.1)$$

where, for convenience, we have used the notation

$$\underline{\rho} = (\rho_1, \dots, \rho_N), \quad \text{grad} \underline{\rho} = (\text{grad} \rho_1, \dots, \text{grad} \rho_N), \quad \underline{\tilde{v}} = (\underline{v}_1, \dots, \underline{v}_N). \quad (3.2)$$

We assume that the response functions are isotropic and that  $c_\alpha^+$  and  $\underline{l}_\alpha^+$  are consistent with (2.5). Further, in order to make our theory consistent with material frame indifference, we require that

$$f(\underline{\rho}, \theta, \text{grad} \underline{\rho}, \text{grad} \theta, \underline{\tilde{v}}) = f(\underline{\rho}, \theta, \text{grad} \underline{\rho}, \text{grad} \theta, \underline{\tilde{v}} + \underline{\tilde{a}}) \quad (3.3)$$

whenever  $f$  is one of the response functions in (3.1) and  $\underline{\tilde{a}} = (\underline{a}_1, \dots, \underline{a}_N)$ . Clearly, (3.3) implies that

$$\sum \frac{\partial f}{\partial \underline{v}_{\tilde{\alpha}}} = 0 \quad (3.4)$$

and that

$$\underline{f}(\underline{\rho}, \theta, \text{grad} \underline{\rho}, \text{grad} \theta, \underline{\tilde{v}}) = \underline{f}(\underline{\rho}, \theta, \text{grad} \underline{\rho}, \text{grad} \theta, \underline{\tilde{u}}), \quad (3.5)$$

where

$$\underline{\tilde{u}} = (\underline{u}_1, \dots, \underline{u}_N) \quad (3.6)$$

with  $\underline{u}_\alpha$  the diffusion velocity (1.1)<sub>4</sub>.

We also assume that:<sup>1</sup>

(A) Given a point  $\underline{x}_0 \in \mathbb{R}^N$ , a neighborhood  $\mathcal{P}_0$  of  $\underline{x}_0$ , an initial density distribution  $\vec{\rho}_0 : \mathcal{P}_0 \rightarrow (\mathbb{R}^+)^N$ , a time interval  $[0, t_0)$ , a temperature field  $\theta : \mathcal{P}_0 \times [0, t_0) \rightarrow \mathbb{R}^+$ , and a velocity distribution  $\vec{v} : \mathcal{P}_0 \times [0, t_0) \rightarrow \mathcal{V}^N$ ; there exists a  $\mathcal{P} \subset \mathcal{P}_0$  with  $\underline{x}_0 \in \mathcal{P}$ , a  $\tau \in (0, t_0)$ , and a solution  $\vec{\rho} : \mathcal{P} \times [0, \tau) \rightarrow (\mathbb{R}^+)^N$  of (2.1), i.e. of

$$\dot{\rho}_\alpha + \rho_\alpha \operatorname{div} \underline{v}_\alpha + \underline{u}_\alpha(\vec{\rho}, \vec{v}) \cdot \operatorname{grad} \rho_\alpha = \rho(\vec{\rho}) c_\alpha^+(\vec{\rho}, \theta, \operatorname{grad} \vec{\rho}, \operatorname{grad} \theta, \vec{v}) \quad (3.7)$$

on  $\mathcal{P} \times (0, \tau)$ , such that

$$\vec{\rho}(\underline{x}, 0) = \vec{\rho}_0 \quad \text{for } \underline{x} \in \mathcal{P}. \quad (3.8)$$

Of course, in (3.7)  $\underline{u}_\alpha(\vec{\rho}, \vec{v})$  and  $\rho(\vec{\rho})$  are given by (1.1).

An array  $(\vec{\rho}, \theta, \vec{v}, \psi, \vec{T}, s, \underline{q}, j, \vec{l}^+, \vec{c}^+)$  of fields on  $\mathcal{P} \times [0, \tau)$  (with values in appropriate spaces) will be called a constitutive process if it is consistent with the constitutive assumption (3.1) and balance of mass (2.1). Here  $\vec{T} = (T_1, \dots, T_N)$ ,  $\vec{l}^+ = (l_1^+, \dots, l_N^+)$ , and  $\vec{c}^+ = (c_1^+, \dots, c_N^+)$ . Assumption (A) insures that given  $\underline{x}_0, \vec{\rho}_0, \vec{v}$ , and  $\theta$  there exists an associated constitutive process in some neighborhood of  $\underline{x}_0$ .

---

<sup>1</sup>Bowen [1969] was the first to notice that an assumption of this type is necessary when chemical reactions are present. See also Coleman and Gurtin [1967] who utilize an assumption of this type in a slightly different context.

Note that by (1.1), (2.10)<sub>1</sub>, and (3.1),

$$\underline{k} = \underline{k}(\vec{\rho}, \theta, \text{grad}\vec{\rho}, \text{grad}\theta, \vec{v}).$$

For convenience, we introduce the following notation:

$$\text{div}_{\underline{0}\underline{\sim}} \underline{k} = \Sigma \frac{\partial \underline{k}}{\partial \underline{\rho}_\alpha} \cdot \text{grad} \underline{\rho}_\alpha + \frac{\partial \underline{k}}{\partial \theta} \cdot \text{grad} \theta, \quad (3.9)$$

$$\text{div}_{\underline{1}\underline{\sim}} \underline{k} = \Sigma \frac{\partial \underline{k}}{\partial (\text{grad} \underline{\rho}_\alpha)} \cdot \text{grad}^2 \underline{\rho}_\alpha + \frac{\partial \underline{k}}{\partial (\text{grad} \theta)} \cdot \text{grad}^2 \theta;$$

thus

$$\text{div} \underline{k} = \text{div}_{\underline{0}\underline{\sim}} \underline{k} + \text{div}_{\underline{1}\underline{\sim}} \underline{k} + \Sigma \left( \frac{\partial \underline{k}}{\partial \underline{v}_\alpha} \right)^T \cdot \text{grad} \underline{v}_\alpha. \quad (3.10)$$

#### 4. Consequences of the Second Law.

Given a constitutive process<sup>1</sup> (2.2) and (2.3) can be used to determine the body forces  $b_{\alpha}$  and the heat supply  $r$  necessary to sustain the process. On the other hand, the inequality (2.4) -- or equivalently (2.12) -- will be satisfied in every process if and only if certain restrictions are placed on the response functions. The next theorem lists these restrictions.<sup>2</sup>

---

<sup>1</sup>Note that, by definition, the mass balance relations (2.1) are satisfied by every constitutive process.

<sup>2</sup>Cf. Müller [1968, §5], Doria [1969, §5], Bowen and Wiese [1969, §5], Bowen [1969, §6].

Theorem 4.1. A necessary and sufficient condition that every constitutive process obey the reduced dissipation inequality (2.12) is that the following four statements be true:

(i) The total stress is a pressure:

$$\underline{T} = - p \underline{1}. \quad (4.1)$$

(ii)  $\psi$ ,  $p$ , and  $s$  are independent of  $\text{grad} \vec{\rho}$ ,  $\text{grad} \theta$ , and  $\vec{v}$ :

$$\psi = \psi(\vec{\rho}, \theta), \quad p = p(\vec{\rho}, \theta), \quad s = s(\vec{\rho}, \theta); \quad (4.2)$$

moreover,

$$p = \rho \Sigma \rho_{\alpha} \frac{\partial \psi}{\partial \rho_{\alpha}}, \quad s = - \frac{\partial \psi}{\partial \theta}. \quad (4.3)$$

(iii) The constituent stress  $\underline{T}_{\alpha}$  is given by

$$\underline{T}_{\alpha} = \left( \frac{\partial \underline{k}}{\partial \underline{v}_{\alpha}} \right)^T - \rho \rho_{\alpha} \frac{\partial \psi}{\partial \rho_{\alpha}} \underline{1}. \quad (4.4)$$

(iv) In every constitutive process

$$\text{div}_{\underline{0}} \underline{k} + \rho \Sigma \frac{\partial \psi}{\partial \rho_{\alpha}} [\rho c_{\alpha}^{+} - \underline{u}_{\alpha} \cdot \text{grad} \rho_{\alpha}] + \Sigma \underline{n}_{\alpha} \cdot \underline{u}_{\alpha} + \frac{1}{\theta^2} \underline{g} \cdot \text{grad} \theta \leq 0, \quad (4.5)$$

$$\text{div}_{\underline{1}} \underline{k} = 0.$$

Proof. The proof of sufficiency follows upon direct substitution. To establish the necessity of (i)-(iv) we assume that every constitutive process obeys (2.12), or equivalently, in view of the constitutive assumption (3.1), balance of mass (2.1), and (3.10),

$$\begin{aligned}
& \rho \left( \frac{\partial \psi}{\partial \theta} + s \right) \dot{\theta} - \Sigma \left[ \rho \rho_{\alpha} \frac{\partial \psi}{\partial \rho_{\alpha}} \underline{1} + \underline{T}_{\alpha} - \left( \frac{\partial \underline{k}}{\partial \underline{v}_{\alpha}} \right)^T \right] \cdot \text{grad} \underline{v}_{\alpha} + \\
& + \rho \left[ \Sigma \frac{\partial \psi}{\partial (\text{grad} \rho_{\alpha})} \cdot \overline{\text{grad} \rho_{\alpha}} + \frac{\partial \psi}{\partial (\text{grad} \theta)} \cdot \overline{\text{grad} \theta} + \Sigma \frac{\partial \psi}{\partial \underline{v}_{\alpha}} \cdot \dot{\underline{v}}_{\alpha} \right] + \quad (4.6) \\
& + \rho \Sigma \frac{\partial \psi}{\partial \rho_{\alpha}} \left[ \rho c_{\alpha}^+ - \underline{u} \cdot \text{grad} \rho_{\alpha} \right] + \text{div}_{\underline{0}} \underline{k} + \text{div}_{\underline{1}} \underline{k} + \Sigma \underline{n}_{\alpha} \cdot \underline{u}_{\alpha} + \frac{1}{\theta^2} \underline{g} \cdot \text{grad} \theta \leq 0.
\end{aligned}$$

It follows from Lemma 6.1 that  $\dot{\theta}$ ,  $\overline{\text{grad} \rho_{\alpha}}$ ,  $\overline{\text{grad} \theta}$ ,  $\dot{\underline{v}}_{\alpha}$ , and  $\text{grad} \underline{v}_{\alpha}$  can be specified arbitrarily in (4.6), and this observation yields (4.2)<sub>1,3</sub>, (4.3)<sub>2</sub>, and (4.4). Next, it follows from (2.10) and (3.1) that (3.3) (and hence (3.4)) holds with  $f = \underline{k}$ ; thus, if we sum (4.4) from 1 to  $\alpha$ , we are led to (4.1) and (4.3)<sub>1</sub>. Next,  $\text{div}_{\underline{1}} \underline{k}$  is the only term in (4.6) involving second gradients; thus it follows from (3.9)<sub>2</sub> and Lemma 6.1 that

$$\text{sym} \left( \frac{\partial \underline{k}}{\partial (\text{grad} \rho_{\alpha})} \right) = \text{sym} \left( \frac{\partial \underline{k}}{\partial (\text{grad} \theta)} \right) = \underline{0}, \quad (4.7)$$

which implies (4.5)<sub>2</sub>. Finally, (4.1)-(4.4), (4.5)<sub>2</sub>, and (4.6) yield (4.5)<sub>1</sub>.  $\square$

We assume for the remainder of the paper that (i)-(iv) of Theorem 4.1 hold.

Let  $\underline{n}$  be a unit vector. The quantity

$$\underline{t}_{\alpha}(\underline{n}) = \underline{T}_{\alpha} \underline{n} \quad (4.8)$$

is the stress vector for constituent  $\alpha$  (corresponding to the unit



normal  $\underline{n}$ ). By (3.1)  $\underline{t}_\alpha(\underline{n})$  is a function of  $(\underline{\rho}, \theta, \text{grad}\underline{\rho}, \text{grad}\theta, \underline{v})$ .

We call the matrix  $\left\| \frac{\partial \underline{t}_\alpha(\underline{n})}{\partial \underline{v}_\beta} \right\|$  (with tensor entries) the stress-diffusion matrix.

Theorem 4.2. The stress-diffusion matrix is symmetric;  
i.e.

$$\frac{\partial \underline{t}_\alpha(\underline{n})}{\partial \underline{v}_\beta} = \left( \frac{\partial \underline{t}_\beta(\underline{n})}{\partial \underline{v}_\alpha} \right)^T \quad (4.9)$$

for every unit vector  $\underline{n}$  and all constituents  $\alpha, \beta$ .

Proof. Let  $\kappa = \underline{k} \cdot \underline{n}$ . Then (4.4) implies

$$\underline{t}_\alpha(\underline{n}) = \frac{\partial \kappa}{\partial \underline{v}_\alpha} - \rho \rho_\alpha \frac{\partial \psi}{\partial \rho_\alpha} \underline{n}; \quad (4.10)$$

since the second term (4.10) is independent of  $\underline{v}$ , if we differentiate  $\underline{t}_\alpha(\underline{n})$  with respect to  $\underline{v}_\beta$ , we are led at once to (4.9).  $\square$

We call the quantity

$$\mu_\alpha = \mu_\alpha(\underline{\rho}, \theta) = \rho \left[ \frac{\partial \psi}{\partial \rho_\alpha} - \frac{1}{N} \sum_\beta \frac{\partial \psi}{\partial \rho_\beta} \right] \quad (4.11)$$

the chemical potential of  $\alpha$ ; note that

$$\sum \mu_\alpha = 0. \quad (4.12)$$

In view of (2.5)<sub>1</sub> and (4.11), we can rewrite (4.5) as follows:

$$\operatorname{div}_{\underline{\underline{O}}} \underline{\underline{k}} + \Sigma (\underline{\underline{n}}_{\alpha} - \rho \frac{\partial \psi}{\partial \rho_{\alpha}} \operatorname{grad} \rho_{\alpha}) \cdot \underline{\underline{u}}_{\alpha} + \frac{1}{\theta} \underline{\underline{q}} \cdot \operatorname{grad} \theta + \rho \Sigma \mu_{\alpha} c_{\alpha}^{+} \leq 0. \quad (4.13)$$

Further, by (1.1), (1.2)<sub>2</sub>, (2.1), (2.5)<sub>1</sub>, (2.9), (2.11), (4.1)-(4.3), and (4.11), the energy equation (2.3) takes the form

$$\begin{aligned} \rho (\theta \dot{s} + \frac{1}{2} \Sigma c_{\alpha} u_{\alpha}^2) &= -\operatorname{div}(\underline{\underline{q}} + \underline{\underline{j}} - \Sigma \mu_{\alpha} \rho_{\alpha} \underline{\underline{u}}_{\alpha}) - \Sigma (\rho_{\alpha} \underline{\underline{u}}_{\alpha} \otimes \underline{\underline{u}}_{\alpha}) \cdot \operatorname{grad} \underline{\underline{v}} \\ &+ \Sigma \rho_{\alpha} \underline{\underline{u}}_{\alpha} \cdot (\underline{\underline{b}}_{\alpha} - \operatorname{grad} \mu_{\alpha}) - \rho \Sigma \mu_{\alpha} c_{\alpha}^{+} + pr. \end{aligned} \quad (4.14)$$

In the classical theory of fluid mixtures the diffusive energy flux has the form  $\underline{\underline{j}} = \Sigma \mu_{\alpha} \rho_{\alpha} \underline{\underline{u}}_{\alpha}$ , so that the divergence term in (4.14) reduces to  $-\operatorname{div} \underline{\underline{q}}$ . The next theorem shows that in the general theory the diffusive energy flux is approximated by its classical counterpart.

Theorem 4.3. The diffusive energy flux  $\underline{\underline{j}} = \underline{\underline{j}}(\underline{\underline{\rho}}, \theta, \operatorname{grad} \underline{\underline{\rho}}, \operatorname{grad} \theta, \underline{\underline{v}})$  vanishes when the diffusion velocities vanish. Moreover,

$$\underline{\underline{j}} = \Sigma \mu_{\alpha} \rho_{\alpha} \underline{\underline{u}}_{\alpha} + o(|\underline{\underline{u}}|^2) \quad \text{as} \quad \underline{\underline{u}} \rightarrow \underline{\underline{0}}. \quad (4.15)$$

Proof. Let  $\Omega = (\underline{\underline{\rho}}, \theta, \operatorname{grad} \underline{\underline{\rho}}, \operatorname{grad} \theta)$ . Since the material is isotropic, it follows from (2.10)<sub>1</sub> that  $\underline{\underline{k}} = \underline{\underline{k}}(\Omega, \underline{\underline{v}})$  is an isotropic function. Thus, by (4.7) and Lemma 6.2,

$$\underline{\underline{k}}(\Omega, \underline{\underline{0}}) = \underline{\underline{0}}, \quad (4.16)^1$$

<sup>1</sup>For binary mixtures Doria [1969, Eq. (6.35)] establishes the stronger result that  $\underline{\underline{k}} = \varphi \underline{\underline{u}}$  ( $\underline{\underline{u}} = \underline{\underline{u}}_1 = -\underline{\underline{u}}_2$ ) with  $\varphi = \varphi(\rho_1, \rho_2, \theta, |\underline{\underline{u}}|)$ , which implies, by (4.4) and (4.2), that  $\underline{\underline{T}}_{\alpha}$  is independent of  $\operatorname{grad} \rho$  and  $\operatorname{grad} \theta$ ; unfortunately, these results are restricted to binary mixtures.

and we conclude from (2.10)<sub>1</sub> that

$$\underline{j}(\Omega, \vec{0}) = 0. \quad (4.17)$$

Next, by (1.1)<sub>3,4</sub>,

$$\frac{\partial u_{\beta}}{\partial v_{\alpha}} = (\delta_{\alpha\beta} - c_{\alpha}) \underline{1}; \quad (4.18)$$

thus if we differentiate (2.10)<sub>1</sub> with respect to  $v_{\alpha}$  and use (2.7), (4.1), and (4.4), we arrive at

$$\frac{\partial \underline{j}}{\partial v_{\alpha}} = (\rho \rho_{\alpha} \frac{\partial \psi}{\partial \rho_{\alpha}} - c_{\alpha} p) \underline{1} \quad \text{when} \quad \vec{v} = \vec{0}. \quad (4.19)$$

Therefore

$$\underline{j}(\Omega, \vec{v}) = \Sigma (\rho \rho_{\alpha} \frac{\partial \psi}{\partial \rho_{\alpha}} - c_{\alpha} p) v_{\alpha} + o(|\vec{v}|^2) \quad \text{as} \quad \vec{v} \rightarrow \vec{0}. \quad (4.20)$$

But by (3.5),  $\underline{j}(\Omega, \vec{v}) = \underline{j}(\Omega, \vec{u})$ ; thus, by (1.2)<sub>2</sub>,

$$\underline{j}(\Omega, \vec{u}) = \Sigma \rho \rho_{\alpha} \frac{\partial \psi}{\partial \rho_{\alpha}} u_{\alpha} + o(|\vec{u}|^2) \quad \text{as} \quad \vec{u} \rightarrow \vec{0}; \quad (4.21)$$

and this result, in conjunction with (1.2)<sub>2</sub> and (4.11), implies the desired result (4.15).  $\square$

### 5. Results near Equilibrium.<sup>1</sup>

Let  $(\vec{\rho}_0, \theta_0)$  be given, and, for convenience, let

$$\vec{f}_\alpha = \text{grad} \rho_\alpha, \quad \vec{f} = \text{grad} \vec{\rho}, \quad \vec{g} = \text{grad} \theta, \quad (5.1)$$

$$\mathcal{A} = (\vec{\rho}, \theta, \vec{f}, \vec{g}, \vec{v}), \quad \mathcal{A}_0 = (\vec{\rho}_0, \theta_0, \vec{0}, \vec{0}, \vec{0}).$$

We call  $\mathcal{A}_0$  an equilibrium state provided

$$c_\alpha^+(\mathcal{A}_0) = 0 \quad (5.2)$$

for every constituent  $\alpha$ . Let  $h$  denote the left-hand side of (4.5), or equivalently, of (4.13). It then follows from (3.9)<sub>1</sub>, (3.1), (2.10), and (4.11) that

$$\begin{aligned} h = h(\mathcal{A}) = \Sigma \frac{\partial k}{\partial \rho_\alpha} \cdot \vec{f}_\alpha + \frac{\partial k}{\partial \theta} \cdot \vec{g} + \Sigma (n_\alpha - \rho \frac{\partial \psi}{\partial \rho_\alpha}) \cdot \vec{u}_\alpha + \\ + \frac{1}{\theta} \vec{g} \cdot \vec{g} + \rho \Sigma \mu_\alpha c_\alpha^+, \end{aligned} \quad (5.3)$$

and it is clear from Theorem 4.1 and Lemma 6.1 that

$$h(\mathcal{A}) \leq 0 \quad (5.4)$$

for every  $\mathcal{A}$  in the domain of the response functions.

If  $\mathcal{A}_0$  is an equilibrium state, then (5.2) and (5.3) imply that  $h(\mathcal{A}_0) = 0$ , so that

---

<sup>1</sup>In the absence of chemical reactions ( $c_\alpha^+ \equiv 0$ ) every  $\mathcal{A}_0$  of the form (5.1)<sub>5</sub> is a strong equilibrium state; thus all of the results of this section apply also to diffusion without chemical reactions.

$$h(\mathcal{N}) \text{ is a maximum at } \mathcal{N} = \mathcal{N}_0. \quad (5.5)$$

Thus

$$\left( \frac{\partial h}{\partial \rho_\alpha} \right)_0 = \left( \frac{\partial h}{\partial \theta} \right)_0 = 0, \quad (5.6)$$

where the subscript "o" indicates that the corresponding function is to be evaluated at  $\mathcal{N} = \mathcal{N}_0$ , and we have the following result:

$$\sum_{\alpha} \dot{\mu}_{\alpha} \left( \frac{\partial c_{\alpha}^{+}}{\partial \rho_{\beta}} \right)_0 = \sum_{\alpha} \dot{\mu}_{\alpha} \left( \frac{\partial c_{\alpha}^{+}}{\partial \theta} \right)_0 = 0. \quad (5.7)^1$$

Here  $\dot{\mu}_{\alpha} = \mu_{\alpha}(\vec{\rho}_0, \theta_0)$ . In addition, since  $c_{\alpha}^{+}$  is an isotropic function,

$$\left( \frac{\partial c_{\alpha}^{+}}{\partial \vec{f}_{\beta}} \right)_0 = \left( \frac{\partial c_{\alpha}^{+}}{\partial \vec{g}_{\beta}} \right)_0 = \left( \frac{\partial c_{\alpha}^{+}}{\partial \vec{v}_{\beta}} \right)_0 = 0. \quad (5.8)$$

It follows from (5.2), (5.7), and (5.8) that

$$\Sigma c_{\alpha}^{+}(\mathcal{N}) \dot{\mu}_{\alpha} = O(|\mathcal{N} - \mathcal{N}_0|^2) \text{ as } \mathcal{N} \rightarrow \mathcal{N}_0, \quad (5.9)$$

where

$$|\mathcal{N} - \mathcal{N}_0| = |\vec{\rho} - \vec{\rho}_0| + |\theta - \theta_0| + |\vec{f}| + |g| + |\vec{v}|. \quad (5.10)$$

We say that an equilibrium state  $\mathcal{N}_0$  is strong provided

$$\Sigma c_{\alpha}^{+}(\mathcal{N}) \dot{\mu}_{\alpha} = O(|\mathcal{N} - \mathcal{N}_0|^3) \text{ as } \mathcal{N} \rightarrow \mathcal{N}_0. \quad (5.11)$$

---

<sup>1</sup>Cf. Bowen [1969, Eq. (7.19)].

Remark. To see that this is a natural generalization of the usual notion of strong equilibrium, we assume, for the time being, that there are  $R$  independent chemical reactions:

$$c_{\alpha}^{+} = \sum_{r=1}^R \nu_{\alpha r} J_r, \quad (5.12)$$

where  $\nu_{\alpha r}$  is the stoichiometric coefficient of constituent  $\alpha$  in the reaction  $r$  divided by the molecular mass of  $\alpha$ , and  $J_r = J_r(\mathcal{A})$  is the reaction rate of reaction  $r$ . The chemical affinity of reaction  $r$  is defined by

$$A_r = A_r(\vec{\rho}, \theta) = \sum_{\alpha} \nu_{\alpha r} \mu_{\alpha}(\vec{\rho}, \theta). \quad (5.13)$$

In this instance it is customary to call  $\mathcal{A}_0$  a "strong equilibrium state" provided<sup>1</sup>

$$J_r(\mathcal{A}_0) = A_r(\vec{\rho}_0, \theta_0) = 0. \quad (5.14)$$

In view of (5.12), the first of (5.14) implies (5.2). Further, by (5.12) - (5.14),

$$\sum_{\alpha} c_{\alpha}^{+}(\mathcal{A}) \dot{\mu}_{\alpha} \equiv 0. \quad (5.15)$$

Thus our notion of a strong equilibrium state is somewhat weaker than the standard definition. For all of our results it suffices to use the definition containing (5.11).

---

<sup>1</sup>Truesdell [1969, p. 107]. See also Bowen [1969, p. 121].

With a view toward determining the behavior of the response functions near equilibrium, we first determine some of the more obvious consequences of isotropy. First of all there exist scalar functions  $p_\alpha(\vec{\rho}, \theta)$  such that

$$\vec{T}_\alpha(\vec{\rho}, \theta, \vec{0}, \vec{0}, \vec{0}) = -p_\alpha(\vec{\rho}, \theta) \underline{1}. \quad (5.16)$$

Next, if  $\omega_0$  is an equilibrium state, then, clearly,

$$\vec{z}(\omega_0) = \left( \frac{\partial \vec{z}}{\partial \rho_\alpha} \right)_0 = \left( \frac{\partial \vec{z}}{\partial \theta} \right)_0 = \underline{0} \quad \text{for} \quad \vec{z} = \underline{g} \quad \text{or} \quad \underline{t}_\beta^+. \quad (5.17)$$

Further, there exist scalars  $\kappa, \kappa_\alpha, \gamma_{\alpha\beta}, \gamma_\alpha$ , and  $\lambda_{\alpha\beta}$  such that

$$\begin{aligned} \left( \frac{\partial \underline{g}}{\partial \underline{z}} \right)_0 &= -\kappa \underline{1}, & \left( \frac{\partial \underline{g}}{\partial \underline{v}_\alpha} \right)_0 &= -\kappa_\alpha \underline{1}, \\ \left( \rho_\alpha \frac{\partial \underline{t}_\alpha^+}{\partial \underline{v}_\beta} \right)_0 &= -\gamma_{\alpha\beta} \underline{1}, & \left( \rho_\alpha \frac{\partial \underline{t}_\alpha^+}{\partial \underline{g}} \right)_0 &= -\gamma_\alpha \underline{1}, \end{aligned} \quad (5.18)$$

$$\left( \frac{1}{\rho_\alpha} \frac{\partial p_\alpha}{\partial \rho_\beta} \right)_0 \underline{1} - \left( \frac{\partial \underline{t}_\alpha^+}{\partial \underline{f}_\beta} \right)_0 = \lambda_{\alpha\beta} \underline{1},$$

and, in view of (2.5), (3.4), (4.18), and (5.2),

$$\sum_\alpha \kappa_\alpha = \sum_\alpha \gamma_{\alpha\beta} = \sum_\beta \gamma_{\alpha\beta} = \sum_\alpha \gamma_\alpha = 0. \quad (5.19)$$

We call  $\kappa$  the conductivity and  $\|\gamma_{\alpha\beta}\|$  the momentum supply matrix. The matrix  $\|\lambda_{\alpha\beta}\|$  is of importance in applications. Indeed, if we consider the linearized system of momentum equations appropriate for small departures from the equilibrium

state  $\mathcal{V}_0$ , then (after dividing by  $\rho_\alpha$ ) the term involving  $\text{grad}\rho_\alpha$  in the  $\alpha$ -th equation has the form  $\sum_\beta \lambda_{\alpha\beta} \text{grad}\rho_\beta$ . For this reason we call  $\|\lambda_{\alpha\beta}\|$  the elasticity matrix.<sup>1</sup>

Theorem 5.1. Let  $\mathcal{V}_0$  be a strong equilibrium state.

Then

$$\lambda_{\beta\alpha} = \left( \frac{\partial^2 (\rho\psi)}{\partial \rho_\alpha \partial \rho_\beta} \right)_0, \quad (5.20)$$

$$\left( \frac{\partial g}{\partial \xi_\alpha} \right)_0 = 0, \quad (5.21)$$

$$\frac{\kappa}{\theta_0} a^2 + \sum_\beta \left[ \frac{\kappa}{\theta_0} \beta + \gamma_\beta + \left( \frac{\partial p_\beta}{\partial \theta} \right)_0 + \left( \rho \rho_\beta \frac{\partial s}{\partial \rho_\beta} \right)_0 \right] a w_\beta + \quad (5.22)$$

$$+ \sum_{\alpha, \beta} \gamma_{\alpha\beta} w_\alpha w_\beta \geq 0 \quad \text{for all } a, w_1, \dots, w_N \in \mathbb{R};$$

so that, in particular, the conductivity  $\kappa \geq 0$  and the momentum supply matrix  $\|\gamma_{\alpha\beta}\|$  is positive semi-definite.

Proof. By (4.11) and (5.11),

<sup>1</sup>In fact, Gurtin [1971] has shown that in the purely mechanical theory the linearized equations for the densities have the form

$$\ddot{\rho}_\alpha = \rho_{0\alpha} \sum_\beta \lambda_{\alpha\beta} \Delta \rho_\beta + \text{terms involving } \dot{\rho}_\beta \text{ and } \text{div} \tilde{b}_\beta.$$

Thus the elasticity matrix plays an essential role when studying the propagation of small-amplitude disturbances.



$$\left( \frac{\partial^2 (\Sigma \mu_\alpha c_\alpha^+)}{\partial \tilde{a} \partial \tilde{d}} \right)_0 = \underset{\sim}{0} \quad \text{whenever} \quad \tilde{a} = \underline{f}_\beta, \underline{g}, \text{ or } \underline{v}_\beta \quad \text{and} \quad (5.23)$$

$$\tilde{d} = \underline{f}_\gamma, \underline{g}, \text{ or } \underline{v}_\gamma.$$

Next, it follows from (4.16), (5.3), and (5.23) that

$$\left( \frac{\partial^2 \tilde{h}}{\partial \tilde{f}_\alpha \partial \tilde{f}_\beta} \right)_0 = \underset{\sim}{0}. \quad (5.24)$$

Thus, in view of (5.5),

$$\left( \frac{\partial^2 \tilde{h}}{\partial \tilde{f}_\alpha \partial \tilde{g}} \right)_0 = \left( \frac{\partial^2 \tilde{h}}{\partial \tilde{f}_\alpha \partial \tilde{v}_\beta} \right)_0 = \underset{\sim}{0}. \quad (5.25)$$

The first of (5.25), in conjunction (5.3), (4.16), and (5.23), yields (5.21). On the other hand, the second of (5.25), (5.3), and (5.23) imply

$$\left( \frac{\partial^2 \tilde{k}}{\partial \rho_\alpha \partial \tilde{v}_\beta} \right)_0^T + \sum_\gamma \left( \frac{\partial \tilde{u}_\gamma}{\partial \tilde{v}_\beta} \right)_0^T \left( \frac{\partial \tilde{n}_\gamma}{\partial \tilde{f}_\alpha} \right)_0 - \left( \rho \frac{\partial \psi}{\partial \rho_\alpha} \right)_0 \left( \frac{\partial \tilde{u}_\alpha}{\partial \tilde{v}_\beta} \right)_0^T = \underset{\sim}{0}. \quad (5.26)$$

By (2.5)<sub>2</sub>, (2.10)<sub>2</sub>, and (5.2),

$$\left( \frac{\partial \tilde{n}_\alpha}{\partial \tilde{a}} \right)_0 = \left( \rho_\alpha \frac{\partial \tilde{t}_\alpha^+}{\partial \tilde{a}} \right)_0 \quad \text{and} \quad \sum \left( \frac{\partial \tilde{n}_\alpha}{\partial \tilde{a}} \right)_0 = \underset{\sim}{0} \quad \text{whenever} \quad (5.27)$$

$$a = \underline{f}_\beta, \underline{g}, \text{ or } \underline{v}_\beta,$$

and (4.4), (4.18), (5.16), (5.18)<sub>5</sub>, (5.26), and (5.27) yield, after some manipulation, the result (5.20).

Next, by (5.5),

$$\tilde{a} \cdot \left( \frac{\partial^2 h}{\partial \tilde{g}^2} \right)_0 \tilde{a} + \sum_{\tilde{\beta}} \tilde{a} \cdot \left( \frac{\partial^2 h}{\partial \tilde{v}_{\tilde{\beta}} \partial \tilde{g}} \right)_0 \tilde{w}_{\tilde{\beta}} + \sum_{\alpha, \tilde{\beta}} \tilde{w}_{\alpha} \cdot \left( \frac{\partial^2 h}{\partial \tilde{v}_{\tilde{\beta}} \partial \tilde{v}_{\alpha}} \right)_0 \tilde{w}_{\tilde{\beta}} \leq 0 \quad (5.28)$$

for all vectors  $\tilde{a}, \tilde{w}_1, \dots, \tilde{w}_N$ . If we take  $\tilde{a} = a \tilde{e}$  and  $\tilde{w}_{\alpha} = w_{\alpha} \tilde{e}$  in (5.28), where  $\tilde{e}$  is a unit vector, and use (4.3), (4.4), (4.16), (4.18), (5.3), (5.16), (5.18), (5.23), and (5.27), we are led to (5.22).  $\square$

As a direct consequence of (5.20) we have the following important result.

Corollary 5.1. Let  $\mathcal{V}_0$  be a strong equilibrium state.  
Then the elasticity matrix is symmetric:

$$\lambda_{\alpha\beta} = \lambda_{\beta\alpha}. \quad (5.29)$$

The next corollary follows from (1.1)<sub>4</sub>, (5.17)-(5.19), and (5.21); it asserts that near a strong equilibrium state to within terms of  $O(|\mathcal{V} - \mathcal{V}_0|^2)$   $\tilde{g}$  depends linearly on  $\text{grad}\theta$  and  $\vec{u}$ .

Corollary 5.2. Let  $\mathcal{V}_0$  be a strong equilibrium state.  
Then

$$\tilde{g} = -\kappa \tilde{g} - \sum_{\alpha} \kappa_{\alpha} u_{\alpha} + O(|\mathcal{V} - \mathcal{V}_0|^2) \text{ as } \mathcal{V} \rightarrow \mathcal{V}_0, \quad (5.30)^1$$

<sup>1</sup>Cf. Müller [1968, Eq. (7.28)<sub>1</sub>], Doria [1969, Eq. (7.31)], Bowen and Garcia [1970, Eq. (6.14)]. All of the above treat diffusion without chemical reactions.

where  $g = g(\mathcal{N})$ .

In view of (1.1), we can take  $\rho, c_1, \dots, c_N$  as independent variables in place of  $\rho_1, \dots, \rho_N$ , i.e., e.g.,

$$\mu_\alpha(\rho_1, \dots, \rho_N, \theta) = \mu_\alpha(c_1 \rho, \dots, c_N \rho, \theta) = \mu_\alpha(\rho, \theta, c_1, \dots, c_N). \quad (5.31)$$

By (1.2)<sub>1</sub> and (4.12), the vectors  $\vec{c} = (c_1, \dots, c_N)$  and  $\vec{\mu} = (\mu_1, \dots, \mu_N)$  both lie on planes in  $\mathbb{R}^N$  of dimension  $N-1$ .

We assume that the mapping

$$\vec{c} \mapsto \vec{\mu}(\rho, \theta, \vec{c}) \quad (5.32)$$

is invertible in some neighborhood of  $\mathcal{N}_0$ . Then in this neighborhood we can express the mass supply as follows:

$$c_\alpha^+ = c_\alpha^+(\rho, \theta, \vec{\mu}, \vec{f}, g, \vec{v}). \quad (5.33)$$

Let

$$\tau_{\alpha\beta} = - \left( \frac{\partial c_\alpha^+}{\partial \mu_\beta} \right)_0; \quad (5.34)^1$$

we call  $\|\tau_{\alpha\beta}\|$  the mass supply matrix. The next theorem shows that to within terms of  $O(|\mathcal{N} - \mathcal{N}_0|^2)$   $c_\alpha^+$  depends only on  $\mu_1, \dots, \mu_N$ .

---

<sup>1</sup> The derivative  $\frac{\partial c_\alpha^+}{\partial \mu_1}, \dots, \frac{\partial c_\alpha^+}{\partial \mu_N}$  lies in the "tangent space"

$\{(\varphi_1, \dots, \varphi_N) \in \mathbb{R}^N \mid \sum \varphi_\alpha = 0\}$ .

Theorem 5.2. Let  $\underline{\nu}_0$  be a strong equilibrium state  
and assume that the mapping (5.32) is invertible in some  
neighborhood of  $\underline{\nu}_0$ . Then

$$c_{\alpha}^{+} = -\sum_{\beta} \tau_{\alpha\beta} (\mu_{\beta} - \mu_{\beta}^{\circ}) + o(|\underline{\nu} - \underline{\nu}_0|^2), \quad (5.35)$$

where  $c_{\alpha}^{+} = c_{\alpha}^{+}(\underline{\nu})$ . Moreover, the mass supply matrix  $\|\tau_{\alpha\beta}\|$   
is positive semi-definite and

$$\sum_{\alpha} \mu_{\alpha}^{\circ} \tau_{\alpha\beta} = 0. \quad (5.36)$$

Proof. First, since  $c_{\alpha}^{+}$  is isotropic,

$$\left( \frac{\partial c_{\alpha}^{+}}{\partial \underline{f}_{\beta}} \right)_0 = \left( \frac{\partial c_{\alpha}^{+}}{\partial \underline{g}} \right)_0 = \left( \frac{\partial c_{\alpha}^{+}}{\partial \underline{v}_{\beta}} \right)_0 = \underline{0}. \quad (5.37)$$

If

$$\underline{\underline{f}} = \underline{\underline{0}}, \quad \underline{\underline{g}} = \underline{\underline{0}}, \quad \underline{\underline{v}} = \underline{\underline{0}} \quad (5.38)$$

then, letting

$$\mu'_{\alpha} = \mu_{\alpha} - \mu_{\alpha}^{\circ}, \quad \rho' = \rho - \rho_0, \quad \theta' = \theta - \theta_0, \quad (5.39)$$

$$\varepsilon = \sum |\mu'_{\alpha}| + |\rho'| + |\theta'|,$$

we conclude from (5.2), (5.3), (5.11), (5.33), and (5.34) that

$$0 \geq \sum c_{\alpha}^{+} \mu_{\alpha} = \sum c_{\alpha}^{+} \mu'_{\alpha} + o(\varepsilon^3) = \sum_{\alpha, \beta} \left[ -\tau_{\alpha\beta} \mu'_{\beta} + \left( \frac{\partial c_{\alpha}^{+}}{\partial \rho} \right)_0 \rho' + \left( \frac{\partial c_{\alpha}^{+}}{\partial \theta} \right)_0 \theta' \right] \mu'_{\alpha} + o(\varepsilon^3) \quad (5.40)$$

as  $\epsilon \rightarrow 0$ . Thus

$$\left( \frac{\partial c^+}{\partial \rho} \right)_0 = \left( \frac{\partial c^+}{\partial \theta} \right)_0 = 0 \quad (5.41)$$

and  $\|\tau_{\alpha\beta}\|$  must be positive semi-definite. Finally, the Taylor expansion of (5.33) about  $\psi_0$  reduces to (5.35) when account is taken of (5.2), (5.37), and (5.41).  $\square$

6. Two Lemmas.

Lemma 6.1. Let  $\underline{\mathcal{A}}^* = (\underline{\rho}^*, \theta^*, \underline{f}^*, \underline{g}^*, \underline{v}^*)$  be an arbitrary element of the domain of the response functions. Let  $\underline{x}_0 \in \mathbb{R}^3$ ;  $\omega \in \mathbb{R}$ ;  $\underline{a}_\alpha, \underline{d}_\alpha, \underline{b} \in \mathbb{V}$ ; and suppose that  $\underline{F}_\alpha, \underline{G}$ , and  $\underline{L}_\alpha$  are symmetric tensors. Then there exists a constitutive process whose domain contains  $(\underline{x}_0, 0)$  such that when  $\underline{x} = \underline{x}_0$  and  $t = 0$ :

$$\underline{\mathcal{A}} \equiv (\underline{\rho}, \theta, \text{grad} \underline{\rho}, \text{grad} \theta, \underline{v}) = \underline{\mathcal{A}}^*,$$

$$\dot{\theta} = \omega, \quad \dot{\underline{v}}_\alpha = \underline{a}_\alpha, \quad \overline{\text{grad} \rho_\alpha} = \underline{d}_\alpha, \quad \overline{\text{grad} \theta} = \underline{b}, \quad (6.1)$$

$$\text{grad}^2 \rho_\alpha = \underline{F}_\alpha, \quad \text{grad}^2 \theta = \underline{G}, \quad \text{grad} \underline{v}_\alpha = \underline{L}_\alpha.$$

Proof. Let  $\underline{v}^*$  denote the mixture velocity corresponding to  $\underline{v}^* = (\underline{v}_1^*, \dots, \underline{v}_N^*)$  and  $\underline{\rho}^* = (\rho_1^*, \dots, \rho_N^*)$ , let

$$\underline{r} = \underline{x} - \underline{x}_0, \quad (6.2)$$

and let

$$\begin{aligned} \theta(\underline{x}, t) &= \theta^* + t[\omega - \underline{g}^* \cdot \underline{v}^*] + [\underline{g}^* + t(\underline{b} - \underline{G} \underline{v}^*)] \cdot \underline{r} + \frac{1}{2} \underline{r} \cdot \underline{G} \underline{r}, \\ \underline{v}_\alpha(\underline{x}, t) &= \underline{v}_\alpha^* + t[\underline{a}_\alpha - \underline{L}_\alpha \underline{v}^*] + \underline{L}_\alpha \underline{r} + \frac{1}{2} \underline{r}^2 \underline{e}_\alpha, \\ \rho_{0\alpha}(\underline{x}, t) &= \rho_\alpha^* + \underline{f}_\alpha \cdot \underline{r} + \frac{1}{2} \underline{r} \cdot \underline{F}_\alpha \underline{r}, \end{aligned} \quad (6.3)$$

where  $\underline{e}_\alpha$  is, as yet, unspecified. Then, in view of (1.3),  $\theta$  and  $\underline{v}$  satisfy (6.1)<sub>1,2,3,5,7,8</sub>. Further, since  $\theta^*$  and  $\rho_\alpha^*$  are strictly positive, there exists a neighborhood  $\mathcal{P}_0$  of  $\underline{x}_0$  and a  $t_0 > 0$  such that  $\theta > 0$  on  $\mathcal{P}_0 \times [0, t_0)$  and  $\rho_{0\alpha} > 0$

on  $\mathcal{P}_0$ . Thus we conclude from assumption (A) and the ensuing discussion that there exists a neighborhood  $\mathcal{P} \subset \mathcal{P}_0$  of  $\underline{x}_0$ , a time  $\tau \in (0, t_0)$ , and a constitutive process on  $\mathcal{P} \times [0, \tau)$  corresponding to  $\theta, \vec{v}$ , and  $\vec{\rho}_0$ . Moreover, (3.8) and (6.3)<sub>3</sub> imply that the corresponding density field  $\vec{\rho}$  obeys (6.1)<sub>1,6</sub>. Thus to complete the proof we only need to establish (6.1)<sub>4</sub>. By (2.1), (1.1), and (3.1), it is clear that  $\text{grad} \dot{\rho}_\alpha + \rho_\alpha \text{grad div} \vec{v}_{\sim\alpha}$  can be considered a function of  $(\mathcal{N}, \text{grad}^2 \vec{\rho}, \text{grad}^2 \theta, \text{grad} \vec{v}_{\sim\alpha})$ ; let  $\lambda$  denote the value of this function at  $\mathcal{N} = \mathcal{N}^*$ ,  $\text{grad}^2 \rho_\alpha = \underline{F}_{\sim\alpha}$ ,  $\text{grad}^2 \theta = \underline{G}$ ,  $\text{grad} \vec{v}_{\sim\alpha} = \underline{L}_{\sim\alpha}$  (i.e. the value when  $\underline{x} = \underline{x}_0$ ,  $t = 0$ ). Then, letting  $\underline{L} = \text{grad} \vec{v}_{\sim\alpha}(\underline{x}_0, 0)$ , we conclude from (1.4) and (6.1)<sub>1</sub> that

$$\overline{\text{grad} \dot{\rho}_\alpha}(\underline{x}_0, 0) = -\rho_\alpha^* \text{grad div} \vec{v}_{\sim\alpha}(\underline{x}_0, 0) + \lambda - \underline{L}^T \underline{f}_{\sim\alpha}^*. \quad (6.4)$$

But by (6.3)<sub>2</sub>,

$$\text{grad div} \vec{v}_{\sim\alpha}(\underline{x}_0, 0) = \underline{e}_{\sim\alpha}; \quad (6.5)$$

thus (6.1)<sub>4</sub> will be satisfied provided we take

$$\underline{e}_{\sim\alpha} = \frac{1}{\rho_\alpha^*} [\lambda - \underline{d}_{\sim\alpha}^* - \underline{L}^T \underline{f}_{\sim\alpha}^*]. \quad \square \quad (6.6)$$

Let  $\mathcal{Q}$  denote the orthogonal group  $\mathcal{U}$ . A function  $\underline{f} : \mathcal{U}^M \rightarrow \mathcal{U}$  is isotropic provided

$$\underline{f}(\underline{Q}\underline{w}) = \underline{Q}\underline{f}(\underline{w}) \quad (6.7)$$

for every  $\underline{w} = (w_1, \dots, w_M) \in \mathcal{V}^M$  and every  $\underline{Q} \in \mathcal{Q}$ , where

$$\underline{Qw} = (Qw_1, \dots, Qw_M). \quad (6.8)$$

Lemma 6.2.<sup>1</sup> Let  $f : \mathcal{V}^M \rightarrow \mathcal{V}$  be an isotropic function of class  $C^3$ , and assume that for every  $\underline{w} \in \mathcal{V}^M$  and every  $m \in \{1, \dots, M\}$  the tensor  $\frac{\partial f(\underline{w})}{\partial w_m}$  is skew. Then  $\underline{f} \equiv \underline{0}$ .

Proof. Choose an orthonormal basis for  $\mathcal{V}$  and let  $f\langle i \rangle$  and  $w_m\langle i \rangle$  ( $i = 1, 2, 3$ ) denote the corresponding components of  $\underline{f}$  and  $w_m$ . By hypothesis,

$$\frac{\partial f\langle i \rangle}{\partial w_m\langle j \rangle} = -\frac{\partial f\langle j \rangle}{\partial w_m\langle i \rangle}, \quad (6.9)$$

so that

$$\frac{\partial f\langle i \rangle}{\partial w_m\langle i \rangle} = 0. \quad (6.10)$$

Equations (6.9) and (6.10) imply that

$$\frac{\partial^2 f\langle i \rangle}{\partial w_m\langle j \rangle \partial w_n\langle j \rangle} = -\frac{\partial^2 f\langle j \rangle}{\partial w_m\langle j \rangle \partial w_n\langle i \rangle} = 0; \quad (6.11)$$

hence

$$\frac{\partial^3 f\langle i \rangle}{\partial w_m\langle j \rangle \partial w_n\langle k \rangle \partial w_p\langle \ell \rangle} = 0, \quad (6.12)$$

since two of  $i, j, k$ , and  $\ell$  must coincide. Therefore  $\underline{f}$  must have the form

$$\underline{f}(\underline{w}) = \underline{f}(\underline{0}) + \underline{F}(\underline{w}) + \underline{G}(\underline{w}, \underline{w}), \quad (6.13)$$

<sup>1</sup>For  $M = 3$  this theorem is a corollary of a result obtained by Doria [1969, §6] using different methods. See also Müller [1968, §6]. When I first established this result I was unaware of Doria's result.



where  $\underline{F} : \mathcal{V}^M \rightarrow \mathcal{V}$  is linear and  $\underline{G} : \mathcal{V}^M \times \mathcal{V}^M \rightarrow \mathcal{V}$  is symmetric and bilinear. It follows from (6.13) and (6.7) with  $\underline{Q} = -\underline{1}$  that  $\underline{f}(\underline{0}) = \underline{Q}$  and  $\underline{G} = \underline{Q}$ ; therefore

$$\underline{f}(\underline{w}) = \underline{F}(\underline{w}) = \sum_{m=1}^M \underline{F}_m w_m, \quad (6.14)$$

where each  $\underline{F}_m$  is a tensor. Further,

$$\underline{F}_m = \frac{\partial \underline{f}(\underline{w})}{\partial w_m}, \quad (6.15)$$

so that  $\underline{F}_m$  is skew. Finally, by (6.7) and (6.14),

$$\underline{Q} \underline{F}_m = \underline{F}_m \underline{Q} \quad (6.16)$$

for every  $\underline{Q} \in \mathcal{O}$ , and the only skew tensor with this property is  $\underline{F}_m = \underline{Q}$ .  $\square$

## REFERENCES

- [1940] Eckart, C., The thermodynamics of irreversible processes, II. Fluid mixtures, *Phys. Rev.* 58, 269-275.
- [1957] Truesdell, C., Sulle basi della termomeccanica, *Rend. Accad. Lincei* 22, 33-88, 158-166. English transl. in Rational Mechanics of Materials, *Intl. Sci. Rev. Ser.* 292-305, Gordon and Breach, New York, 1965.
- [1959] Meixner, J. and H. G. Reik, Thermodynamik der irreversiblen Prozesse, The Encyclopedia of Physics, Vol. 111/2, edited by S. Flügge, Springer, Berlin-Göttingen-Heidelberg.
- Nachbar, W., F. Williams, and S. S. Penner, The conservation equations for independent coexistent continua and for multicomponent reacting gas mixtures, *Quart. Appl. Math.* 17, 43-54.
- [1962] de Groot, S. R. and P. Mazur, Non-Equilibrium Thermodynamics, Interscience, New York.
- [1963] Coleman, B. D. and W. Noll, The thermodynamics of elastic materials with heat conduction and viscosity, *Arch. Rational Mech. Anal.* 13, 167-178
- [1964] Kelly, P. D., A reacting continuum, *Int. J. Engng. Sci.* 2, 129-153.
- [1965] Eringen, A. C. and J. D. Ingram, A continuum theory of chemically reacting media, I, *Int. J. Engng. Sci.* 3, 197-212.
- Green, A. E. and P. M. Naghdi, A dynamical theory of interacting continua, *Int. J. Engng. Sci.* 3, 231-241.
- [1966] Crochet, M. J. and P. M. Naghdi, Constitutive equations for flow of fluid through an elastic solid, *Int. J. Engng. Sci.* 4, 383-401.
- Green, A. E. and T. R. Steel, Constitutive equations for interacting continua, *Int. J. Engng. Sci.* 4, 483-500.
- Mills, N., Incompressible mixture of Newtonian fluids, *Int. J. Engng. Sci.* 4, 97-112.
- [1967] Atkin, R. J., Constitutive theory for a mixture of an isotropic elastic solid and a non-Newtonian fluid, *Zeit. angew. Math. Phys.* 18, 803-825.

- Bowen, R. M., Towards a thermodynamics and mechanics of mixtures, Arch. Rational Mech. Anal. 24, 370-403.
- Coleman, B. D. and M. E. Gurtin, Thermodynamics with internal state variables, J. Chem. Phys. 47, 597-613.
- Crochet, M. J. and P. M. Naghdi, Small motions superposed on large static deformations in porous media, Acta Mech. 4, 315-335.
- Green, A. E. and P. M. Naghdi, A theory of mixtures, Arch. Rational Mech. Anal. 24, 243-263.
- Ingram, J. D. and A. C. Eringen, A continuum theory of chemically reacting continua, II. Constitutive equations of reacting fluid mixtures, Int. J. Engng. Sci. 5, 289-322.
- Mills, N., A theory of multi-component mixtures, Quart. J. Mech. Appl. Math. 20, 499-508.
- [1968] Dunwoody, N. T. and I. Müller, A thermodynamic theory of two chemically reacting ideal gases with different temperatures, Arch. Rational Mech. Anal. 29, 344-369.
- Green, A. E. and P. M. Naghdi, A note on mixtures, Int. J. Engng. Sci. 6, 631-635.
- Müller, I., A thermodynamic theory of mixtures of fluids, Arch. Rational Mech. Anal. 28, 1-39.
- Truesdell, C., Sulle basi della termodinamica delle miscele, Rend. Accad. Naz. Lincei, (8) 44, 381-383.
- [1969] Bowen, R. M., The thermochemistry of a reacting mixture of elastic materials with diffusion, Arch. Rational Mech. Anal. 34, 97-127.
- Bowen, R. M. and J. C. Wiese, Diffusion in mixtures of elastic materials, Int. J. Engng. Sci. 7, 689-722.
- Doria, M. L., Some general results for non-reacting binary mixtures of fluids, Arch. Rational Mech. Anal. 32, 343-368.
- Green, A. E. and P. M. Naghdi, On basic equations for mixtures, Quart. J. Mech. Appl. Math. 22, 427-438.

Truesdell, C., Rational Thermodynamics: A Course of Lectures on Selected Topics, McGraw-Hill, New York.

[1970] Bowen, R. M. and D. J. Garcia, On the thermodynamics of mixtures and several temperatures, Int. J. Engng. Sci. 8, 63-83.

Craine, R. E., A. E. Green, and P. M. Naghdi, A mixture of viscous elastic materials with different constituent temperatures, Quart. J. Mech. Appl. Math. 23, 171-184.

Dunwoody, N. T., A thermomechanical theory of diffusion in solid-fluid mixtures, Arch. Rational Mech. Anal. 38, 348-371.

[1971] Gurtin, M. E., Some remarks on the mechanical theory of fluid mixtures, forthcoming.

Carnegie-Mellon University  
Pittsburgh, Pa. 15213 U.S.A.