

NAMT
92-042

**Anisotropic Motion of an Interface
Relaxed by the Formation of
Infinitesimal Wrinkles**

**Morton E. Gurtin
H.M. Soner
Carnegie Mellon University**

**P.E. Souganidis
University of Wisconsin**

Research Report No. 92-NA-042

December 1992

Sponsors

**U.S. Army Research Office
Research Triangle Park
NC 27709**

**National Science Foundation
1800 G Street, N.W.
Washington, DC 20550**

ANISOTROPIC MOTION OF AN INTERFACE RELAXED BY THE FORMATION OF
INFINITESIMAL WRINKLES

by

M. E. Gurtin and H. M. Soner

Department of Mathematics, Carnegie Mellon University

P. E. Souganidis

Department of Mathematics, University of Wisconsin, Madison

1. INTRODUCTION.

1.1. MATHEMATICAL THEORY.

In this paper we discuss the motion, in the plane, of a region $\Omega(t)$ whose boundary-curve evolves from a given region Ω_0 according to an equation

$$(1.1) \quad B(\theta)V = G(\theta)K - U$$

with V the **normal velocity** and K the **curvature**. (Our sign convention is such that the positive normal-direction is outward from $\partial\Omega = \partial\Omega(t)$, and $K < 0$ when $\partial\Omega$ is a circle.) Here $B(\theta)$ and $G(\theta)$ are given functions of the **normal-angle** θ , which is the counterclockwise angle from a fixed axis to the outward normal of $\partial\Omega$, and U is a given constant.

For $B(\theta)$ and $G(\theta)$ continuous and strictly positive, (1.1) is a parabolic equation that is well understood, with fairly well-behaved solutions.¹ There are, however, situations of physical importance for which $G(\theta) = 0$ over certain angle-intervals and for which $G(\theta)$ need not be continuous (cf. §1.2). Here we will develop a fairly complete theory of (1.1) under the following assumptions:

(1.2a) G is piecewise continuous and ≥ 0 , and continuous on any interval of strict positivity;

¹Cf. Angenent [Ag]; Chen, Giga, and Goto [CGG]; Soner [Sol]; Barles, Soner, and Souganidis [BSS].

(1.2b) B is continuous and >0 .

In some instances we will add the hypothesis:

(1.3) B has polar diagram a straight line on any angle interval for which $G=0$,

which is based on the underlying physics.

Because of the lack of continuity of G as well as the degeneracy of (1.1) when $G=0$, it is convenient to discuss this equation within the weak framework of viscosity solutions. This approach to geometric equations, initiated by Evans and Spruck [ES1] and Chen, Giga, and Goto [CGG], is based on the use of level sets to characterize evolving curves, an idea due to Sethian [Se], Osher and Sethian [OS], and Barles [Ba]. Here — to study (1.1) — we will use this approach as well as an intrinsic approach given by Soner [So] and Barles, Soner, and Souganidis [BSS]. The difficulties concerning (1.1) result from the *discontinuous* nature of G ; the degeneracy of the equation, at angles θ with $G(\theta)=0$, causes no great difficulty; were G continuous, most of our results would follow from those in [CGG].

Our main results, for evolution from a given compact region Ω_0 , consist of: a theorem of existence and local uniqueness; a global comparison theorem² for level-set solutions.

1.2. PHYSICAL BACKGROUND.

There are situations of interest in which the motion of a phase interface is essentially independent of the behavior of the corresponding bulk phases. One of the first models of such phenomena was proposed by Mullins [Mu] to study the *planar* motion of grain boundaries; the resulting evolution equation has the form³

$$(1.4) \quad V = K$$

after an appropriate scaling Equation (1.4) is a parabolic PDE with a large literature;⁴ its major consequence ([GH], [Gr]) is that all such boundary

²This comparison theorem was established independently by Ohnuma and Sato [OS], whose proof is different (and more concise) than ours.

³Allen and Cahn [AC] and Rubinstein, Sternberg, and Keller [RSK] deduce the equation $V=K$ as a formal approximation to the Landau-Ginzburg equation, a result established rigorously in [BSS], [ESS], [Ch], and [DS]. See also [ORS], [OwS], [RS].

curves, irrespective of their initial shape, shrink to a point in finite time, with asymptotic shape a circle.

Mullins's theory was generalized in [G1,§8], [AG1] to include anisotropy and the possibility of a difference in bulk energies between phases. The resulting equation is

$$(1.5) \quad b(\theta)V = g(\theta)K - U,$$

where $g(\theta)$, the **energy modulus**, is given by

$$(1.6) \quad g(\theta) = f(\theta) + f''(\theta)$$

with $f(\theta) > 0$ the **interfacial energy**; U is the **relative energy** of the material in Ω ; and $b(\theta) > 0$, the **kinetic modulus**, is a material function. The presence of the angle θ reflects anisotropy, and the particular form in which f appears in (1.6) is a consequence of thermodynamics. In fact, a consequence of (1.5) and (1.6) is the thermodynamic inequality

$$(1.7) \quad \frac{d}{dt} \left\{ \int_{\partial\Omega(t)} f(\theta) ds + U \text{area}(\Omega(t)) \right\} = - \int_{\partial\Omega(t)} b(\theta) V^2 ds.$$

When

$$(1.8) \quad g(\theta) > 0$$

evolution according to (1.5) is governed by a parabolic PDE and the underlying problem is not much different than that for the equation $V=K$. What makes (1.5) nonstandard is the possibility of interfacial energies that satisfy

$$(1.9) \quad g(\theta) < 0$$

⁴Cf. Brakke [Br], Sethian [Se], Abresch and Langer [AL], Gage and Hamilton [GH], Grayson [Gr], Osher and Sethian [OS], Evans and Spruck [ES1-3], Chen, Giga, and Goto [CGG], Goto and Sato [GS], Almgren, Taylor, and Wang [ATW], Taylor, Cahn, and Handwerker [TCH], and the references therein.

$$(1.12) \quad G(\theta) = F(\theta) + F''(\theta).$$

The next question we must answer is what is an appropriate kinetic modulus for the infinitesimally wrinkled curve. If $\Gamma(t)$ is a finite wrinkling whose facets have θ_1 and θ_2 as normal angles, then $\Gamma(t)$ evolves as a rigid body with constant velocity ω defined by [AG1]

$$(1.16) \quad \omega \cdot N(\theta_1) = -b(\theta_1)^{-1}U, \quad \omega \cdot N(\theta_2) = -b(\theta_2)^{-1}U$$

(although $\Gamma(t)$ is allowed to shrink or grow tangentially). Since ω depends on the particular wrinkling only through θ_1 and θ_2 , it seems reasonable to suppose that *infinitesimal wrinklins* with θ_1 and θ_2 as normal angles also evolve with rigid velocity ω , and this is equivalent to replacing the kinetic modulus $b(\theta)$ between θ_1 and θ_2 by an effective modulus $B(\theta)$ that agrees with $b(\theta)$ at θ_1 and θ_2 and has polar diagram between θ_1 and θ_2 a straight line:

$$(1.17) \quad B(\theta)^{-1} = \mu_1(\theta)b(\theta_1)^{-1} + \mu_2(\theta)b(\theta_2)^{-1}$$

This procedure defines an *effective kinetic modulus* $B(\theta)$ for all θ [G2]: $B(\theta) > 0$ is continuous; $B(\theta) = b(\theta)$ for all GS angles θ ; the polar diagram of $B(\theta)$ is a straight line over normal-angle intervals with $f(\theta) > F(\theta)$.

We will refer to G and B derived in this manner as the **effective moduli** corresponding to f and g .

We are therefore led to the **relaxed evolution equation** (1.1) with B and G the effective moduli corresponding to f and g [G2]. It is important to note that this relaxed equation coincides with our original system (1.5) at all GS angles θ . Note also that, because of the construction of $G(\theta)$, no matter how smooth $f(\theta)$ is,

$$(1.18) \quad G(\theta) \text{ will generally be discontinuous}$$

whenever the angle θ changes from GS to GUS; this property of $G(\theta)$

renders the relaxed evolution equation nonstandard. In addition, $G(\theta)=0$ whenever θ is GUS, so that (1.1) *degenerates to hyperbolic* at GUS angles.

Our main results of physical interest are:

1° Viscosity solutions of (1.1) not only satisfy (1.5) away from corners, but, what is most interesting, such solutions automatically satisfy the force balance (1.11) across corners.

2° If (θ_1, θ_2) is a GUS angle-interval, then a wedge whose two sides have normal angles θ_1 and θ_2 and evolve according to $b(\theta_1)V=-U$ and $b(\theta_2)V=-U$, respectively, is a solution of the basic equations (1.5) and (1.11) [AG1, §9]. We show that our choice of the effective moduli G and B is the only possible choice if all such wedges are to be viscosity solutions of (1.1). What makes this result so interesting is that $G(\theta)$ and $B(\theta)$ differ from $g(\theta)$ and $b(\theta)$ only at angles θ that are not globally stable, and wedges by definition do not involve such θ .

3° For $U < 0$ and Ω_0 large enough, $t^{-1}\Omega(t)$ converges to a dilation of the Wulff region for $1/B(\theta)$.⁹

⁹This result, conjectured by Angenent and Gurtin [AG1], was proved by Soner [So] for $G > 0$ and B with a convex polar diagram, and extended in [AG2] to general $B > 0$.

2. CLASSICAL EVOLUTIONS. WRINKLINGS AND WEDGES.

Throughout the paper we restrict attention to energies $f(\theta)$ and kinetic moduli $b(\theta)$ that are consistent with the following hypotheses:

(2.1a) f is C^2 and >0 ;

(2.1b) each convexifying tangent to the Frank diagram \mathcal{F} intersects \mathcal{F} at most at two angles, and there are at most a finite number of such tangents;

(2.1c) $g(\theta) > 0$ at each GS angle θ ;

(2.1d) b is continuous and strictly positive.

We begin with a discussion of regions whose boundaries evolve according to (1.5), but with normal angles constrained to be GS, so that (1.5) is parabolic. Such boundaries will generally contain corners - consistent with (1.11) - for which the jump in normal angle removes angles of backward parabolicity of (1.5).¹⁰ Not all initial data are consistent with evolutions of this type; in particular, the initial region Λ must be **admissible** in the sense that¹¹

(2.2a) Λ is closed with $\partial\Lambda$ piecewise C^2 , and at each point of smoothness the (outward) normal angle θ is GS (so that $g(\theta) > 0$);

(2.2b) (1.11) is satisfied.

For each $t \in (0, T)$, let $\Omega(t) \subset \mathbb{R}^2$ be given. Then $\Omega(t)$ is a **classical evolution** in $(0, T)$ if:

(2.3a) $\Omega(t)$ is admissible at each $t \in (0, T)$;

(2.3b) the evolution equation (1.5) is satisfied on each interval of smoothness of $\partial\Omega(t)$ (up to the endpoints).

If, in addition,

$$(2.4) \quad \Omega(0^+) = \Omega_0.$$

¹⁰The motivation for considering such regions can be found in [AG1,S9], [AG2,S2], [G2,S11].

¹¹An assumption of piecewise smoothness for a boundary curve Γ will always contain the tacit assumption that Γ is locally graphlike, so that, e.g., sets with boundary a "figure 8" are ruled out.

then $\Omega(t)$ is a classical evolution from Ω_0 .

Theorem 2.1 (Existence and Uniqueness of Classical Evolutions [AG2]).
Let Ω_0 be bounded and admissible. Then there is a unique maximal classical evolution $\Omega(t)$, $t \in [0, T_{\max})$, from Ω_0 . Moreover, $\partial\Omega(t)$ is piecewise C^∞ at each $t \in (0, T_{\max})$.

By definition, if the boundary curve $\partial\Omega(t)$ of a classical evolution $\Omega(t)$ has a corner corresponding to an angle jump from θ_1 to θ_2 , then (θ_1, θ_2) is a GUS angle-interval and $C(\theta_1) = C(\theta_2)$. Suppose that θ_1, θ_2 is such a pair. Then we can construct classical evolutions, called (θ_1, θ_2) -**wrinklings** (Figure 2.1), whose normal angles jump back and forth between θ_1 and θ_2 [AG1, S9]; the flat portions of the wrinkling with angle θ_i ($i=1,2$) are then called θ_i -**facets**. By (1.5), each θ_i -facet evolves according to

$$(2.5) \quad V = -b(\theta_i)U,$$

and from this we may conclude that the wrinkling itself evolves as a rigid body with velocity ω given by (1.16). A (θ_1, θ_2) -wrinkling with a single corner is called a (θ_1, θ_2) -**wedge**. A (θ_1, θ_2) -wedge $\Omega(t)$ is prescribed by specifying: (i) whether $\Omega(t)$ is convex or concave; (ii) the position of the corner at some time.

Suppose that $\partial\Omega_0$ is a piecewise flat curve whose normal angle jumps back and forth between θ_1 and θ_2 , with $\partial\Omega_0$ the c -level set of an auxiliary function Φ_0 ; i.e.,

$$(2.6) \quad \partial\Omega_0 = \{ \mathbf{x} \in \mathbb{R}^2 : \Phi_0(\mathbf{x}) = c \}, \quad \Omega_0 = \{ \mathbf{x} \in \mathbb{R}^2 : \Phi_0(\mathbf{x}) \geq c \}.$$

Then Ω_0 is the initial set of a (θ_1, θ_2) -wrinkling $\Omega(t)$ if and only if

$$(2.7) \quad \Omega(t) = \Omega_0 + t\omega = \{ \mathbf{x} \in \mathbb{R}^2 : \Phi(t, \mathbf{x}) \geq c \}, \quad \Phi(t, \mathbf{x}) = \Phi_0(\mathbf{x} - t\omega).$$

3. VISCOSITY SOLUTIONS. RELAXED EVOLUTIONS.

We will use the relaxed equation (1.1) to discuss evolution from an initial region that has normal-angles θ with $g(\theta) < 0$. In the derivation of (1.1), G and B are the effective moduli for f and b , but we will generally require only that G and B satisfy (1.2).

a. DEFINITIONS.

We are interested in the relaxed evolution problem defined by the relaxed equation (1.1) supplemented by the initial condition (2.4):

$$(E) \quad B(\theta)V = G(\theta)K - U, \quad \Omega(0^+) = \Omega_0.$$

Suppose that B , G , and Ω_0 are such that (E) has a smooth solution $\Omega(t)$ with $\partial\Omega(t)$ the c -level set of an auxiliary function Φ :

$$(3.3) \quad \partial\Omega(t) = \{ \mathbf{x} \in \mathbb{R}^2 : \Phi(t, \mathbf{x}) = c \}, \quad \Omega(t) = \{ \mathbf{x} \in \mathbb{R}^2 : \Phi(t, \mathbf{x}) \geq c \}.$$

Assume further that Φ is a smooth function whose spatial gradient $\nabla\Phi$ has $|\nabla\Phi(t, \mathbf{x})|$ never zero on $\partial\Omega(t)$. Then Φ satisfies the PDE

$$(3.4) \quad \Phi_t = \mathcal{F}(\nabla\Phi, \nabla^2\Phi),$$

where $\nabla^2\Phi$ is the Hessian matrix of second spatial derivatives of Φ , while

$$(3.5) \quad \begin{aligned} \mathcal{F}(\mathbf{p}, \mathbf{A}) &= B(\theta)^{-1} \{ G(\theta)T(\theta) \cdot \mathbf{A}T(\theta) - U|\mathbf{p}| \} \\ &= B(\theta)^{-1} \{ G(\theta) \operatorname{tr}[(I - \bar{\mathbf{p}}\bar{\mathbf{p}})\mathbf{A}] - U|\mathbf{p}| \}, \\ \theta &= \sin^{-1}(-\bar{\mathbf{p}}_2), \quad \bar{\mathbf{p}} = \mathbf{p}/|\mathbf{p}| \end{aligned}$$

for all vectors $\mathbf{p} \neq 0$ and all symmetric matrices \mathbf{A} . Thus solving (E) at least formally reduces to solving (3.4) subject to an initial condition

$$(3.10) \quad \Phi(\mathbf{x}, 0) = \Phi_0(\mathbf{x})$$

for all $\mathbf{x} \in \mathbb{R}^2$, where Φ_0 is an auxiliary function satisfying

$$(3.9) \quad \partial\Omega_0 = \{ \mathbf{x} \in \mathbb{R}^2 : \Phi_0(\mathbf{x}) = c \}, \quad \Omega_0 = \{ \mathbf{x} \in \mathbb{R}^2 : \Phi_0(\mathbf{x}) \geq c \}.$$

\mathcal{F} defined by (3.5) has two chief properties upon which much of the level-set theory of (3.4) is based: the **geometric property**

$$(3.12) \quad \mathcal{F}(\lambda \mathbf{p}, \lambda \mathbf{A} + \nu \mathbf{p} \otimes \mathbf{p}) = \lambda \mathcal{F}(\mathbf{p}, \mathbf{A})$$

for all $\lambda \geq 0$, $\nu \in \mathbb{R}$; and the **elliptic property**

$$(3.13) \quad \mathcal{F}(\mathbf{p}, \mathbf{A} + \mathbf{B}) \geq \mathcal{F}(\mathbf{p}, \mathbf{A})$$

whenever \mathbf{B} is symmetric and positive semi-definite.

The level-set method is not intrinsic, since it requires data irrelevant to the problem: namely the values of Φ_0 away from an arbitrary small neighborhood of $\partial\Omega_0$. A method of circumventing this is to work with the characteristic function¹²

$$(3.14) \quad u(t, \mathbf{x}) = \chi_{\Omega(t)}(\mathbf{x})$$

of the region $\Omega(t)$. It is reasonable to expect that u should, in some sense, satisfy (3.4), an expectation motivated by viewing u as the limit of a sequence $\{\Phi_k\}$ of functions Φ_k consistent with (3.3) for, say, $c = \frac{1}{2}$. We will use the theory of viscosity solutions¹³ to define the sense in which u satisfies (3.4).

Let h be a bounded scalar function on a subset \mathcal{H} of \mathbb{R}^n ; then h^* and h_* , respectively, denote the **upper and lower semicontinuous envelopes** of h defined on $\text{cl } \mathcal{H}$ by

$$(3.6) \quad h^*(\mathbf{z}) = \limsup_{\mathbf{q} \rightarrow \mathbf{z}} h(\mathbf{q}), \quad h_*(\mathbf{z}) = \liminf_{\mathbf{q} \rightarrow \mathbf{z}} h(\mathbf{q})$$

¹²Cf. [BSS].

¹³Cf. Crandall and Lions [CL], Crandall, Evans, and Lions [CEL], Jensen [Je]. A recent article of Crandall, Ishii, and Lions [CIL] provides an excellent survey of the subject.

$q \in \mathcal{K}$ $q \in \mathcal{K}$

Let u be a bounded function on $[0, \infty) \times \mathbb{R}^2$. Then u is a **viscosity subsolution** of (3.4) if, for every (scalar) test function $w \in C^{1,2}((0, \infty) \times \mathbb{R}^2)$,

$$(3.7) \quad w_t(t_0, \mathbf{x}_0) \leq \mathcal{F}^*(\nabla w(t_0, \mathbf{x}_0), \nabla^2 w(t_0, \mathbf{x}_0))$$

at every local maximum of $u^* - w$; u is a **viscosity supersolution** of (3.4) if, for every such w ,

$$(3.8) \quad w_t(t_0, \mathbf{x}_0) \geq \mathcal{F}_*(\nabla w(t_0, \mathbf{x}_0), \nabla^2 w(t_0, \mathbf{x}_0))$$

at every local minimum of $u_* - w$; u is a **viscosity solution** of (3.4) if u is both a viscosity subsolution and a viscosity supersolution of (3.4) [CGG]. We will also use viscosity subsolutions, supersolutions, and solutions on finite time intervals $(0, T)$.

Let $\Omega(t)$, $t \geq 0$, be given, and define $u(t, \mathbf{x}) = \chi_{\Omega(t)}(\mathbf{x})$. Then $\Omega(t)$, $t \geq 0$, is a χ -**subsolution** or a χ -**supersolution** of (1.1) according as u is a viscosity subsolution or a viscosity supersolution of (3.4) and $\Omega(t)$ is uniformly bounded on compact time intervals; $\Omega(t)$, $t \geq 0$, is a **relaxed evolution** if it is both a χ -subsolution and a χ -supersolution of (1.1).

Let

$$\Omega^*(0^+) = \left\{ \mathbf{x} \in \mathbb{R}^2 : \limsup_{t \rightarrow 0^+, \mathbf{y} \rightarrow \mathbf{x}} u^*(t, \mathbf{y}) = 1 \right\},$$

$$\Omega_*(0^+) = \left\{ \mathbf{x} \in \mathbb{R}^2 : \liminf_{t \rightarrow 0^+, \mathbf{y} \rightarrow \mathbf{x}} u_*(t, \mathbf{y}) = 1 \right\},$$

so that $\Omega^*(0^+)$ is closed, while $\Omega_*(0^+)$ is open. Then $\Omega(t)$, $t \geq 0$, is:

- (a) a χ -**subsolution** of (1.1) compatible with Ω_0 if it is a χ -subsolution and $\Omega^*(0^+) \subseteq \text{cl } \Omega_0$;
- (b) a χ -**supersolution** of (1.1) compatible with Ω_0 if it is a χ -supersolution and $\Omega_*(0^+) \supseteq \text{int } \Omega_0$;
- (c) a **relaxed evolution** from Ω_0 if it is a χ -subsolution of (1.1)

compatible with Ω_0 as well as a χ -supersolution of (1.1) compatible with Ω_0 .

Note that, if a relaxed evolution is to take on initial data Ω_0 in a classical sense, then Ω_0 must be regular (i.e., $\text{cl } \Omega_0 = \text{cl}(\text{int } \Omega_0)$).

One should expect lack of (global) uniqueness for relaxed evolutions from a given initial set;¹⁴ with this in mind, we introduce the following definitions: the **upper** and **lower envelopes** $\mathcal{U}(t)$ and $\mathcal{L}(t)$ for relaxed evolutions from an initial set Ω_0 are defined at each $t \geq 0$ by¹⁵

$$\mathcal{U}(t) = \text{cl} \left\{ \text{union of all values at } t \text{ of } \chi\text{-subsolutions of (1.1)} \right. \\ \left. \text{compatible with } \Omega_0 \right\},$$

$$\mathcal{L}(t) = \text{int} \left\{ \text{intersection of all values at } t \text{ of } \chi\text{-supersolutions of (1.1)} \right. \\ \left. \text{compatible with } \Omega_0 \right\};$$

the **graph up to time** T of a time-dependent set $A(t)$ is defined by

$$\text{graph}_T A = \bigcup_{0 \leq t \leq T} [A(t) \times \{t\}];$$

the time

$$T_{\text{uniq}} = \sup \left\{ T : \text{graph}_T \mathcal{U} = \text{cl}(\text{graph}_T \mathcal{L}) \text{ and } \text{int}(\text{graph}_T \mathcal{U}) = \text{graph}_T \mathcal{L} \right\}$$

is the **uniqueness time** for relaxed evolutions from Ω_0 and, for $T_{\text{uniq}} > 0$,

$$\Omega_{\text{uniq}}(t) = \mathcal{U}(t) = \text{cl } \mathcal{L}(t), \quad t \in [0, T_{\text{uniq}})$$

is the **unique relaxed evolution** from Ω_0 .

b. EXISTENCE AND UNIQUENESS.

We assume throughout this subsection that

¹⁴For G continuous there are conditions that guarantee the uniqueness of solutions [BSS], [So]. For motion by mean curvature and smooth initial data ($V=K$) uniqueness holds generically [ES3].

¹⁵Cf. [So, Section 11].

G and B satisfy (1.2);

Ω_0 is a prescribed initial domain, assumed compact.

(Note that we do not require the consistency of B with (1.3)).

Theorem 3.1 (Existence and Local Uniqueness of Relaxed Evolutions).

(a) *there is at least one relaxed evolution from Ω_0 ;*

(b) *the upper and lower envelopes are relaxed evolutions from Ω_0 .*

If, in addition, $\partial\Omega_0$ is C^3 , then

(c) *the uniqueness time for relaxed evolutions from Ω_0 is strictly positive.*¹⁶

We postpone, until Section 8, the proof of this theorem and the next.

Let $\hat{M}([0, T] \times \mathbb{R}^2)$ denote the set of all bounded functions on $[0, T] \times \mathbb{R}^2$ that are equal to a constant outside of a large ball; i.e., $\varphi \in \hat{M}([0, T] \times \mathbb{R}^2)$ if and only if there are constants α and R such that $\varphi(t, \mathbf{x}) = \alpha$ for $|\mathbf{x}| \geq R$; here α and R may depend on φ . We define $\hat{M}(\mathbb{R}^2)$ similarly. Finally,

$$\hat{M}([0, \infty) \times \mathbb{R}^2) = \bigcap_{T > 0} \hat{M}([0, T] \times \mathbb{R}^2);$$

i.e., $\varphi \in \hat{M}([0, \infty) \times \mathbb{R}^2)$ if and only if for every T there are constants α_T and R_T satisfying $\varphi(t, \mathbf{x}) = \alpha_T$ for $|\mathbf{x}| \geq R_T$ and $t \in [0, T]$.

Let Φ_0 be an **auxiliary** function for the initial set Ω_0 ; that is, a continuous function $\Phi_0 \in \hat{M}(\mathbb{R}^2)$ satisfying (3.9). Then $\Phi \in \hat{M}([0, \infty) \times \mathbb{R}^2)$ is a **level-set solution** of (1.1) if Φ is a *continuous* viscosity solution of (3.4); if, in addition, Φ satisfies the initial condition (3.10), then Φ **corresponds to Φ_0** .

Theorem 3.2 (Existence and Uniqueness of Level-Set Solutions).¹⁷

There is a unique level-set solution of (1.1) corresponding to any given

¹⁶If Ω_0 is strictly star-shaped, then $T_{\text{uniq}} = \infty$ (cf. [So, Section 9]; a more general condition is given by [BSS, Section 4]).

¹⁷For G continuous and nonnegative uniqueness and existence follow from Theorem 6.8 of [CGG].

choice of auxiliary function Φ_0 for Ω_0 . Moreover, the upper and lower envelopes for relaxed evolutions from Ω_0 are given by

$$(5.7) \quad \mathcal{U}(t) = \{ \mathbf{x} : \Phi(t, \mathbf{x}) \geq c \}, \quad \mathcal{L}(t) = \{ \mathbf{x} : \Phi(t, \mathbf{x}) > c \}.$$

Thus the sets $\{ \mathbf{x} : \Phi(t, \mathbf{x}) = c \}$, $\{ \mathbf{x} : \Phi(t, \mathbf{x}) \geq c \}$, and $\{ \mathbf{x} : \Phi(t, \mathbf{x}) > c \}$ are independent of the choice of auxiliary function Φ_0 .¹⁸

c. COMPARISON.

In this subsection we state comparison theorems related to weak solutions of (1.1) and (3.4). We assume throughout that

G and B satisfy (1.2);

we do not require (1.3). The next theorem is the key technical result of the paper.

Theorem 3.3.¹⁹ Let $\varphi \in \hat{M}([0, T] \times \mathbb{R}^2)$ be a viscosity subsolution and $\psi \in \hat{M}([0, T] \times \mathbb{R}^2)$ a viscosity supersolution, both of (3.4) on $(0, T) \times \mathbb{R}^2$. Then

$$(5.1) \quad \sup_{[0, T] \times \mathbb{R}^2} (\varphi^* - \psi_*) = \sup_{\mathbf{y} \in \mathbb{R}^2} [\varphi^*(0, \mathbf{y}) - \psi_*(0, \mathbf{y})]$$

Suppose that $\Omega_1(t)$ and $\Omega_2(t)$ are, respectively, a χ -subsolution and a χ -supersolution of (1.1) and set

$$u_i(t, \mathbf{x}) = \chi_{\Omega_i(t)}(\mathbf{x}).$$

Since χ -sub and supersolutions are assumed to be uniformly bounded on compact time intervals, $u_i \in \hat{M}([0, \infty) \times \mathbb{R}^2)$. Then (5.1) with $\varphi = u_1$ and $\psi = u_2$ yields

¹⁸Cf. Theorem 7.1 of [CGG] for the case in which G is continuous and nonnegative.

¹⁹Ohnuma and Sato [OS] have independently established this theorem using a completely different method of proof.

$$(5.2) \quad (u_1)^*(t, \mathbf{x}) - (u_2)_*(t, \mathbf{x}) \leq \sup_{\mathbf{y} \in \mathbb{R}^2} [(u_1)^*(0, \mathbf{y}) - (u_2)_*(0, \mathbf{y})]$$

and we have

Corollary 3.1 (Weak Comparison). *Let $\Omega_1(t)$ be a χ -subsolution and $\Omega_2(t)$ a χ -supersolution of (1.1). Suppose that, for all \mathbf{x} ,*

$$(5.3) \quad (u_1)^*(0, \mathbf{x}) \leq (u_2)_*(0, \mathbf{x}).$$

Then for all $t \geq 0$,

$$(5.4) \quad \text{cl } \Omega_1(t) \subseteq \text{int } \Omega_2(t).$$

Condition (5.3) follows if (5.4) is satisfied at $t=0^+$. Unfortunately, (5.3) is stronger than the requirement: $\Omega_1(0) \subseteq \Omega_2(0)$.

We say that (1.1) with initial data Ω_0 has **strong comparison** in $(0, T)$ if

$$(5.5a) \quad \text{graph}_T \Omega_1 = \text{cl}(\text{int}(\text{graph}_T \Omega_2))$$

for all $t \in (0, T)$ for every χ -subsolution $\Omega_1(t)$ of (1.1) compatible with Ω_0 and χ -supersolution $\Omega_2(t)$ of (1.1) compatible with Ω_0 .

The next result follows from the definitions of the upper and lower envelopes $\mathcal{U}(t)$ and $\mathcal{L}(t)$ and the uniqueness time T_{uniq} for relaxed evolutions from Ω_0 .

Theorem 3.4. *Let $\Omega_1(t)$ be a χ -subsolution and $\Omega_2(t)$ a χ -supersolution, both of (1.1) and both compatible with Ω_0 . Then for all $t \geq 0$,*

$$\text{cl } \Omega_1(t) \subseteq \mathcal{U}(t), \quad \text{int } \Omega_2(t) \supseteq \mathcal{L}(t).$$

Thus (1.1) with initial data Ω_0 has strong comparison in $(0, T_{\text{uniq}})$.

4. RELATION BETWEEN CLASSICAL AND RELAXED EVOLUTIONS.

Our next theorem shows that our choice of effective moduli G and B for the relaxed problem is the only possible choice, at least if wedges are to be relaxed evolutions; what makes this result so interesting is that $G(\theta)$ and $B(\theta)$ differ from $g(\theta)$ and $b(\theta)$ only at angles θ that are not globally stable, and wedges by definition do not involve such θ .

Theorem 4.1 (Effective Moduli are Canonical). *Let f and b be consistent with (2.1), let G and B be consistent with (1.2), and let $G(\theta) = f(\theta) + f''(\theta)$ and $B(\theta) = b(\theta)$ for all GS angles θ . Then all wedges are relaxed evolutions only if G and B are the effective moduli for f and b .*

Proof. It suffices to show that $G(\theta)=0$ and $B(\theta)$ satisfies (1.17) on any GUS angle-interval (θ_1, θ_2) . Choose such an angle-interval (θ_1, θ_2) . Consider a (θ_1, θ_2) -wedge $\Omega(t)$ with corner at the origin at $t=1$, and let ω be the corresponding rigid velocity defined by (1.16). Assume that $\Omega(t)$ is a relaxed evolution, so that $u(t, \mathbf{x}) = \chi_{\Omega(t)}(\mathbf{x})$ is a viscosity solution of (3.4).

Let $\Omega(t)$ be convex, and let

$$w(t, \mathbf{x}, \theta) = 1 - [\mathbf{x} - (t-1)\omega] \cdot \mathbf{N}(\theta)$$

for all (t, \mathbf{x}) and all $\theta \in (\theta_1, \theta_2)$. Then by (2.7),

$$\Omega(t) = \{ \mathbf{x} \in \mathbb{R}^2 : w(t, \mathbf{x}, \theta) \geq 1, \theta = \theta_1, \theta_2 \}$$

Fix $\theta \in (\theta_1, \theta_2)$. Then,

$$u^*(t, \mathbf{x}) - w(t, \mathbf{x}, \theta) \leq u^*(1, 0) - w(1, 0, \theta) = 0$$

for all (t, \mathbf{x}) near $(1, 0)$. Thus, since u is a viscosity solution of (3.4),

$$w_t(1, 0, \theta) \leq \mathcal{F}^*(\nabla w(1, 0, \theta), \nabla^2 w(1, 0, \theta)).$$

Further,

$$w_t(1,0,\theta) = \omega \cdot N(\theta), \quad \nabla w(1,0,\theta) = -N(\theta), \quad \nabla^2 w(1,0,\theta) = 0,$$

and (3.5) yields

$$\mathcal{F}^*(\nabla w(1,0,\theta), \nabla^2 w(1,0,\theta)) = -U/B(\theta);$$

hence

$$(4.1) \quad B(\theta) \leq -U[\omega \cdot N(\theta)]^{-1}.$$

Now let $\Omega(t)$ be concave, and let

$$w(t,\mathbf{x}) = w(t,\mathbf{x},\theta) = -[\mathbf{x} - (t-1)\omega] \cdot N(\theta).$$

with θ fixed. Then $u_* - w$ has a local minimum at $(t,\mathbf{x}) = (1,0)$, so that, arguing as before,

$$B(\theta) \geq -U[\omega \cdot N(\theta)]^{-1}.$$

Thus, appealing to (4.1),

$$(4.2) \quad B(\theta) = -U[\omega \cdot N(\theta)]^{-1} \text{ for all } \theta \in (\theta_1, \theta_2),$$

and (1.17) follows from (1.16), (4.2), and (1.12).

Next, to show that $G(\theta) = 0$ on (θ_1, θ_2) , we again take $\Omega(t)$ to be convex, and let

$$\hat{w}(t,\mathbf{x},\theta) = 1 - [\mathbf{x} - (t-1)\omega] \cdot N(\theta) + \beta[\mathbf{x} - (t-1)\omega]^2$$

for all (t,\mathbf{x}) and all $\theta \in (\theta_1, \theta_2)$. Fix θ and write $\hat{w}(t,\mathbf{x}) = \hat{w}(t,\mathbf{x},\theta)$.

We first show that, given any $\beta \in \mathbb{R}$, $u_* - \hat{w}$ has a local maximum at $(t,\mathbf{x}) = (1,0)$. Choose (τ,\mathbf{y}) with $u^*(\tau,\mathbf{y}) = 1$. Then

$$[\mathbf{y} - (\tau - 1)\boldsymbol{\omega}] \cdot \mathbf{N}(\theta_i) \leq 0,$$

$i=1,2$, and, since $\theta \in (\theta_1, \theta_2)$,

$$[\mathbf{y} - (\tau - 1)\boldsymbol{\omega}] \cdot \mathbf{N}(\theta) \leq -\alpha < 0,$$

$\alpha = \alpha(\theta)$. Therefore if (τ, \mathbf{y}) is close enough to $(1, \mathbf{0})$ that $|\mathbf{y} - (\tau - 1)\boldsymbol{\omega}| \leq \alpha/2\beta$, then

$$\begin{aligned} 1 = u^*(\tau, \mathbf{y}) &\leq 1 + \frac{1}{2}\alpha|\mathbf{y} - (\tau - 1)\boldsymbol{\omega}| \\ &\leq 1 - [\mathbf{x} - (\tau - 1)\boldsymbol{\omega}] \cdot \mathbf{N}(\theta) + \beta[\mathbf{x} - (\tau - 1)\boldsymbol{\omega}]^2 = \hat{w}(\tau, \mathbf{y}). \end{aligned}$$

Further,

$$\hat{w}(\tau, \mathbf{y}) \geq 0 \quad \text{if} \quad |\mathbf{y} - (\tau - 1)\boldsymbol{\omega}| + \beta|\mathbf{y} - (\tau - 1)\boldsymbol{\omega}|^2 \leq 1,$$

and hence

$$u^*(\tau, \mathbf{y}) - \hat{w}(\tau, \mathbf{y}) \leq 0 = u^*(1, \mathbf{0}) - \hat{w}(1, \mathbf{0})$$

for all (τ, \mathbf{y}) sufficiently close to $(1, \mathbf{0})$; thus

$$\hat{w}_t(1, \mathbf{0}) \leq \mathcal{F}^*(\nabla \hat{w}(1, \mathbf{0}), \nabla^2 \hat{w}(1, \mathbf{0})).$$

Further,

$$\begin{aligned} \hat{w}_t(1, \mathbf{0}) &= \boldsymbol{\omega} \cdot \mathbf{N}(\theta), \quad \nabla \hat{w}(1, \mathbf{0}) = -\mathbf{N}(\theta), \quad \nabla^2 \hat{w}(1, \mathbf{0}) = 2\beta \mathbf{I}, \\ \mathcal{F}^*(\nabla \hat{w}(1, \mathbf{0}), \nabla^2 \hat{w}(1, \mathbf{0})) &= (2\beta G(\theta) - U)/B(\theta); \end{aligned}$$

hence

$$(4.3) \quad \boldsymbol{\omega} \cdot \mathbf{N}(\theta) \leq (2\beta G(\theta) - U)/B(\theta),$$

and this must hold for all $\beta \in \mathbb{R}$ and $\theta \in (\theta_1, \theta_2)$. On the other hand, by

(4.2), $\omega \cdot N(\theta) = -U/B(\theta)$, and (4.3) can hold for all β only if $G(\theta) = 0$. \square

Theorem 4.2 (Classical Evolutions are Relaxed Evolutions). *Let Ω_0 be bounded and admissible. Let G and B be the effective moduli corresponding to f and b , with f and b consistent with (2.1). Let $\Omega(t)$, $t \in [0, T_{\max})$, be the maximal classical evolution from Ω_0 . Then the uniqueness time T_{uniq} for relaxed evolutions from Ω_0 satisfies $T_{\text{uniq}} \geq T_{\max}$ and $\Omega(t)$ coincides with the unique relaxed evolution $\Omega_{\text{uniq}}(t)$ for all $t \in [0, T_{\max})$.*

Proof. Let $\Omega(t)$, $0 < t < T_{\max}$ be a classical evolution. We will show only that $\Omega(t)$ is a χ -subsolution; the proof that $\Omega(t)$ is a χ -supersolution is analogous. Let $u(t, \mathbf{x}) = \chi_{\Omega(t)}(\mathbf{x})$. Suppose that for a test function w

$$u^*(t, \mathbf{x}) - w(t, \mathbf{x}) \leq u^*(t_0, \mathbf{x}_0) - w(t_0, \mathbf{x}_0) = 0$$

for all (t, \mathbf{x}) near (t_0, \mathbf{x}_0) .

Case 1: $\mathbf{x}_0 \in \text{int } \Omega(t_0)$. Then $u^*(t, \mathbf{x}) = 1$ for all (t, \mathbf{x}) near (t_0, \mathbf{x}_0) and

$$w_t(t_0, \mathbf{x}_0) = 0, \quad \nabla w(t_0, \mathbf{x}_0) = 0, \quad \nabla^2 w(t_0, \mathbf{x}_0) \geq 0.$$

Hence

$$\mathcal{F}^*(\nabla w(t_0, \mathbf{x}_0), \nabla^2 w(t_0, \mathbf{x}_0)) \geq 0$$

and (3.7) is satisfied.

Case 2: $\mathbf{x}_0 \in \mathbb{R}^2 \setminus \overline{\Omega}(t_0)$. Then $u^*(t, \mathbf{x}) = 0$ for all (t, \mathbf{x}) near (t_0, \mathbf{x}_0) and an analysis similar to that of Case 1 yields (3.7).

Case 3: $\mathbf{x}_0 \in \partial\Omega(t_0)$ and $\nabla w(t_0, \mathbf{x}_0) = 0$. Then $w_t(t_0, \mathbf{x}_0) = 0$, since the normal velocity V of $\partial\Omega(t)$ is finite. Moreover, the definition of the upper semicontinuous envelope yields

$$\mathcal{F}^*(\nabla w(t_0, \mathbf{x}_0), \nabla^2 w(t_0, \mathbf{x}_0)) = G(\theta) B(\theta)^{-1} \max \{ \mathbf{q} \cdot \nabla^2 w(t_0, \mathbf{x}_0) \mathbf{q} : |\mathbf{q}|=1 \}.$$

We claim that the quantity $\max \{ \dots \}$ is nonnegative. Indeed, $\mathbf{x}_0 \in \partial\Omega(t_0)$ and $t_0 < T_{\max}$; hence \mathbf{x}_0 is not an isolated point of $\Omega(t_0)$ and there is a sequence $\{\mathbf{x}_n\}$ with $\mathbf{x}_n \in \partial\Omega(t_0)$, $\mathbf{x}_n \neq \mathbf{x}_0$, and $\mathbf{x}_n \rightarrow \mathbf{x}_0$. By choosing a subsequence, if necessary, we may assume that $(\mathbf{x}_n - \mathbf{x}_0) / |\mathbf{x}_n - \mathbf{x}_0|$ is convergent, say to \mathbf{e} . Then

$$\begin{aligned} w(t_0, \mathbf{x}_n) &\geq u^*(t_0, \mathbf{x}_n) = 1 = w(t_0, \mathbf{x}_0), \\ \mathbf{e} \cdot \nabla^2 w(t_0, \mathbf{x}_0) \mathbf{e} &= 2 \lim_{n \rightarrow \infty} [w(t_0, \mathbf{x}_n) - w(t_0, \mathbf{x}_0)] / |\mathbf{x}_n - \mathbf{x}_0|^2, \end{aligned}$$

so that

$$0 = w_t(t_0, \mathbf{x}_0) \leq \mathcal{F}^*(\nabla w(t_0, \mathbf{x}_0), \nabla^2 w(t_0, \mathbf{x}_0)).$$

Case 4: \mathbf{x}_0 belongs to a smooth part of $\partial\Omega(t_0)$ and $|\nabla w(t_0, \mathbf{x}_0)| \neq 0$. Then the normal angle θ , the curvature K , and the normal velocity V of $\partial\Omega(t)$ at $t=t_0$ and \mathbf{x}_0 satisfy

$$\begin{aligned} N(\theta) &= -\nabla w(t_0, \mathbf{x}_0) / |\nabla w(t_0, \mathbf{x}_0)|, \\ K &\leq \operatorname{div}[\nabla w / |\nabla w|](t_0, \mathbf{x}_0), \\ V &= w_t(t_0, \mathbf{x}_0) / |\nabla w(t_0, \mathbf{x}_0)|, \end{aligned}$$

and (3.7) follows from (1.5).

Case 5: \mathbf{x}_0 is a corner point of $\partial\Omega(t_0)$ and $|\nabla w(t_0, \mathbf{x}_0)| \neq 0$. Let (θ_1, θ_2) be the GUS angle-interval that defines the corner, and let $\mathbf{z}(t)$ with $\mathbf{z}(t_0) = \mathbf{x}_0$ denote the trajectory of the corner for t near t_0 . Then $\partial\Omega(t)$ must have a "convex-type" corner of the type shown in Figure 4.1 near $\mathbf{z}(t)$, with curvature $K(\mathbf{x}, t) \leq 0$ for \mathbf{x} near but not equal to $\mathbf{z}(t)$. Thus and by (1.5), for such \mathbf{x} the normal velocity of $\partial\Omega(t)$ must satisfy

$$(4.4) \quad V(\mathbf{x}, t) \leq -U/b(\theta(\mathbf{x}, t)).$$

On the other hand, consider the simple (θ_1, θ_2) -wedge $\partial\Lambda(t)$ which has $\Lambda(t)$ convex and has corner at \mathbf{x}_0 at time t_0 , and let ω be the rigid velocity of the wedge as defined by (1.16). Then, since this wedge must have normal velocity $V = -U/b(\theta_i)$ on each of its facets, we may conclude from (4.4) that there is a ball \mathcal{B} centered at \mathbf{x}_0 such that $\Lambda(t) \cap \mathcal{B} \subseteq \Omega(t) \cap \mathcal{B}$ for all t near t_0 with $t \leq t_0$; thus

$$\Lambda(t) \cap \mathcal{B} \subseteq \{ \mathbf{x} : w(t, \mathbf{x}) \geq 1 \}.$$

Further, since $\Lambda(t)$ moves with rigid velocity ω and \mathbf{x}_0 is the the corner point of $\Lambda(t_0)$,

$$\mathbf{x}_0 + (t - t_0)[\omega \cdot \mathbf{N}(\alpha)]\mathbf{N}(\alpha) \in \Lambda(t)$$

for all t and all $\alpha \in [\theta_1, \theta_2]$. Thus for such α and for t near t_0 with $t \leq t_0$,

$$w(t, \mathbf{x}_0 + (t - t_0)[\omega \cdot \mathbf{N}(\alpha)]\mathbf{N}(\alpha)) \geq 1 = w(t_0, \mathbf{x}_0)$$

and it follows that

$$(4.5) \quad w_t(t_0, \mathbf{x}_0) + [\omega \cdot \mathbf{N}(\alpha)][\nabla w(t_0, \mathbf{x}_0) \cdot \mathbf{N}(\alpha)] \leq 0$$

for all $\alpha \in [\theta_1, \theta_2]$.

Now let $\alpha \in [\theta_1, \theta_2]$ be the angle defined by

$$\nabla w(t_0, \mathbf{x}_0) / |\nabla w(t_0, \mathbf{x}_0)| = -\mathbf{N}(\alpha);$$

then (4.5) yields

$$(4.6) \quad w_t(t_0, \mathbf{x}_0) \leq \omega \cdot \mathbf{N}(\alpha) |\nabla w(t_0, \mathbf{x}_0)|,$$

and, by (4.2),

$$(4.7) \quad w_t(t_0, \mathbf{x}_0) \leq -U|\nabla w(t_0, \mathbf{x}_0)|/B(\alpha).$$

If $\alpha \in (\theta_1, \theta_2)$, then $G(\alpha) = 0$ and (3.5) yields

$$(4.8) \quad \mathcal{F}^*(\nabla w(t_0, \mathbf{x}_0), \nabla^2 w(t_0, \mathbf{x}_0)) = -U|\nabla w(t_0, \mathbf{x}_0)|/B(\alpha),$$

so that, by (4.7), (3.7) is satisfied. If α equals θ_1 or θ_2 , then $G(\alpha)$ is generally nonzero, but the definition of \mathcal{F}^* yields

$$\mathcal{F}^*(\mathbf{p}, \mathbf{A}) \geq \limsup_{n \rightarrow \infty} \mathcal{F}^*(\mathbf{p}_n, \mathbf{A})$$

for any sequence $\mathbf{p}_n \rightarrow \mathbf{p}$. Let $\mathbf{p} = \nabla w(t_0, \mathbf{x}_0)$ and choose a sequence so that $G(\theta_n) = 0$ for all n , where θ_n is defined by $N(\theta_n) = \mathbf{p}_n / |\mathbf{p}_n|$. Then (4.8) is replaced by

$$(4.9) \quad \mathcal{F}^*(\nabla w(t_0, \mathbf{x}_0), \nabla^2 w(t_0, \mathbf{x}_0)) \geq -U|\nabla w(t_0, \mathbf{x}_0)|/B(\alpha),$$

which, with (4.7), yields (3.7).

We have only to show that

$$(4.10) \quad T_{\text{uniq}} \geq T_{\text{max}}.$$

Given Ω_0 we can construct a one-parameter family of *admissible* initial domains $\Omega_0(\delta)$ ($|\delta| \leq \delta_0$ for some $\delta_0 > 0$) satisfying

$$(4.11a) \quad \Omega_0(\delta) \subset \Omega_0(\delta') \quad \text{if } \delta > \delta',$$

$$(4.11b) \quad \lim_{\delta' \rightarrow \delta} \Omega_0(\delta') = \Omega_0(\delta),$$

the limit being in the *Hausdorff metric*.²⁰ Let $\Omega(t; \delta)$, ($t \in [0, T_{\text{max}}(\delta))$, $|\delta| \leq \delta_0$) be the unique maximal evolution from the initial data $\Omega_0(\delta)$ (cf. Theorem 2.1). Since for $t \in (0, T_{\text{max}}(\delta))$, $\partial\Omega(t; \delta)$ is piecewise smooth, we may use the compactness lemma [AG2, Lemma 8.3] and the uniqueness of

²⁰For bounded sets the Hausdorff metric $d_H(A, B)$ is the largest of the distances $\sup\{\text{dist}(\mathbf{x}, A) : \mathbf{x} \in B\}$ and $\sup\{\text{dist}(\mathbf{x}, B) : \mathbf{x} \in A\}$.

classical solutions to show that: (i) $T_{\max}(\delta)$ is lower semicontinuous in δ (cf. also the proof of [AG2, Lemma 8.2]); (ii) by Corollary 3.1, $\Omega(t;\delta) \subset \Omega(t;\delta')$ if $\delta \geq \delta'$; (iii) for fixed t , the map $\delta \mapsto \Omega(t;\delta)$ is continuous in the Hausdorff metric. (Assertion (iii) is proved by showing that any limit point of $\Omega(t;\delta)$ as $\delta \rightarrow \delta'$ is a classical solution with initial data $\Omega_0(\delta')$ and hence by uniqueness is equal to $\Omega(t;\delta')$.

Fix $T < T_{\max} = T_{\max}(0)$. By the lower semicontinuity of $T_{\max}(\delta)$ there is a $\delta(T) > 0$ satisfying

$$T_{\max}(\delta) \geq T \quad \text{for all } |\delta| \leq \delta(T).$$

For $(t, \mathbf{x}) \in [0, T] \times \mathbb{R}^2$ define

$$\begin{aligned} \Phi(t, \mathbf{x}) &= \inf \{ \delta : |\delta| \leq \delta(T), \mathbf{x} \in \Omega(t; \delta) \}, \\ &= -\delta(T) \quad \text{if the set above is empty.} \end{aligned}$$

Since $\Omega(t; \delta) \subseteq \Omega(t; \delta')$ for $\delta \geq \delta'$,

$$\{ \mathbf{x} : \Phi(t, \mathbf{x}) \geq \delta \} = \Omega(t; \delta)$$

whenever $|\delta| \leq \delta(T)$ and $t \in [0, T]$. Moreover, for each δ , $\Omega(t; \delta)$ is a classical and therefore relaxed evolution from $\Omega_0(\delta)$. From this one can show that Φ is a continuous viscosity solution of (1.1), so that, by Theorem 3.2,

$$\mathcal{U}(t) = \{ \mathbf{x} : \Phi(t, \mathbf{x}) \geq 0 \} = \Omega(t; 0) = \Omega(t), \quad \mathcal{L}(t) = \{ \mathbf{x} : \Phi(t, \mathbf{x}) > 0 \}.$$

Since

$$\lim_{\delta \downarrow 0} \Omega(t; \delta) = \Omega(t; 0)$$

for every $\mathbf{x} \in \mathcal{U}(t)$, there are $\delta_n > 0$ and $\mathbf{x}_n \rightarrow \mathbf{x}$ such that $\mathbf{x}_n \in \Omega(t; \delta_n) \subset \mathcal{L}(t)$. Hence $\text{cl } \mathcal{L}(t) = \mathcal{U}(t)$ for all $t \in [0, T]$. An analogous

argument shows that $\text{int}\mathcal{U}(t)=\mathcal{L}(t)$ at each $t\in[0,T]$. Hence $T_{\text{uniq}}\geq T$, and the desired conclusion follows, since $T < T_{\text{max}}$ was chosen arbitrarily. \square

Remark 4.1. For bounded, admissible initial data Ω_0 there exist a maximal existence time T_{max} and a classical evolution $\hat{\Omega}(t)$, $t\in[0,T_{\text{max}})$, from Ω_0 . There is also (at least one) relaxed evolution $\Omega(t)$, $t\in[0,\infty)$, from Ω_0 , and, by Theorem 4.2,

$$\hat{\Omega}(t) = \Omega(t) \text{ for all } t\in[0,T_{\text{max}}).$$

Hence the relaxed evolution represents a weak extension of the classical evolution $\hat{\Omega}(t)$ after $\hat{\Omega}(t)$ develops a singularity at $t = T_{\text{max}}$.

5. CONVERGENCE.

Throughout this section

G and B satisfy (1.2);

Ω_0 is a prescribed initial domain, assumed compact;

T_{uniq} is the uniqueness time for relaxed evolutions from Ω_0 .

a. GENERAL RESULTS.

We say that a sequence $\{\Omega_0^n\}$ of compact domains approximates Ω_0 provided the signed distance to Ω_0^n approaches the signed distance to Ω_0 , uniformly on \mathbb{R}^2 .

Theorem 5.1 (Convergence of Relaxed Evolutions). Assume that $T_{\text{uniq}} > 0$. Let $\{\Omega_0^n\}$ approximate Ω_0 . For each integer n , let $\Omega^n(t)$, $t \in [0, \infty)$, be a relaxed evolution from Ω_0^n . Then, for each $t \in [0, T_{\text{uniq}})$, $\Omega^n(t)$ converges, in the Hausdorff topology, to the unique relaxed evolution from Ω_0 .

We now state a result that holds for all time. The proof will be given at the end of this section, as will the proof of the theorem just stated.

Theorem 5.2 (Convergence of Level-Set Solutions). Let Φ be the unique level-set solution corresponding to an auxiliary function Φ_0 for Ω_0 . Let $\{\Phi_n\}$ be a sequence of level-set solutions of (1.1) such that

$$\lim_{n \rightarrow \infty} \Phi_n(0, \mathbf{x}) = \Phi_0(\mathbf{x})$$

uniformly on \mathbb{R}^2 . Then

$$\lim_{n \rightarrow \infty} \Phi_n(t, \mathbf{x}) = \Phi(t, \mathbf{x})$$

uniformly on compact subsets of $[0, \infty) \times \mathbb{R}^2$.

b. INFINITESIMALLY WRINKLED SOLUTIONS AS LIMITS OF SOLUTIONS FROM ADMISSIBLE INITIAL DOMAINS.

If the initial domain Ω_0 is *admissible*, then there is a classical evolution from Ω_0 up to a maximal existence time T_{\max} ; each relaxed evolution from Ω_0 (unique up to $T_{\text{uniq}} \geq T_{\max}$) supplies a weak extension of this classical solution for times greater than T_{\max} .

Suppose that Ω_0 is *not admissible* (for example, suppose that $\partial\Omega_0$ has normal angles θ for which $g(\theta) < 0$). Then the notion of a classical evolution from Ω_0 breaks down, since classical evolutions are required to be admissible and hence to have globally stable normal-angles. On the other hand, there is a relaxed evolution from Ω_0 . The derivation of the relaxed formulation is based on allowing the boundary curve to develop infinitesimal wrinkles whenever its normal angle is not globally stable. We now use Theorem 6.1 to give a partial justification of this procedure, under the assumption that $\partial\Omega_0$ is C^1 and piecewise C^2 , and that $T_{\text{uniq}} > 0$.

We first approximate Ω_0 by a sequence $\{\Omega_0^n\}$ of *admissible* bounded domains. We accomplish this by dividing $\partial\Omega_0$ into curves whose normal angles are GS, interspaced with curves whose normal angles are GUS. We approximate $\partial\Omega_0$ by leaving the GS curves unchanged, but replacing each GUS curve by a wrinkled curve. If Γ is such a GUS curve, then the normal angles of Γ lie in a GUS angle-interval (θ_1, θ_2) with θ_1 and θ_2 angles for a corner consistent with (1.11). We replace Γ by a wrinkled curve \mathcal{W} such that: the endpoints of \mathcal{W} coincide with those of Γ ; the facet angles of \mathcal{W} are θ_1 and θ_2 ; \mathcal{W} lies in an arbitrary small neighborhood of Γ . The replacement for $\partial\Omega_0$ constructed in this manner is admissible and arbitrarily close to $\partial\Omega_0$ in the required sense.

For each n , we let $\Omega^n(t)$, $t \in [0, \infty)$, be a relaxed evolution from the admissible initial domain Ω_0^n . Then, by Theorem 6.1, for each $t \in [0, T_{\text{uniq}})$, $\Omega^n(t)$ converges, in the Hausdorff topology, to the unique relaxed evolution from Ω_0 .

c. PROOFS.

Proof of Theorem 5.1.

1° Let

$$\Phi_0(\mathbf{x}) = \begin{cases} \text{dist}(\mathbf{x}, \partial\Omega_0) \wedge 1, & \mathbf{x} \in \Omega_0, \\ -(\text{dist}(\mathbf{x}, \partial\Omega_0) \wedge 1), & \mathbf{x} \notin \Omega_0. \end{cases}$$

Then Φ_0 is an auxiliary function for Ω_0 as defined in Section 3, and there is a unique level-set solution Φ of (1.1) corresponding to Φ_0 . For each n , let $\Phi_{0n}(\mathbf{x})$ and $\Phi_n(t, \mathbf{x})$ be defined in the same manner using Ω_0^n as the initial set. Since $\{\Omega_0^n\}$ approximates Ω_0 , $\Phi_{0n}(\mathbf{x})$ converges to $\Phi_0(\mathbf{x})$, uniformly for $\mathbf{x} \in \mathbb{R}^2$. Thus Theorem 5.2, which will be proved subsequently, implies that $\Phi_n(t, \mathbf{x})$ converges to $\Phi(t, \mathbf{x})$, uniformly on compact subsets of $[0, \infty) \times \mathbb{R}^2$.

2° Let $\mathcal{B}(r)$ denote the (closed) ball of radius r centered at the origin. Since Ω_0 is compact and $\{\Omega_0^n\}$ approximates Ω_0 , there is an R_0 such that $\Omega_0^n \subset \mathcal{B}(R_0)$ for all n . Set

$$\mu = |U|(\inf \mathcal{B}(\theta))^{-1}.$$

Then

$$\mathcal{B}_0(t) = \mathcal{B}(R_0 + \mu t)$$

is a χ -supersolution of (1.1) compatible with Ω_0 , and hence

$$\Psi(t, \mathbf{x}) = \begin{cases} -1, & \mathbf{x} \notin \text{int} \mathcal{B}_0(t), \\ 1, & \mathbf{x} \in \text{int} \mathcal{B}_0(t). \end{cases}$$

is a viscosity supersolution of (3.4) with $\Psi(0, \mathbf{x}) \geq \Phi_n(0, \mathbf{x})$. Thus Theorem 3.3 with Ψ as supersolution and Φ_n as subsolution yields $\Psi \geq \Phi_n$. In particular,

$$\Phi_n(t, \mathbf{x}) \leq -1, \quad \mathbf{x} \notin \text{int} \mathcal{B}_0(t).$$

3° By Theorem 3.4,

$$(5.1a) \quad \{ \mathbf{x} : \Phi_n(t, \mathbf{x}) > 0 \} \subseteq \Omega^n(t) \subseteq \{ \mathbf{x} : \Phi_n(t, \mathbf{x}) \geq 0 \}$$

for $t \geq 0$. Therefore

$$(5.1b) \quad \Omega^n(t) \subseteq \mathcal{B}_0(t)$$

for $t \geq 0$. Also, on $[0, T_{\text{uniq}})$ the unique relaxed evolution from Ω_0 is given by

$$(5.2) \quad \Omega(t) = \{ \mathbf{x} : \Phi(t, \mathbf{x}) \geq 0 \}.$$

4° For $\delta > 0$, let

$$\mathcal{U}(t; \delta) = \{ \mathbf{x} : \Phi(t, \mathbf{x}) \geq -\delta \}, \quad \mathcal{L}(t; \delta) = \{ \mathbf{x} : \Phi(t, \mathbf{x}) > \delta \}.$$

Since $\Phi_n(t, \mathbf{x})$ converges to $\Phi(t, \mathbf{x})$ locally uniformly, we may use (5.1b) to conclude that there is an $n(\delta)$ such that, for all $n \geq n(\delta)$ and $t \in [0, T_{\text{uniq}})$,

$$\mathcal{U}(t; \delta) \supseteq \{ \mathbf{x} : \Phi_n(t, \mathbf{x}) \geq 0 \}, \quad \mathcal{L}(t; \delta) \subseteq \{ \mathbf{x} : \Phi_n(t, \mathbf{x}) > 0 \}.$$

Hence (5.1a) yields

$$(5.3) \quad \mathcal{L}(t; \delta) \subseteq \Omega^n(t) \subseteq \mathcal{U}(t; \delta)$$

for all $n \geq n(\delta)$ and $t \in [0, T_{\text{uniq}})$.

5° Using the arguments of step 2, we can show that

$$\mathcal{U}(t; \delta), \mathcal{L}(t; \delta) \subseteq \mathcal{B}_0(t).$$

6° Our next step will be to show that, for every $t \in [0, T_{\text{uniq}})$, the Hausdorff distance

$$d_\delta = d_H(\mathcal{U}(t;\delta), \mathcal{L}(t;\delta))$$

satisfies

$$(5.4) \quad d_\delta \rightarrow 0 \quad \text{as} \quad \delta \rightarrow 0.$$

Since $\mathcal{L}(t;\delta) \subseteq \mathcal{U}(t;\delta)$, we may use the definitions of $\mathcal{L}(t;\delta)$ and $\mathcal{U}(t;\delta)$ to conclude that

$$\begin{aligned} d_\delta &= \sup\{d(t, \mathbf{x}; \delta) : \Phi(t, \mathbf{x}) \geq -\delta\}, \\ d(t, \mathbf{x}; \delta) &= \inf\{|\mathbf{x} - \mathbf{y}| : \Phi(t, \mathbf{x}) \geq \delta\}. \end{aligned}$$

Choose $\mathbf{x}(\delta)$ satisfying

$$(5.5) \quad \begin{aligned} \Phi(t, \mathbf{x}(\delta)) &\geq -\delta, \\ d(t, \mathbf{x}(\delta); \delta) &\geq d_\delta - \delta. \end{aligned}$$

Since $\mathbf{x}(\delta) \in \mathcal{U}(t;\delta) \subset \mathcal{B}_0(t)$ and $\mathcal{B}_0(t)$ is compact, there is a sequence (also denoted by δ) such that $\mathbf{x}(\delta) \rightarrow \mathbf{x}_0$ as $\delta \downarrow 0$; hence

$$\Phi(t, \mathbf{x}_0) = \lim_{\delta \downarrow 0} \Phi(t, \mathbf{x}(\delta)) \geq 0,$$

and $\mathbf{x}_0 \in \mathcal{U}(t)$. Also, $t \in [0, T_{\text{uniq}})$; hence we may conclude from the definition of T_{uniq} that $\mathcal{U}(t) = \text{cl } \mathcal{L}(t)$, and there is a sequence $\mathbf{y}_m \rightarrow \mathbf{x}_0$, $\mathbf{y}_m \in \mathcal{L}(t)$, or equivalently, $\Phi(t, \mathbf{y}_m) > 0$. Thus, for all $\delta < \Phi(t, \mathbf{y}_m)$,

$$d(t, \mathbf{x}(\delta); \delta) \leq |\mathbf{x}(\delta) - \mathbf{y}_m|.$$

Now let δ tend to zero and then m to infinity to obtain

$$\lim_{\delta \downarrow 0} d(t, \mathbf{x}(\delta); \delta) = 0,$$

and this, with (5.5), implies (5.4).

7° By (5.2) and (5.3),

$$d_H(\Omega^n(t), \Omega(t)) \leq d_H(\mathcal{U}(t; \delta), \mathcal{L}(t; \delta)) = d_\delta$$

for every $t \in [0, T_{\text{uniq}})$, $\delta > 0$, and $n > n(\delta)$. Therefore, by (5.3),

$$\lim_{n \rightarrow \infty} d_H(\Omega^n(t), \Omega(t)) = 0$$

for all $t \in [0, T_{\text{uniq}})$. \square

Proof of Theorem 5.2.

1° Since Φ_0 is bounded and $\Phi_n(0, \mathbf{x})$ converges uniformly to $\Phi_0(\mathbf{x})$, there is a $\kappa > 0$ such that

$$(5.6) \quad |\Phi_n(0, \mathbf{x})| \leq \kappa$$

for all $\mathbf{x} \in \mathbb{R}^2$. Since $\psi \equiv \kappa$ is a solution of (3.4), the inequality (5.6) and the comparison theorem 3.3 yield $\Phi_n(t, \mathbf{x}) \leq \kappa$. Similarly, $\psi \equiv -\kappa$ yields $\Phi_n(t, \mathbf{x}) \geq -\kappa$. Hence

$$(5.7) \quad |\Phi_n(t, \mathbf{x})| \leq \kappa$$

for all $(t, \mathbf{x}) \in [0, \infty) \times \mathbb{R}^2$.

2° For $(t, \mathbf{x}) \in [0, \infty) \times \mathbb{R}^2$, define

$$\Phi^+(t, \mathbf{x}) = \limsup_{\substack{n \rightarrow \infty \\ (s, \mathbf{y}) \rightarrow (t, \mathbf{x})}} \Phi_n(s, \mathbf{y}),$$

$$\Phi^-(t, \mathbf{x}) = \liminf_{\substack{n \rightarrow \infty \\ (s, \mathbf{y}) \rightarrow (t, \mathbf{x})}} \Phi_n(s, \mathbf{y}).$$

Then Φ^+ is a viscosity subsolution and Φ^- a viscosity supersolution of (3.4) in $(0, \infty) \times \mathbb{R}^2$ (cf. [FS; §2.6, §7.4]).

3° Theorem 3.3 applied to the subsolution Φ^+ and the supersolution Φ^- yields

$$\Phi^+(t, \mathbf{x}) - \Phi^-(t, \mathbf{x}) \leq \sup_{\mathbf{y}} [\Phi^+(0, \mathbf{x}) - \Phi^-(0, \mathbf{x})].$$

Note that Φ_n is locally uniformly convergent if and only if $\Phi^+ = \Phi^-$. Also, by construction, $\Phi^+ \geq \Phi_0 \geq \Phi^-$. Hence to prove local uniform convergence of Φ_n it suffices to show that

$$(5.8) \quad \Phi^+(0, \mathbf{x}) = \Phi_0(\mathbf{x}) = \Phi^-(0, \mathbf{x}),$$

which we shall accomplish in the next three steps.

4° Let

$$g^* = \sup \{ G(\theta) : \theta \in [0, 2\pi) \}, \quad \alpha = |U|.$$

For $\mathbf{x} \in \mathbb{R}^2$ and $\delta > 0$, define (cf. (5.6))

$$\Psi(t, \mathbf{x}; \mathbf{y}, \delta) = \begin{cases} 0, & |\mathbf{x} - \mathbf{y}| \leq R(t; \delta), \\ -2\kappa, & |\mathbf{x} - \mathbf{y}| > R(t; \delta), \end{cases}$$

where $R(t; \delta)$ is a solution of

$$\begin{aligned} dR(t; \delta)/dt &= -g^* R(t; \delta)^{-1} - \alpha, & t \in (0, T(\delta)), \\ R(0; \delta) &= \delta, \end{aligned}$$

with $T(\delta) < +\infty$ the first time t for which $R(t; \delta) = 0$. Then

$$\{ \mathbf{x} : |\mathbf{x} - \mathbf{y}| \leq R(t; \delta) \}, \quad t \in (0, T(\delta))$$

is a classical subsolution of the relaxed equation (1.1), and hence Ψ is a

viscosity subsolution of (3.4) on $(0, T(\delta)) \times \mathbb{R}^2$.

5° Fix $\mathbf{y} \in \mathbb{R}^2$ and let $\beta_0 = \Phi_0(\mathbf{y})$. Then, for all $\beta < \beta_0$, there are $\delta > 0$ and η_0 such that

$$\Psi(0, \mathbf{x}; \mathbf{y}, \delta) + \beta \leq \Phi_n(0, \mathbf{x})$$

for all $\mathbf{x} \in \mathbb{R}^2$ and $n \geq \eta_0$. Since Ψ is a viscosity subsolution of (3.4), it is clear from the form of this equation that $\Psi + \beta$ is also a viscosity subsolution of (3.4).

6° We now use Theorem 3.3 with subsolution $\Psi + \beta$ and supersolution Φ_n to obtain

$$\Psi(t, \mathbf{x}; \mathbf{y}, \delta) + \beta \leq \Phi_n(t, \mathbf{x})$$

for all $(t, \mathbf{x}) \in [0, T(\delta)) \times \mathbb{R}^2$; hence

$$\Psi(t, \mathbf{x}; \mathbf{y}, \delta) + \beta \leq \Phi^-(t, \mathbf{x})$$

for all $(t, \mathbf{x}) \in [0, T(\delta)) \times \mathbb{R}^2$. Applying this inequality at $(t, \mathbf{x}) = (0, \mathbf{y})$ yields $\beta \leq \Phi^-(0, \mathbf{y})$ for all $\beta < \beta_0 = \Phi_0(\mathbf{y})$. Therefore $\Phi_0(\mathbf{y}) = \Phi^-(0, \mathbf{y})$.

7° To show that $\Phi_0(\mathbf{y}) = \Phi^+(0, \mathbf{y})$, we follow the procedure of the three previous steps replacing Ψ with the supersolution

$$\hat{\Psi}(t, \mathbf{x}; \mathbf{y}, \delta) = \begin{cases} 0, & |\mathbf{x} - \mathbf{y}| \leq R(t; \delta), \\ 2\kappa, & |\mathbf{x} - \mathbf{y}| > R(t; \delta), \end{cases}$$

of (3.4). \square

6. LARGE-TIME ASYMPTOTICS.

In this section we discuss the large-time asymptotics of relaxed evolutions, assuming throughout that:

G and B satisfy (1.2) and (1.3);

Ω_0 is a prescribed initial domain, assumed compact.

In particular, we will prove that, for $U < 0$ and Ω_0 large enough, $t^{-1}\Omega(t)$ converges to a dilation of the Wulff region for $1/B(\theta)$. This result, conjectured by Angenent and Gurtin [AG1], was proved by Soner [So] for $G > 0$ and B with a convex polar diagram, and extended in [AG2] to general $B > 0$. We here follow the ideas of [So, §12-13].

Let $\Omega(t) \subset \mathbb{R}^2$, $t \geq 0$, be given. Then $\Omega(t)$ **vanishes in finite time** if there is a $T > 0$ such that

$$\Omega(t) = \emptyset \text{ for all } t > T.$$

Given a function $\varphi > 0$ on $(0, \infty)$ and a set $A \subset \mathbb{R}^2$, we write

$$\Omega(t) \sim \varphi(t)A \text{ as } t \rightarrow \infty$$

if there are functions $\varphi_1, \varphi_2 > 0$ on $(0, \infty)$ such that

$$\varphi_1(t)A \subset \Omega(t) \subset \varphi_2(t)A$$

for all sufficiently large t , and

$$\varphi_i(t)/\varphi(t) \rightarrow 1 \text{ as } t \rightarrow \infty \quad (i=1,2).$$

The **Wulff region** $W(h)$ for a given function $h(\theta)$ (cf., e.g., [G2]) is the set

$$W(h) = \{ \mathbf{x} \in \mathbb{R}^2 : \mathbf{x} \cdot \mathbf{N}(\theta) \leq h(\theta), \quad \theta \in [0, 2\pi] \}.$$

Our main result of this section is

Theorem 6.1 (Asymptotic behavior of relaxed evolutions). *Let $\Omega(t)$ be a relaxed evolution from Ω_0 .*

(a) *If $U > 0$, then $\Omega(t)$ vanishes in finite time.*

(b) *If $U < 0$ with $|U|$ sufficiently large, then*

$$\Omega(t) \sim t|U|W(1/B) \quad \text{as } t \rightarrow \infty.$$

Assertion (a) is a direct consequence of

Lemma 6.1. *Let $\Omega(t)$ be a χ -subsolution of (1.1) compatible with Ω_0 . Choose α_0 such that*

$$(6.1) \quad \text{int } \Omega_0 \subset \alpha_0 W(1/B).$$

Then, for $t > 0$,

$$(6.2) \quad \Omega(t) \subset (-Ut + \alpha_0)W(1/B).$$

Proof. The right side of (6.2), denoted by $\Lambda(t)$, is a χ -solution of

$$B(\theta)V = -U$$

for $t > 0$ [So, §12]. Since $W(1/B)$ is convex, $\Lambda(t)$ has curvature ≤ 0 . Thus, since $G \geq 0$, $\Lambda(t)$ is a χ -supersolution of (1.1); (6.2) then follows from (6.1) and Corollary 3.1. \square

Assertion (b) is more difficult to prove; for that reason we first give a simple proof under a more stringent hypothesis on B . To state this hypothesis, let \mathcal{D} denote the differential operator defined on functions $H(\theta)$ by

$$\mathcal{D}H = H + H''.$$

Then the polar diagram of H is convex at angles θ for which $\mathcal{D}H(\theta) \geq 0$, strictly convex at angles with $\mathcal{D}H(\theta) > 0$. We now establish (b) under the assumption that, for some constant $C > 0$,

$$(6.3) \quad G \leq C\mathcal{D}(1/B)$$

on $[0, 2\pi]$, so that the polar diagram of $1/B$ is convex; in fact, strictly convex at angles θ with $G(\theta) > 0$. Granted (6.3), (b) follows from Lemma 6.1 and

Lemma 6.2. *Assume that (6.3) is satisfied, and that $U < 0$ and sufficiently large that*

$$(6.4) \quad \alpha_0 \text{ int} W(1/B) \subset \Omega_0, \quad \alpha_0 = -2C/U.$$

Let $\alpha(t)$ be the solution of

$$\alpha'(t) = -U - C/\alpha(t), \quad \alpha(0) = \alpha_0.$$

Then any relaxed evolution $\Omega(t)$ compatible with Ω_0 satisfies, for $t > 0$,

$$(6.5) \quad \alpha(t)W(1/B) \subset \Omega(t).$$

Proof. Let $\Lambda(t) = \alpha(t)W(1/B)$. Then $\Lambda(t)$ is a χ -solution of (1.1) with G replaced by $C\mathcal{D}(1/B)$ [So §12]. Since $W(1/B)$ is smooth and $0 \leq G \leq C\mathcal{D}(1/B)$, $\Lambda(t)$ is a χ -subsolution of (1.1) (with G); hence (6.5) follows from (6.4) and Corollary 3.1. \square

Proof of Theorem 6.1.

1° Let

$$g = \sup G(\theta), \quad b = \sup B(\theta),$$

and assume that $|U|$ is sufficiently large that

$$\alpha_0 \text{ int } \mathcal{B}_1 \subset \Omega_0, \quad \alpha_0 = -2g/U,$$

where \mathcal{B}_1 is the unit ball in \mathbb{R}^2 . Let $\alpha(t)$ be the solution of

$$b\alpha'(t) = -U - g/\alpha(t), \quad \alpha(0) = \alpha_0,$$

and let $\Lambda(t) = \alpha(t)\mathcal{B}_1$. Then $\Lambda(t)$ is a classical solution of the isotropic equation $bV = gK - U$. Also, $V > 0$, since $\alpha'(t) > 0$ for all $t > 0$; consequently, $\Lambda(t)$ is a χ -subsolution of (1.1) compatible with Ω_0 , and, by Corollary 3.1, $\Lambda(t) \subset \Omega(t)$. Further, $\alpha(t) \rightarrow \infty$ as $t \rightarrow \infty$; hence:

$$(6.6) \quad \text{as } t \rightarrow \infty, \quad \Omega(t) \text{ expands to fill the entire space.}$$

2° Let $W_n \subset \mathbb{R}^2$ be a sequence of strictly convex, closed domains, with smooth boundary, satisfying

$$(6.7) \quad (1 - n^{-1})W(1/B) \subset W_n \subset W(1/B).$$

Further, let $\gamma_n(\theta)$ denote the support function of W_n :

$$\gamma_n(\theta) = \sup \{ \mathbf{x} \cdot \mathbf{N}(\theta) : \mathbf{x} \in W_n \}$$

(so that $W_n = W(\gamma_n)$). Since W_n is strictly convex and ∂W_n is smooth, γ_n is smooth and $\mathcal{D}\gamma_n > 0$. Also,

$$1/B(\theta) \geq \sup \{ \mathbf{x} \cdot \mathbf{N}(\theta) : \mathbf{x} \in W(1/B) \},$$

and hence, by (6.7),

$$(6.8) \quad 1/B \geq \gamma_n.$$

3° Let

$$c_n = g \left\{ \inf_{\theta} \mathcal{D}\gamma_n(\theta) \right\}^{-1}.$$

By 2°, $c_n < \infty$, but c_n may diverge to $+\infty$ as $n \rightarrow \infty$.

Choose $t_n > 0$ satisfying

$$(6.9) \quad \alpha_n = \text{int } W_n \subset \Omega(t_n), \quad \alpha_n = -2c_n/U,$$

and let $\alpha_n(t)$ be the solution of

$$\begin{aligned} \alpha_n'(t) &= -U - c_n/\alpha_n(t), \quad t > t_n, \\ \alpha_n(t_n) &= \alpha_n. \end{aligned}$$

Then $\Lambda_n(t) = \alpha_n(t)W_n$ is a χ -solution of

$$\gamma_n(\theta)^{-1}V = c_n \mathcal{D}\gamma_n(\theta)K - U$$

for $t > t_n$ [So, §12]. Further, (6.8), the definition of c_n , the convexity of $\Lambda_n(t)$, and the positivity of V imply that

$$B(\theta)V \leq \gamma_n(\theta)^{-1}V = c_n \mathcal{D}\gamma_n(\theta)K - U \leq G(\theta)K - U;$$

hence $\Lambda_n(t)$ is a χ -subsolution of (1.1) for $t > t_n$. By Corollary 3.1 and (6.9), $\Lambda_n(t) \subset \Omega(t)$ for all $t > t_n$, and using (6.7) we conclude that

$$(6.7) \quad \alpha_n(t)(1 - n^{-1})W(1/B) \subset \Omega(t), \quad t > t_n.$$

4° For $t > t_1$ define

$$\alpha(t) = \sup \{ (1 - n^{-1})\alpha_n(t) : n \geq 1, t_n \geq t \}.$$

Then $\alpha(t)W(1/B) \subset \Omega(t)$ for $t > t_1$. Also, for each n ,

$$\alpha(t) \geq (1 - n^{-1})\alpha_n(t), \quad t \geq t_n.$$

Hence

$$\liminf_{t \rightarrow \infty} \alpha(t)/t \geq (1 - n^{-1}) \liminf_{t \rightarrow \infty} \alpha_n(t)/t = -(1 - n^{-1}) U$$

for every n , and consequently

$$(6.11) \quad \liminf_{t \rightarrow \infty} \alpha(t)/t \geq -U.$$

5° Summarizing, in 4° and Lemma 6.1 we have shown that

$$\alpha(t)W(1/B) \subset \Omega(t) \subset (-Ut + \alpha_0)W(1/B),$$

which, with (6.11), yields

$$\lim_{t \rightarrow \infty} \alpha(t)(-Ut + \alpha_0)^{-1} = 1;$$

hence $\Omega(t) \sim t|U|W(1/B)$ as $t \rightarrow \infty$. \square

7. PROOFS OF THE COMPARISON THEOREMS.

a. SUB AND SUPERDIFFERENTIALS.

We recall several definitions from the theory of viscosity solutions.²¹ Let φ be a bounded function on $(0, \infty) \times \mathbb{R}^2$, and let \mathcal{A} denote the set of symmetric 2×2 matrices. Then the subdifferential $D^+\varphi(t, \mathbf{x})$ and the superdifferential $D^-\varphi(t, \mathbf{x})$ of φ at $(t, \mathbf{x}) \in (0, \infty) \times \mathbb{R}^2$ are defined by

$$D^+\varphi(t, \mathbf{x}) = \{ (q, \mathbf{p}, \mathbf{A}) \in \mathbb{R} \times \mathbb{R}^2 \times \mathcal{A} : \limsup_{(h, \mathbf{z}) \rightarrow 0} \mathcal{D}(\varphi^*)(h, \mathbf{z}) \leq 0 \},$$

$$D^-\varphi(t, \mathbf{x}) = \{ (q, \mathbf{p}, \mathbf{A}) \in \mathbb{R} \times \mathbb{R}^2 \times \mathcal{A} : \liminf_{(h, \mathbf{z}) \rightarrow 0} \mathcal{D}(\varphi_*)(h, \mathbf{z}) \geq 0 \},$$

where

$$\begin{aligned} \mathcal{D}\Phi(h, \mathbf{z}) &= \mathcal{D}\Phi(t, \mathbf{x}; h, \mathbf{z}; q, \mathbf{p}, \mathbf{A}) = \\ &(|h| + |\mathbf{z}|^2)^{-1} \{ \Phi(t+h, \mathbf{x}+\mathbf{z}) - \Phi(t, \mathbf{x}) - hq - \mathbf{z} \cdot \mathbf{p} - \frac{1}{2} \mathbf{z} \cdot \mathbf{A} \mathbf{z} \}; \end{aligned}$$

we close the sets $D^\pm\varphi(t, \mathbf{x})$ as follows

$$cD^\pm\varphi(t, \mathbf{x}) = \{ \lim_{n \rightarrow \infty} (q_n, \mathbf{p}_n, \mathbf{A}_n) : (q_n, \mathbf{p}_n, \mathbf{A}_n) \in D^\pm\varphi(t_n, \mathbf{x}_n), (t_n, \mathbf{x}_n) \rightarrow (t, \mathbf{x}) \},$$

Then²² $(q, \mathbf{p}, \mathbf{A}) \in D^+\varphi(t, \mathbf{x})$ if and only if there is a $w \in C^{1,2}$ satisfying

$$w_t(t, \mathbf{x}) = q, \quad Dw(t, \mathbf{x}) = \mathbf{p}, \quad D^2w(t, \mathbf{x}) = \mathbf{A},$$

and (t, \mathbf{x}) is a maximum of the difference $(\varphi^* - w)$. Hence φ is a viscosity subsolution of (3.4) if and only if

$$q \leq \mathcal{F}^*(\mathbf{p}, \mathbf{A}) \quad \text{for all } (q, \mathbf{p}, \mathbf{A}) \in D^+\varphi(t, \mathbf{x})$$

²¹Cf., [CIL, §2], [C], [CEL], [FS, §5.4].

²²Cf., e.g., [FS, Prop. 4.1, §5.4]; since φ is not necessarily continuous, the proof given in [FS] must be slightly modified.

and all $(t, \mathbf{x}) \in (0, \infty) \times \mathbb{R}^2$. A limit argument then shows that

$$(7.1) \quad q \leq \mathcal{F}^*(p, A) \quad \text{for all } (q, p, A) \in cD^+\varphi(t, \mathbf{x}).$$

Similarly, φ is a viscosity supersolution of (3.4) if and only if

$$(7.2) \quad q \geq \mathcal{F}_*(p, A) \quad \text{for all } (q, p, A) \in cD^-\varphi(t, \mathbf{x})$$

and all $(t, \mathbf{x}) \in (0, \infty) \times \mathbb{R}^2$.

b. SEMICONVEX AND SEMICONCAVE FUNCTIONS.

Let $C \subset \mathbb{R}^d$ be a convex set. We say that Ψ is **semiconvex** on C if there is a constant κ such that

$$\bar{\Psi}(Y) = \Psi(Y) + \kappa|Y|^2$$

is convex on C ; $\tilde{\Psi}$ is **semiconcave** on C if $-\tilde{\Psi}$ is semiconvex.

Let Ψ be semiconvex. Since $\bar{\Psi}(Y)$ is convex, the set of subdifferentials, standard in convex analysis [C], is given by

$$\partial\bar{\Psi}(Y) = \{ P \in \mathbb{R}^d : \bar{\Psi}(\bar{Y}) \geq \bar{\Psi}(Y) + P \cdot (\bar{Y} - Y), \quad \forall \bar{Y} \in C \}.$$

In addition, wthe directional derivatives of $\bar{\Psi}$ exist and are given by

$$(7.3) \quad (\partial/\partial Z)\bar{\Psi}(Y) = \lim_{r \downarrow 0} r^{-1}[\bar{\Psi}(Y + rZ) - \bar{\Psi}(Y)] = \sup \{ P \cdot Z : P \in \partial\bar{\Psi}(Y) \}$$

for $Z \in \mathbb{R}^d \setminus \{0\}$. We now define

$$\hat{\partial}\Psi(Y) = \partial\bar{\Psi}(Y) + \{-2\kappa Y\} = \{ P : P = \bar{P} - 2\kappa Z, \quad \bar{P} \in \partial\bar{\Psi}(Y) \}.$$

Then, using the formula for the directional derivative of $\bar{\Psi}$,

$$(\partial/\partial Z)\Psi(Y) = \sup \{ P \cdot Z : P \in \hat{\partial}\Psi(Y) \}$$

for $Z \in \mathbb{R}^d \setminus \{0\}$.

The following properties of semiconvex functions are well known:

(7.4) Ψ is differentiable at Y if and only if $\hat{\partial}\Psi(Y)$ is a singleton.

(7.5) if there are sequences $P_n \rightarrow P$, $Y_n \rightarrow Y$ and convex functions $\Psi_n \rightarrow \Psi$ (uniformly on C) satisfying $P_n \in \hat{\partial}\Psi(Y_n)$ for all n , then $P \in \hat{\partial}\Psi(Y)$.

Our next result is an implicit function theorem for semiconvex functions. Let ψ be a semiconvex function on C , and let 0 be an interior point of C . We assume that ψ is differentiable²³ at 0 with a nonzero gradient P_0 ; and, without loss in generality, we assume that $\bar{P}_0 = P_0/|P_0|$ satisfies

$$\bar{P}_0 = (0, 0, \dots, 0, 1).$$

Let δ_1 be a constant satisfying

$$(7.6a) \quad |P - P_0| \leq |P_0|/2, \quad \forall P \in \hat{\partial}\bar{\Psi}(Y), \quad |Y| \leq 2\delta_1,$$

$$(7.6b) \quad P_d \geq |P_0|/2, \quad \forall P \in \hat{\partial}\bar{\Psi}(Y), \quad |Y| \leq 2\delta_1,$$

$$(7.6c) \quad \mathcal{B}(2\delta_1) = \{ Y : |Y| \leq 2\delta_1 \} \subset C.$$

Note that the existence of δ_1 follows from (7.5) (with $\Psi_n = \Psi$ for all n).

For $Y = (Y_1, Y_2, \dots, Y_{d-1}, Y_d) \in \mathbb{R}^d$ we write

$$\mathcal{P}Y = (Y_1, Y_2, \dots, Y_{d-1}) \in \mathbb{R}^{d-1},$$

Theorem 7.1 (Implicit Function Theorem). *There is a $\delta > 0$ and a unique real-valued, Lipschitz continuous function l on $\mathcal{B}(\delta)$ such that, for all Y ,*

²³We make this assumption to simplify the analysis; an analogous result holds under the weaker assumption $\hat{\partial}\bar{\Psi}(0) \cap \{0\} = \emptyset$.

$$(7.7) \quad \Psi(\mathcal{P}Y, I(Y)) = \Psi(Y_d \bar{P}_0) \quad \text{for all } Y = (\mathcal{P}Y, Y_d) \in \mathcal{B}(\delta).$$

Moreover, δ depends only on $|P_0|$, δ_1 , and the Lipschitz constant of Ψ on $\mathcal{B}(\delta_1)$.

Proof.

1° For $|Y| \leq \delta_1$ and $\alpha \in \mathbb{R}$ with $|\alpha| \leq \delta_1$, we define

$$\Phi(\alpha; Y) = \Psi(\mathcal{P}Y, \alpha) - \Psi(Y_d \bar{P}_0).$$

By (7.3),

$$\Psi(\mathcal{P}Y, \alpha) = \Psi(Y) + \int_{Y_d}^{\alpha} (\partial/\partial \bar{P}_0) \Psi(\mathcal{P}Y, \rho) d\rho,$$

$$(\partial/\partial \bar{P}_0) \Psi(\mathcal{P}Y, \rho) = \sup\{P_d : P \in \hat{\partial} \Psi(\mathcal{P}Y, \rho)\}.$$

Thus, by (7.6b), for $|Y| \leq \delta_1$ and $|\alpha| \leq \delta_1$,

$$(7.8a) \quad \Psi(\mathcal{P}Y, \alpha) \geq \Psi(Y) + (\alpha - Y_d) |P_0|/2, \quad \forall \alpha \geq Y_d,$$

$$(7.8b) \quad \Psi(\mathcal{P}Y, \alpha) \leq \Psi(Y) + (\alpha - Y_d) |P_0|/2, \quad \forall \alpha \leq Y_d.$$

2° Our next step will be to show that there is a $\delta \in (0, \delta_1]$ such that

$$(7.9) \quad \Phi(-\delta_1; Y) \leq 0 \leq \Phi(\delta_1; Y)$$

for all $|Y| \leq \delta$. Indeed, by (7.8a),

$$\begin{aligned} \Phi(\delta_1; Y) &= \Psi(\mathcal{P}Y, \delta_1) - \Psi(Y) + \Psi(Y) - \Psi(Y_d \bar{P}_0) \\ &\geq (\delta_1 - Y_d) |P_0|/2 - L |Y - Y_d \bar{P}_0|, \end{aligned}$$

where L is the Lipschitz constant of Ψ on $\mathcal{B}(\delta_1)$. (Ψ is semiconvex on

C and hence Lipschitz continuous on every compact subset interior to C.)
 Let δ be the lesser of $\delta_1/2$ and $\delta_1|P_0|/4L$. Then for all $|Y| \leq \delta$,

$$\begin{aligned}\Phi(\delta_1; Y) &\geq (\delta_1 - \delta)|P_0|/2 - L(|Y| - |Y_d|), \\ &\geq \delta_1|P_0|/4 - 2L\delta \geq 0.\end{aligned}$$

The other inequality in (7.9) is proved similarly, with the same choice for δ .

3° Fix $|Y| \leq \delta$ and consider the map

$$\ell(\alpha) = \Phi(\alpha; Y), \quad \alpha \in [-\delta_1, \delta_1].$$

Then ℓ is continuous on $[-\delta_1, \delta_1]$ with

$$\ell(-\delta_1) \leq 0 \leq \ell(\delta_1).$$

Also, the argument leading to (7.8a) yields, for $\alpha, \beta \in [-\delta_1, \delta_1]$,

$$\ell(\alpha) \geq \ell(\beta) + (\alpha - \beta)|P_0|/2 \quad \text{for } \alpha \geq \beta.$$

Hence there is a unique $\alpha_* \in [-\delta_1, \delta_1]$ such that $\ell(\alpha_*) = 0$.

4° For each $|Y| \leq \delta$ set

$$I(Y) = \alpha_*;$$

Then, for all $|Y| \leq \delta$,

$$0 = \ell(\alpha_*) = \Psi(\mathcal{P}Y, I(Y)) - \Psi(Y_d \bar{P}_0),$$

and (7.7) is satisfied.

5° Our last step will be to show that I is Lipschitz continuous. Let $|Y|, |X| \leq \delta$ be given. Then (7.8a) and the Lipschitz continuity of ψ yield

$$\begin{aligned}\Phi(I(X) + \rho; Y) &= \Psi(\mathcal{P}Y, I(X) + \rho) - \Psi(Y_d \bar{P}_0) \\ &\geq \Psi(\mathcal{P}X, I(X) + \rho) - \Psi(X_d \bar{P}_0) - 2L|X - Y| \\ &= \Psi(\mathcal{P}X, I(X) + \rho) - \Psi(\mathcal{P}X, I(X)) + \Psi(\mathcal{P}X, I(X)) - \Psi(X_d \bar{P}_0) - 2L|X - Y|\end{aligned}$$

$$\geq \rho |P_0|/2 + \Phi(I(X);X) - 2L|X - Y| \geq 0$$

provided

$$(7.10) \quad \rho \geq 4L|X - Y|/|P_0|.$$

A similar argument shows that

$$\Phi(I(X) - \rho; Y) \leq 0$$

if ρ satisfies (7.10). Hence $I(Y) \in [I(X) - \rho, I(X) + \rho]$. \square

Remark 7.1. Note that the Lipschitz constant of I is $\leq 4L/|P_0|$, with L the Lipschitz constant of Ψ on $\mathcal{B}(\delta_1)$.

The next result, the key technical contribution of the paper, will be used in an essential manner in the proof of Theorem 3.3. For $p \neq 0$, let

$$\theta(p) = \sin^{-1}(-\bar{p}_2), \quad \bar{p} = p/|p|.$$

Proposition 7.1. Let v be a semiconvex function on $[0, \infty) \times \mathbb{R}^2$ (so that $v(t, x) + \kappa(t^2 + |x|^2)$ is convex for some constant κ). Suppose that v is differentiable at (t_0, x_0) with

$$p_0 = \nabla v(t_0, x_0) \neq 0.$$

Then there exist $(t_n, x_n) \rightarrow (t_0, x_0)$ and $(q_n, p_n, A_n) \in cD^*v(t_n, x_n)$ such that

$$(7.11a) \quad \lim_{n \rightarrow \infty} (q_n, p_n) = (v_t(t_0, x_0), p_0), \quad |p_n| \neq 0,$$

$$(7.11b) \quad \liminf_{n \rightarrow \infty} \min\{\theta(p_n) - \theta(p_0), T(p_n, A_n)\} \leq 0,$$

where

$$T(p,A) = \text{trace}[(I - \bar{p} \otimes \bar{p})A].$$

To motivate the proof of this proposition, assume, for the moment, that v is smooth. For $r \in \mathbb{R}$, let

$$h(r) = \theta(\nabla v(t_0, \mathbf{x}_0 + r\mathbf{w})), \quad \mathbf{w} = ((p_0)_2, -(p_0)_1).$$

Then

$$h'(0) = T(p_0, \nabla^2 v(t_0, \mathbf{x}_0)).$$

Hence if $T(p_0, \nabla^2 v(t_0, \mathbf{x}_0)) > 0$ then $h(r) \leq h(0)$ for $r > 0$.

This argument works only for smooth v . However, if v is semiconvex, then its second derivative is bounded from below. We will use this lower bound to prove (7.11b), with \hat{H} playing the role of h (cf. 7°).

The argument given above indicates the possible validity of the following assertion, which is dual to (7.11b):

$$(7.11c) \quad \liminf_{n \rightarrow \infty} \min\{ \theta(p_0) - \theta(p_n), T^*(p_n, A_n) \} \leq 0$$

Indeed, the proof of Proposition 7.1 with minor changes establishes the existence of a sequence satisfying (7.11a) and (7.11c). One might believe further that

$$(7.11d) \quad \liminf_{n \rightarrow \infty} \min\{ \theta(p_n) - \theta(p_0), -T(p_n, A_n) \} \leq 0,$$

but (7.11d) is not valid, the reason being that, since v is assumed semiconvex, its second derivatives are necessarily bounded only from below, but the proof of (7.11d) requires an upper bound on the second derivatives; in fact, (7.11d) holds for semiconcave functions.

The proof of Proposition 7.1 will utilize the following result, which

connects the subdifferentials of convex analysis to cD^- and cD^+ . We omit the proof; similar results may be found in [FS, §2.8 and Chapter 5].

Lemma 7.1. *Let v be a semiconvex function on $[0, \infty) \times \mathbb{R}^2$. Then*

$$(q, p, A) \in cD^+v(t, x) \cup cD^-v(t, x)$$

only if

$$(7.12) \quad (q, p) \in \hat{\partial}v(t, x).$$

Conversely, (7.12) implies that

$$(q, p, -2\kappa I) \in D^-v(t, x),$$

where κ is the constant appearing in the definition of semiconvexity.

Proof of Proposition 7.1.

1° Let (y, z) denote a generic point of \mathbb{R}^2 . Assume, without loss in generality, that v is defined on $\mathbb{R} \times \mathbb{R}^2$, that $(t_0, x_0) = (0, 0, 0)$, and that

$$p_0 = |p_0|(0, 1).$$

If

$$\liminf_{\epsilon \downarrow 0} \inf \{ T(p, A) : (q, p, A) \in cD^+v(t, y, z), |t| + |y| + |z| \leq \epsilon \} \leq 0,$$

then (7.11b) follows directly; we therefore assume that there are $\gamma, \epsilon_1 > 0$ such that

$$(7.13) \quad T(p, A) \geq \gamma, \quad \forall (q, p, A) \in cD^+v(t, y, z), (t, y, z) \in \mathcal{B}(\epsilon_1).$$

The semicontinuity of v yields the existence of a $\delta_1 > 0$ satisfying (7.6), and hence of a $\delta_1 > 0$ such that, for all $(q, p) \in \hat{\partial}v(t, y, z)$,

$(t,y,z) \in \mathcal{B}(2\delta_1)$:

$$(7.14a) \quad |p - p_0| \leq |p_0|/4,$$

$$(7.14b) \quad p_2 \geq |p_0|/4.$$

2° Since v is semiconvex, v is locally Lipschitz continuous. Therefore, by Rademacher's Theorem, v is differentiable almost everywhere; we define $H(t,y,z)$ at the points of differentiability by

$$H(t,y,z) = \theta(\nabla v(t,y,z)).$$

3° By Theorem 7.1, there are a $\delta \in (0, \delta_1]$ and a function $I(t,y,z)$ satisfying

$$(7.15) \quad v(t,y,I(t,y,z)) = v(0,0,z), \quad \forall (t,y,z) \in \mathcal{B}(\delta).$$

Further, by Remark 7.1, $I(t,y,z)$ is Lipschitz continuous on $\mathcal{B}(\delta)$ with Lipschitz constant no more than $4L/|p_0|$, where L is the Lipschitz constant of v on $\mathcal{B}(\delta_1)$

4° Our next step is to show that the map $(t,y,z) \mapsto (t,y,I(t,y,z))$ with domain $\mathcal{B}(\delta)$ is one-to-one. Suppose $(t,y,I(t,y,z)) = (\bar{t}, \bar{y}, I(\bar{t}, \bar{y}, \bar{z}))$. Then $(t,y) = (\bar{t}, \bar{y})$ and

$$v(0,0,z) = v(t,y,I(t,y,z)) = v(\bar{t}, \bar{y}, I(\bar{t}, \bar{y}, \bar{z})) = v(0,0,\bar{z}).$$

Since $p_0 = |p_0|(0,1)$, (7.14b) yields

$$(7.16) \quad |v(0,0,\alpha) - v(0,0,\bar{\alpha})| \geq |\alpha - \bar{\alpha}||p_0|/2$$

for all $(0,0,\alpha), (0,0,\bar{\alpha}) \in \mathcal{B}(2\delta_1)$; hence $z = \bar{z}$.

5° The inverse of the map defined in 4° has the form $(t,y,J(t,y,z))$ with

$$v(t,y,z) = v(0,0,J(t,y,z)),$$

and we may use (7.16) to show that J is Lipschitz continuous.

6° Thus the map $(t,y,z) \mapsto (t,y,I(t,y,z))$ with domain $\mathcal{B}(\delta)$ is one-to-one and Lipschitz, with Lipschitz inverse; hence it transforms null sets into null sets.

7° In view of 6°,

$$\hat{H}(t,y,z) = H(t,y,I(t,y,z))$$

is defined for almost every $(t,y,z) \in \mathcal{B}(\delta)$. We now define, for $0 < \epsilon, \zeta \leq \delta/2$,

$$k(\epsilon, \zeta) = \int_{\mathcal{B}(\epsilon)} [\hat{H}(t,y+\zeta,z) - \hat{H}(t,y,z)] dt dy dz.$$

In 9°-15° we shall show that, for sufficiently small $\epsilon, \zeta > 0$,

$$(7.17) \quad k(\epsilon, \zeta) \geq |\mathcal{B}_1| \delta \zeta \epsilon^3 / 2 |p_0|$$

with δ as in (7.13), where $|\mathcal{B}_1|$ is the volume of the unit ball in \mathbb{R}^3 . The above estimate provides a weak method of proving that $\hat{H}(t,y,z)$ is increasing in y . Indeed, if v were smooth, a direct calculation would yield

$$\hat{H}_y(0,0,z) = |p_0|^{-1} T(p_0, \nabla^2 v(0,0,z)) \geq \delta / |p_0|.$$

(The details are given in 9°.)

We shall assume that (7.17) is valid and complete the proof of (7.11a,b), before proving (7.17).

8° Let \mathcal{O} denote the set of points of differentiability of v . For all sufficiently small $\epsilon, \zeta > 0$ there are $(\bar{t}, \bar{y}, \bar{z}) \in \mathcal{B}(\epsilon)$ satisfying

$$(\bar{t}, \bar{y}, I(\bar{t}, \bar{y}, \bar{z})) \in \mathcal{O}, \quad (\bar{t}, \bar{y} + \zeta, I(\bar{t}, \bar{y} + \zeta, \bar{z})) \in \mathcal{O},$$

$$(7.18) \quad \hat{H}(\bar{t}, \bar{y} + \zeta, \bar{z}) - \hat{H}(\bar{t}, \bar{y}, \bar{z}) \geq \delta \zeta / 2 |p_0|.$$

Moreover, by (7.5) and Lemma 7.1,

$$\lim_{\rho \downarrow 0} (\sup \{ |H(t,y,z) - \theta(\mathbf{p}_0)| : (t,y,z) \in \mathcal{B}(\rho) \cap \mathcal{O} \}) = 0.$$

Since $I(0,0,0) = 0$, by choosing $\varepsilon > 0$ small we can make $|H(\bar{t}, \bar{y}, \bar{z}) - \theta(\mathbf{p}_0)|$ smaller than $\varepsilon \zeta / 2 |\mathbf{p}_0|$. Therefore for every $\zeta = 1/n$ there are $\varepsilon_n \downarrow 0$ and $(\bar{t}_n, \bar{y}_n, \bar{z}_n) \in \mathcal{B}(\varepsilon_n)$ satisfying

$$\begin{aligned} (t_n, y_n, z_n) &:= (\bar{t}_n, \bar{y}_n + n^{-1}, I(\bar{t}_n, \bar{y}_n + n^{-1}, \bar{z}_n)) \in \mathcal{O}, \\ H(t_n, y_n, z_n) &> \theta(\mathbf{p}_0). \end{aligned}$$

This completes the proof of (7.11a,b), granted (7.17).

We now turn to a proof of (7.17)

9° We now assume that v is smooth, a restriction we will later remove using mollification. Since v is smooth, H is defined everywhere. Recall that

$$\begin{aligned} H(t,y,z) &= \theta(\nabla v(t,y,z)), & \hat{H}(t,y,z) &= H(t,y,I(t,y,z)), \\ \theta(\mathbf{p}) &= \sin^{-1}(-\bar{p}_2), & \bar{\mathbf{p}} &= \mathbf{p}/|\mathbf{p}|, \\ v(t,y,I(t,y,z)) &= v(0,0,z). \end{aligned}$$

We claim that

$$(7.19) \quad \hat{H}_y(t,y,z) = T(\nabla v(\boldsymbol{\xi}), \nabla^2 v(\boldsymbol{\xi})) \mathbf{v}_z(\boldsymbol{\xi})^{-1}, \quad \boldsymbol{\xi} = (t,y,I(t,y,z)).$$

This formula may be verified using a direct but tedious calculation; instead we give an indirect derivation, which also motivates our reason for computing \hat{H}_y .

For (t,z) fixed, the parametrized curve $\Gamma : y \mapsto (y, I(t,y,z))$, $|y| \leq \delta$ is a subset of the $v(0,0,z)$ level curve of v . Hence the normal angle of Γ is $\theta = \hat{H}(t,y,z)$ and the curvature is given by

$$\theta_s = T(\nabla v, \nabla^2 v) / |\nabla v|$$

with s the arc length. Thus

$$\hat{H}_y = T(\nabla v, \nabla^2 v) s_y / |\nabla v|, \quad s_y = (1 + I_y^2)^{1/2}.$$

By differentiating (7.15) with respect to y we obtain

$$v_y + v_z I_y = 0;$$

hence $s_y = |\nabla v|/v_z$, which, when substituted into the previous formula, yields (7.19).

10° We continue to assume that v is smooth. The definition of $k(\varepsilon, \xi)$ yields

$$k(\varepsilon, \xi) = \left[\int_{\mathcal{B}(\varepsilon)} \int_0^1 \hat{H}_y(t, y+r\xi, z) dr dt dy dz \right] \xi.$$

Let

$$K(t, y, z) = T(\nabla v(\xi), \nabla^2 v(\xi)) v_z(\xi)^{-1}.$$

Then, by (7.19),

$$k(\varepsilon, \xi) = \left[\int_{\mathcal{B}(\varepsilon)} \int_0^1 K(t, y+r\xi, z) dr dt dy dz \right] \xi.$$

for all $0 < \varepsilon, \xi \leq \delta/2$, and (7.17) follows from (7.13) and (7.14). Thus we have established (7.17) for v smooth. We now remove this restriction; here the manner in which δ depends on v is important. The constant δ comes from Theorem 7.1 and hence depends only on $|\mathcal{D}v(0,0,0)|$, δ_1 , and the Lipschitz constant of v on $\mathcal{B}(\delta_1)$; the constant δ_1 is chosen in 1° and satisfies both (7.6) and (7.14).

11° Let v_n be a mollification of v . Then v_n converges to v uniformly on compact sets; $v_n(t, y, z) + \kappa(t^2 + y^2 + z^2)$ is convex, with κ as in the statement of the proposition; on compact sets, the Lipschitz constant

of v_n is \leq the Lipschitz constant of v .

Let k_n and K_n be defined as in 10°, but with v replaced by v_n . Then 10° yields

$$k_n(\varepsilon, \zeta) = \left[\int_{\mathcal{B}(\varepsilon)} \int_0^1 K_n(t, y+r\zeta, z) dr dt dy dz \right] \zeta.$$

for all $0 < \varepsilon, \zeta \leq \delta_n/2$.

12° Consider a sequence $(t_n, y_n, z_n) \rightarrow (t, y, z)$. Since $Dv_n(t_n, y_n, z_n)$ (the derivative in \mathbb{R}^3) is uniformly bounded in n , it has a subsequence, also denoted by n , such that $Dv_n(t_n, y_n, z_n)$ is convergent with limit (q, p) . Then, by (7.5), $(q, p) \in \hat{\partial}v(t, y, z)$. Thus $Dv_n(t_n, y_n, z_n) \rightarrow Dv(t, y, z)$ for any sequence $(t_n, y_n, z_n) \rightarrow (t, y, z) \in \mathcal{O}$. In particular,

$$(7.22a) \quad \lim_{n \rightarrow \infty} Dv_n(0, 0, 0) = Dv(0, 0, 0)$$

$$(7.22b) \quad \lim_{n \rightarrow \infty} k_n(\varepsilon, \zeta) = k(\varepsilon, \zeta)$$

for all $0 < \varepsilon, \zeta \leq \delta_n/2$ for sufficiently large n .

13° Recall that $\delta_1 > 0$ satisfies (7.6) and (7.14). In view of the previous steps, we may choose $\delta_{1n} \rightarrow \delta_1$, as $n \rightarrow \infty$, satisfying (7.6) and (7.14) with v replaced by v_n . Since the Lipschitz constant of v_n is \leq that of v on each compact set, 12° and the discussion just before 11° imply that $\delta_n \rightarrow \delta$.

14° Recall the definition of k_n given in 11°. We claim that there is a subsequence, also labeled by n , such that

$$(7.23) \quad \liminf_{n \rightarrow \infty} k_n(\varepsilon, \zeta) \geq |\mathcal{B}_1| \delta \varepsilon^3 / 2 |p_0|$$

for all sufficiently small $0 < \varepsilon, \zeta$. Indeed, since v is semiconvex,

$$D^2v = M + \Lambda,$$

where M is an integrable matrix-valued function and Λ is a matrix-valued measure orthogonal to the Lebesgue measure (cf. [J, Proposition 3.3]). Moreover, $\Lambda \geq 0$ and

$$(7.24) \quad \nu \cdot M(t, y, z) \nu \geq -\kappa |\nu|^2, \quad \forall \nu \in \mathbb{R}^3$$

with κ as in the statement of the proposition. Since $v_n = v * m_n$ for some smooth mollifier m_n ,

$$D^2 v_n = M_n + \Lambda_n, \quad M_n = M * m_n, \quad \Lambda_n = \Lambda * m_n.$$

The measure Λ_n has density with respect to Lebesgue measure. Moreover, $\Lambda_n \geq 0$ and M_n satisfies (7.24).

The monotonicity of $T(p, A)$ in A and the positivity of Λ imply that

$$K_n(t, y, z) = T(\nabla v_n(\xi_n), \nabla^2 v_n(\xi_n)) / (\partial/\partial z) v_n(\xi_n), \\ \xi_n = (t, y, I_n(t, y, z)).$$

Suppose $\xi = (t, y, I(t, y, z)) \in \Theta$. Then, by 12°,

$$\nabla v_n(\xi_n) \rightarrow \nabla v(\xi), \quad (\partial/\partial z) v_n(\xi_n) \rightarrow (\partial/\partial z) v(\xi),$$

Further,

$$K_n(t, y, z) \geq T(\nabla v_n(\xi_n), M_n(\xi_n)) / 2|p_0|$$

for all $(t, y, z) \in \mathcal{B}(\delta_n)$ and sufficiently large n . Also $M_n \rightarrow M$ in L^1 . Recall that the map $(t, y, z) \mapsto (t, y, I(t, y, z))$, on $\mathcal{B}(\delta)$, is one-to-one and Lipschitz, with Lipschitz inverse (cf. 6°). Let

$$\hat{M}_n(t, y, z) = M_n(\xi_n), \quad \hat{M}(t, y, z) = M(\xi).$$

Then $\hat{M}_n \rightarrow \hat{M}$ in $L^1(\mathcal{B}(\delta))$. Therefore, by passing to a subsequence, also

labeled by n ,

$$\lim_{n \rightarrow \infty} \hat{M}_n(t, y, z) = \hat{M}(t, y, z)$$

for almost every $(t, y, z) \in \mathcal{B}(\delta)$. Since

$$\nu \cdot M_n(\xi_n) \nu \geq -\kappa |\nu|^2,$$

it follows that

$$T(\nabla v_n(\xi_n), M_n(\xi_n)) \geq -\kappa.$$

Fatou's Lemma then yields

$$\liminf_{n \rightarrow \infty} \int_{\mathcal{B}(\varepsilon)} T(\nabla v_n(\xi_n), M_n(\xi_n)) dt dy dz \geq$$

$$\int_{\mathcal{B}(\varepsilon)} T(\nabla v(\xi), M(\xi)) dt dy dz$$

for all $\varepsilon < \delta/2$. Hence, for $0 < \varepsilon, \xi < \delta/2$,

$$\liminf_{n \rightarrow \infty} k_n(\varepsilon, \xi) \geq \left[\int_{\mathcal{B}(\varepsilon)} \int_0^1 T(t, y+r\xi, z) dr dt dy dz \right] \xi / 2|p_0|,$$

where

$$T(t, y, z) = T(\nabla v(\xi), M(\xi)).$$

Also for $\xi \in \mathcal{O}$, $\xi \notin \text{supp } \Lambda$,

$$(Dv(\xi), M(\xi)) \in D^+v(\xi).$$

Let $\hat{\mathcal{O}} = \mathcal{O} \cap \text{complement}(\text{supp } \Lambda)$. Then $\hat{\mathcal{O}}$ has full measure and, by 6°, so also has $\{(t, y, z) \in \mathcal{B}(\delta) : \xi(t, y, z) \in \hat{\mathcal{O}}\}$. Moreover, by (7.13), $T(t, y, z) \geq \delta$ for

every $(t, y, z) \in \mathcal{B}(\varepsilon_1) \cap \hat{\mathcal{O}}$. We have therefore proved (7.23).

15° The desired result (7.17) follows from (7.21), (7.22), and (7.23). \square

c. SEMICONVEX AND SEMICONCAVE APPROXIMATIONS.

For $\varepsilon > 0$ and $(t, \mathbf{x}) \in [0, \infty) \times \mathbb{R}^2$, we define

$$\begin{aligned}\varphi^\varepsilon(t, \mathbf{x}) &= \sup \{ \varphi^*(s, \mathbf{y}) - (4\varepsilon)^{-2}(|t-s|^4 + |\mathbf{x}-\mathbf{y}|^4) : (s, \mathbf{y}) \in [0, \infty) \times \mathbb{R}^2 \}, \\ \varphi_\varepsilon(t, \mathbf{x}) &= \inf \{ \varphi_*(s, \mathbf{y}) + (4\varepsilon)^{-2}(|t-s|^4 + |\mathbf{x}-\mathbf{y}|^4) : (s, \mathbf{y}) \in [0, \infty) \times \mathbb{R}^2 \}.\end{aligned}$$

These definitions are similar to the sup and inf convolutions of the theory of viscosity solutions [LL,FS,JLS,CIL], in which the second power rather than the fourth is used in the translations. Our reasons for using the fourth power are its simplification of our proof of comparison (cf. Lemma 7.4c).

Let φ be bounded. Then φ^ε is semiconvex. To verify this, choose a maximizer (s_0, \mathbf{y}_0) in the definition of $\varphi^\varepsilon(t_0, \mathbf{x}_0)$, and set

$$r = t_0 - s_0, \quad \mathbf{w} = \mathbf{x}_0 - \mathbf{y}_0.$$

Then

$$\begin{aligned}\varphi^\varepsilon(t_0, \mathbf{x}_0) &= \varphi^*(s_0, \mathbf{y}_0) - (4\varepsilon)^{-2}(r^4 + |\mathbf{w}|^4), \\ \varphi^\varepsilon(t, \mathbf{x}) &\geq \varphi^*(s_0, \mathbf{y}_0) - (4\varepsilon)^{-2}(|t-s_0|^4 + |\mathbf{x}-\mathbf{y}_0|^4)\end{aligned}$$

for all (t, \mathbf{x}) . For $0 < h \leq t_0$ and $\mathbf{z} \in \mathbb{R}^2$, we use this inequality at $(t, \mathbf{x}) = (t_0 \pm h, \mathbf{x}_0 \pm \mathbf{z})$ to obtain

$$\begin{aligned}Q(t_0, \mathbf{x}_0; h, \mathbf{z}) &= \varphi^\varepsilon(t_0 + h, \mathbf{x}_0 + \mathbf{z}) + \varphi^\varepsilon(t_0 - h, \mathbf{x}_0 - \mathbf{z}) - 2\varphi^\varepsilon(t_0, \mathbf{x}_0) \\ &\geq - (4\varepsilon)^{-2}[(r+h)^4 + (r-h)^4 - 2r^4 + |\mathbf{w} + \mathbf{z}|^4 + |\mathbf{w} - \mathbf{z}|^4 - |\mathbf{z}|^4].\end{aligned}$$

Hence

$$\liminf_{(h, \mathbf{z}) \rightarrow 0} [h^2 + |\mathbf{z}|^2]^{-1} Q(t_0, \mathbf{x}_0; h, \mathbf{z}) \geq -3(4\varepsilon)^{-2}[r^2 + |\mathbf{w}|^2].$$

Also,

$$(4\varepsilon)^{-2}[r^4 + |w|^4] = \varphi^*(s_0, y_0) - \varphi^\varepsilon(t_0, x_0) \leq 2\|\varphi\|,$$

with $\|\cdot\|$ the sup norm. Therefore

$$D^2\varphi^\varepsilon \geq -(\kappa/\varepsilon)I \quad \text{in } \mathcal{D}'$$

for some constant κ depending only on $\|\varphi\|$; hence φ^ε is semiconvex. A similar argument shows that φ_ε is semiconcave (cf. [CIL, §3], [FS, §5.4]). Also, as $\varepsilon \downarrow 0$,

$$\varphi^\varepsilon(t, \mathbf{x}) \downarrow \varphi^*(t, \mathbf{x}), \quad \varphi_\varepsilon(t, \mathbf{x}) \uparrow \varphi_*(t, \mathbf{x})$$

for all $(t, \mathbf{x}) \in [0, \infty) \times \mathbb{R}^2$.

The next lemma is similar to [FS, Lemma 7.2, §5.7] (see also [CIL, §3]). Let $M^\varepsilon(t, \mathbf{x})$ denote the set of all maximizers in the definition of $\varphi^\varepsilon(t, \mathbf{x})$ and $m^\varepsilon(t, \mathbf{x})$ the set of all minimizers in the definition of $\varphi_\varepsilon(t, \mathbf{x})$.

Lemma 7.2. Fix $(t, \mathbf{x}) \in (0, \infty) \times \mathbb{R}^2$ and $\varepsilon > 0$. Then

$$(7.24) \quad |t-s|^4 + |\mathbf{x}-\mathbf{y}|^4 \leq 8\|\varphi\|\varepsilon^2,$$

for every $(s, \mathbf{y}) \in M^\varepsilon(t, \mathbf{x}) \cap m^\varepsilon(t, \mathbf{x})$. Suppose

$$(7.25) \quad t > t_\varepsilon := (8\|\varphi\|\varepsilon^2)^{1/4}.$$

Then

$$D^+\varphi^\varepsilon(t, \mathbf{x}) \subset D^+\varphi(s, \mathbf{y})$$

for every $(s, \mathbf{y}) \in M^\varepsilon(t, \mathbf{x})$ and

$$D^-\varphi_\varepsilon(t, \mathbf{x}) \subset D^-\varphi(s, \mathbf{y})$$

for every $(s, \mathbf{y}) \in m^\varepsilon(t, \mathbf{x})$.

Suppose that φ is a viscosity solution of (3.4) in $(0, \infty) \times \mathbb{R}^2$. Then we may use Lemma 7.2 and (7.1) to conclude that φ^ε is a viscosity subsolution of (3.4) in $(t_\varepsilon, \infty) \times \mathbb{R}^2$ and that φ_ε is a viscosity supersolution of (3.4) in $(t_\varepsilon, \infty) \times \mathbb{R}^2$.

Lemma 7.3. *Suppose that*

$$(q, p, A) \in cD^+\varphi^\varepsilon(t_0, \mathbf{x}_0).$$

Then

$$(7.26) \quad A \geq -3(|p|/\varepsilon)^{2/3}I.$$

Proof. We assume, without loss in generality, that $(q, p, A) \in D^+\varphi^\varepsilon(t_0, \mathbf{x}_0)$. Then there is a function $w \in C^{1,2}$ such that

$$w_t(t_0, \mathbf{x}_0) = q, \quad \nabla w(t_0, \mathbf{x}_0) = p, \quad \nabla^2 w(t_0, \mathbf{x}_0) = A$$

and (t_0, \mathbf{x}_0) is a maximizer of $\varphi^\varepsilon - w$. Let

$$\psi(t, \mathbf{x}; s, \mathbf{y}) = \varphi^*(s, \mathbf{y}) - (4\varepsilon)^{-2}(|t-s|^4 + |\mathbf{x}-\mathbf{y}|^4) - w(t, \mathbf{x}).$$

Choose $(s_0, \mathbf{y}_0) \in M^\varepsilon(t_0, \mathbf{x}_0)$. Then ψ has a maximum at $(t_0, \mathbf{x}_0; s_0, \mathbf{y}_0)$. Thus

$$\begin{aligned} \varepsilon^{-2}(\mathbf{y}_0 - \mathbf{x}_0)|\mathbf{y}_0 - \mathbf{x}_0|^2 &= \nabla w(t_0, \mathbf{x}_0) = p, \\ -\varepsilon^{-2}[|\mathbf{y}_0 - \mathbf{x}_0|^2 I + 2(\mathbf{y}_0 - \mathbf{x}_0) \otimes (\mathbf{y}_0 - \mathbf{x}_0)] &\leq \nabla^2 w(t_0, \mathbf{x}_0) = A, \end{aligned}$$

and (7.25) follows. \square

A similar argument yields

$$(7.27) \quad B \leq 3(|p|/\varepsilon)^{2/3}I.$$

for all $(q,p,B) \in cD^-\varphi^\varepsilon(t_0, \mathbf{x}_0)$.

Lemma 7.4. Let $\varepsilon, \beta > 0$ and bounded functions φ, ψ on $[0, \infty) \times \mathbb{R}^2$ be given. Suppose that $(t_0, \mathbf{x}_0) \in (t_\varepsilon, \infty) \times \mathbb{R}^2$ is a maximizer of t_ε (cf. (7.25))

$$W^\varepsilon(t, \mathbf{x}) = \varphi^\varepsilon(t, \mathbf{x}) - \psi_\varepsilon(t, \mathbf{x}) - \beta t.$$

Then:

(a) φ^ε and ψ_ε are differentiable at (t_0, \mathbf{x}_0) with

$$(7.28a) \quad \nabla \varphi^\varepsilon(t_0, \mathbf{x}_0) = \nabla \psi_\varepsilon(t_0, \mathbf{x}_0) =: \mathbf{p}_\varepsilon,$$

$$(7.28b) \quad -\beta + (\varphi^\varepsilon)_t(t_0, \mathbf{x}_0) = (\psi_\varepsilon)_t(t_0, \mathbf{x}_0) =: q_\varepsilon;$$

(b) there are symmetric matrices $\mathbf{A}_\varepsilon \leq \mathbf{B}_\varepsilon$ such that

$$(q_\varepsilon + \beta, \mathbf{p}_\varepsilon, \mathbf{A}_\varepsilon) \in cD^+\varphi^\varepsilon(t_0, \mathbf{x}_0), \quad (q_\varepsilon, \mathbf{p}_\varepsilon, \mathbf{B}_\varepsilon) \in cD^-\psi_\varepsilon(t_0, \mathbf{x}_0);$$

(c) if $\mathbf{p}_\varepsilon = 0$, then $\mathbf{A}_\varepsilon = \mathbf{B}_\varepsilon = 0$.

Proof. (a) Recall that φ^ε and ψ_ε are semiconvex and semiconcave, respectively. Thus there is a κ_ε such that

$$\bar{\varphi}(t, \mathbf{x}) = \varphi^\varepsilon(t, \mathbf{x}) + \kappa_\varepsilon(t^2 + |\mathbf{x}|^2)$$

is convex and

$$\bar{\psi}(t, \mathbf{x}) = \psi_\varepsilon(t, \mathbf{x}) - \kappa_\varepsilon(t^2 + |\mathbf{x}|^2)$$

is concave.

Let $(q_1, \mathbf{p}_1) \in \partial \bar{\varphi}(t_0, \mathbf{x}_0)$, $(q_2, \mathbf{p}_2) \in -\partial(-\bar{\psi})(t_0, \mathbf{x}_0)$. Since (t_0, \mathbf{x}_0) is a maximizer of $W^\varepsilon(t, \mathbf{x})$,

$$\begin{aligned}\bar{\varphi}(t, \mathbf{x}) - \bar{\psi}(t, \mathbf{x}) &= W^\varepsilon(t, \mathbf{x}) + \beta t + 2\kappa_\varepsilon(t^2 + |\mathbf{x}|^2) \\ &\leq W^\varepsilon(t_0, \mathbf{x}_0) + \beta t + 2\kappa_\varepsilon(t^2 + |\mathbf{x}|^2).\end{aligned}$$

Also, by the definition of the subdifferentials $\partial\bar{\varphi}$ and $-\partial(-\bar{\psi})$,

$$\begin{aligned}\bar{\varphi}(t, \mathbf{x}) - \bar{\psi}(t, \mathbf{x}) &\geq \bar{\varphi}(t_0, \mathbf{x}_0) - \bar{\psi}(t_0, \mathbf{x}_0) + (q_1 - q_2)(t - t_0) + (\mathbf{p}_1 - \mathbf{p}_2) \cdot (\mathbf{x} - \mathbf{x}_0) \\ &= W^\varepsilon(t_0, \mathbf{x}_0) + \beta t_0 + 2\kappa_\varepsilon(t_0^2 + |\mathbf{x}_0|^2) + (q_1 - q_2)(t - t_0) + (\mathbf{p}_1 - \mathbf{p}_2) \cdot (\mathbf{x} - \mathbf{x}_0).\end{aligned}$$

Thus

$$(q_1 - q_2 - \beta)(t - t_0) + (\mathbf{p}_1 - \mathbf{p}_2) \cdot (\mathbf{x} - \mathbf{x}_0) \leq 2\kappa_\varepsilon(t^2 + |\mathbf{x}|^2 - t_0^2 - |\mathbf{x}_0|^2)$$

for all (t, \mathbf{x}) , so that

$$q_1 = q_2 + \beta, \quad \mathbf{p}_1 = \mathbf{p}_2$$

for all $(q_1, \mathbf{p}_1) \in \partial\bar{\varphi}(t_0, \mathbf{x}_0)$, $(q_2, \mathbf{p}_2) \in -\partial(-\bar{\psi})(t_0, \mathbf{x}_0)$. Hence $\partial\bar{\varphi}(t_0, \mathbf{x}_0)$ and $-\partial(-\bar{\psi})(t_0, \mathbf{x}_0)$ are singletons, and $\bar{\psi}$ and $\bar{\varphi}$ are differentiable at (t_0, \mathbf{x}_0) . Assertion (a) then follows from the definitions of $\bar{\psi}$ and $\bar{\varphi}$.

(b) Since φ^ε is semiconvex and ψ_ε semiconcave, this assertion follows from (7.25), (7.26), and Jensen's maximum principle [Je], [CIL, §3], [FS, Theorem 5.1, §5.5].

(c) Using (7.26) and (7.27), we obtain

$$-3(|\mathbf{p}_\varepsilon|/\varepsilon)^{2/3} \mathbf{I} \leq \mathbf{A}_\varepsilon \leq \mathbf{B}_\varepsilon \leq 3(|\mathbf{p}_\varepsilon|/\varepsilon)^{2/3} \mathbf{I}. \quad \square$$

d. PROOF OF THEOREM 3.3.

We will prove Theorem 3.3 by contradiction. Suppose that conclusion (5.1) is invalid.

1° By hypothesis, $\varphi \in \hat{M}([0, T] \times \mathbb{R}^2)$ and $\psi \in \hat{M}([0, T] \times \mathbb{R}^2)$; thus there are constants $\alpha, \hat{\alpha}, R$ such that

$$\varphi(t, \mathbf{x}) = \alpha, \quad \psi(t, \mathbf{x}) = \hat{\alpha} \quad \text{for } |\mathbf{x}| \geq R, \quad t \in [0, T].$$

Thus, by (7.24), for all sufficiently small ε ,

$$\varphi^\varepsilon(t, \mathbf{x}) = \alpha, \quad \psi_\varepsilon(t, \mathbf{x}) = \hat{\alpha} \quad \text{for } |\mathbf{x}| \geq R + 1, \quad t \in [0, T].$$

2° Set

$$I = \sup_{\mathbf{x} \in \mathbb{R}^2} [\varphi^*(0, \mathbf{x}) - \psi_*(0, \mathbf{x})]$$

Then $I \geq \alpha - \hat{\alpha}$. Since (5.1) does not hold, there are $(s, \mathbf{y}) \in (0, T] \times \mathcal{B}(R)$ and $\gamma > 0$ such that

$$(7.29) \quad \varphi^*(s, \mathbf{y}) - \psi_*(s, \mathbf{y}) \geq I + \gamma.$$

3° For $\varepsilon, \beta > 0$ consider the function

$$H^\varepsilon(t, \mathbf{x}) = \varphi^\varepsilon(t, \mathbf{x}) - \psi_\varepsilon(t, \mathbf{x}) - \beta t$$

for $(t, \mathbf{x}) \in [0, T] \times \mathbb{R}^2$. Then, by the definitions of φ^ε and ψ_ε ,

$$H^\varepsilon(t, \mathbf{x}) \geq \varphi^*(t, \mathbf{x}) - \psi_*(t, \mathbf{x}) - \beta t$$

for all $(t, \mathbf{x}) \in [0, T] \times \mathbb{R}^2$. In particular,

$$H^\varepsilon(s, \mathbf{y}) \geq I + \gamma - \beta T.$$

Therefore, by 1° and the inequality $I \geq \alpha - \hat{\alpha}$, for $\beta < \gamma/T$, H^ε achieves its maximum at some $(t(\varepsilon), \mathbf{x}(\varepsilon)) \in (0, T] \times \mathcal{B}(R + 1)$.

4° Suppose $t(\varepsilon) \leq t_\varepsilon$ for all sufficiently small $\varepsilon > 0$, where t_ε is defined in (7.25). Since $|\mathbf{x}(\varepsilon)| \leq R + 1$, there is a subsequence, also labeled by ε , such that $(t(\varepsilon), \mathbf{x}(\varepsilon)) \rightarrow (0, \mathbf{z})$ as $\varepsilon \downarrow 0$, and

$$(7.30) \quad I \geq \varphi^*(0, \mathbf{z}) - \psi_*(0, \mathbf{z}).$$

Also,

$$(7.31) \quad \varphi^\varepsilon(t(\varepsilon), \mathbf{x}(\varepsilon)) - \psi_\varepsilon(t(\varepsilon), \mathbf{x}(\varepsilon)) - \beta t(\varepsilon) \geq I + \gamma - \beta T.$$

Choose $(s(\varepsilon), \mathbf{y}(\varepsilon)) \in M^\varepsilon(t(\varepsilon), \mathbf{x}(\varepsilon))$. (Recall that $M^\varepsilon(t, \mathbf{x})$ is the set of all maximizers in the definition of $\varphi^\varepsilon(t, \mathbf{x})$.) Then, by (7.24), $(s(\varepsilon), \mathbf{y}(\varepsilon)) \rightarrow (0, \mathbf{z})$ as $\varepsilon \downarrow 0$. Moreover,

$$\varphi^\varepsilon(t(\varepsilon), \mathbf{x}(\varepsilon)) \leq \varphi_*(s(\varepsilon), \mathbf{y}(\varepsilon))$$

and therefore

$$\limsup_{\varepsilon \downarrow 0} \varphi^\varepsilon(t(\varepsilon), \mathbf{x}(\varepsilon)) \leq \varphi^*(0, \mathbf{z}).$$

Similarly,

$$\liminf_{\varepsilon \downarrow 0} \psi_\varepsilon(t(\varepsilon), \mathbf{x}(\varepsilon)) \geq \psi_*(0, \mathbf{z}).$$

Using (7.30), (7.31), and the inequalities above, we are led to the inequality $\beta \geq \gamma/T$.

5° We now fix

$$\beta = \gamma/2T.$$

Then, in view of the previous step, $t(\varepsilon) \leq t_\varepsilon$ for some small $\varepsilon > 0$. Let $(t_0, \mathbf{x}_0) = (t(\varepsilon), \mathbf{x}(\varepsilon))$. Then, by Lemma 7.4, φ^ε and ψ_ε are differentiable at (t_0, \mathbf{x}_0) and (7.28) are satisfied. Moreover, there are $A_\varepsilon \leq B_\varepsilon$ satisfying Lemma 7.4b. We now write q_0, p_0, A_0, B_0 for $q_\varepsilon, p_\varepsilon, A_\varepsilon, B_\varepsilon$ to emphasize the fact that ε is now fixed. Since φ^ε and ψ_ε are, respectively, a viscosity subsolution and supersolution of (3.4), Lemma 7.4, (7.1), and (7.2) imply that

$$(7.32) \quad q_0 + \beta \leq \mathcal{F}^*(p_0, A_0),$$

$$(7.33) \quad q_0 \geq \mathcal{F}_*(p_0, B_0).$$

6° Suppose that $p_0 = 0$. Then, by Lemma 7.4c, $A_0 = B_0 = 0$ and (7.32) and (7.33) yield

$$q_0 + \beta \leq \mathcal{F}^*(0, 0) = 0 \leq q_0,$$

which contradicts the positivity of β . Hence $p_0 \neq 0$.

7° Suppose that G is continuous at

$$\theta_0 = \theta(p_0).$$

Then for any symmetric matrix A ,

$$\mathcal{F}^*(p_0, A) = \mathcal{F}_*(p_0, A) = \mathcal{F}(p_0, A).$$

Since $A_0 \leq B_0$, the ellipticity property (3.13), (7.32), and (7.33) yield

$$q_0 + \beta \leq \mathcal{F}^*(p_0, A_0) = \mathcal{F}(p_0, A_0) \leq \mathcal{F}(p_0, B_0) = \mathcal{F}_*(p_0, B_0) \leq q_0,$$

which again contradicts the positivity of β .

8° Suppose that G is discontinuous at θ_0 . Then, by (1.2), there is a $\gamma > 0$ such that $G(\theta) = 0$ either for all $\theta \in [\theta_0, \theta_0 + \gamma]$ or for all $\theta \in [\theta_0 - \gamma, \theta_0]$. We will consider only the case in which

$$G(\theta) = 0, \quad \forall \theta \in [\theta_0 - \gamma, \theta_0];$$

the other case is treated similarly. Let

$$\rho = B(\theta_0)^{-1}, \quad G_0 = \rho \lim_{\theta \downarrow \theta_0} G(\theta).$$

Then

$$\mathcal{F}^*(\mathbf{p}_0, \mathbf{A}_0) = -\rho U|\mathbf{p}_0| + G_0(T(\mathbf{p}_0, \mathbf{A}_0))^+,$$

$$\mathcal{F}_*(\mathbf{p}_0, \mathbf{B}_0) = -\rho U|\mathbf{p}_0| - G_0(T(\mathbf{p}_0, \mathbf{B}_0))^-,$$

where $(\alpha)^+ = \max(\alpha, 0)$ and $(\alpha)^- = (-\alpha)^+$.

9° We will analyze three cases separately.

Case $\mathbf{A}_0 \geq 0$. Since $\mathbf{B}_0 \geq \mathbf{A}_0$, it follows that $\mathbf{B}_0 \geq 0$ and $T(\mathbf{p}_0, \mathbf{B}_0) \geq 0$. Then, by (7.33),

$$(7.34) \quad q_0 \geq \mathcal{F}_*(\mathbf{p}_0, \mathbf{B}_0) = -\rho U|\mathbf{p}_0|.$$

We now use Proposition 7.1 to construct

$$(7.35) \quad (q_n, \mathbf{p}_n, \mathbf{A}_n) \in cD^+\varphi^\varepsilon(t_n, \mathbf{x}_n),$$

$$(q_n, \mathbf{p}_n) \rightarrow (q_0 + \beta, \mathbf{p}_0), \quad (t_n, \mathbf{x}_n) \rightarrow (t_0, \mathbf{x}_0)$$

satisfying (7.11b). By (7.35)

$$q_n \leq \mathcal{F}^*(\mathbf{p}_n, \mathbf{A}_n)$$

and therefore

$$(7.36) \quad q_0 + \beta \leq \liminf_{n \rightarrow \infty} \mathcal{F}^*(\mathbf{p}_n, \mathbf{A}_n).$$

On the other hand, (7.11b) implies that

$$(7.37) \quad \liminf_{n \rightarrow \infty} \mathcal{F}^*(\mathbf{p}_n, \mathbf{A}_n) \leq -\rho U|\mathbf{p}_0|.$$

Indeed, (7.11b) yields either $\theta(\mathbf{p}_n) < \theta_0$ or

$$(7.38) \quad \liminf_{n \rightarrow \infty} T(\mathbf{p}_n, \mathbf{A}_n) \leq 0.$$

In the first case $G(\theta(\mathbf{p}_n)) = 0$ and

$$\mathcal{F}^*(\mathbf{p}_n, \mathbf{A}_n) = -\rho U|\mathbf{p}_n|,$$

and hence (7.37) follows from the convergence of \mathbf{p}_n to \mathbf{p}_0 . On the other hand, (7.38) and 8° yield

$$\mathcal{F}^*(\mathbf{p}_n, \mathbf{A}_n) \leq -\rho U|\mathbf{p}_n| + \max_{\theta} [G(\theta)B(\theta)^{-1}](T(\mathbf{p}_n, \mathbf{A}_n))^+,$$

which implies (7.37). Now combine (7.36) and (7.37) to obtain

$$\varepsilon + \beta \leq \rho U|\mathbf{p}_0|,$$

which, with (3.4), contradicts the positivity of ρ .

Case $B_0 \leq 0$. Then $A_0 \leq 0$ and

$$\mathcal{F}^*(\mathbf{p}_0, \mathbf{A}_0) = -\rho U|\mathbf{p}_0|,$$

and we may use Proposition 7.1 with $-\psi_\varepsilon$ and argue exactly as in the previous case to obtain a contradiction.

Case $A_0 < 0 < B_0$. Then

$$q_0 + \beta \leq \mathcal{F}^*(\mathbf{p}_0, \mathbf{A}_0) = -\rho U|\mathbf{p}_0| = \mathcal{F}_*(\mathbf{p}_0, \mathbf{B}_0) = \mathcal{F}_*(\mathbf{p}_0, \mathbf{B}_0) \leq q_0,$$

which once again contradicts the positivity of β . \square

8. PROOF OF THEOREMS 3.1 AND 3.2.

Proof of Theorem 3.2. The uniqueness of a level-set solution of (1.1) corresponding to an auxiliary function Φ_0 follows from Theorem 3.3.

Let $\{\theta_1, \theta_2, \dots, \theta_M\}$ be the set of points of discontinuity of G . For n a sufficiently large positive integer, let G_n be the continuous 2π -periodic function with $G_n(\theta) = G(\theta)$ for $|\theta - \theta_k| \geq 1/n$, $k=1, 2, \dots, M$, and $G_n(\theta)$ linear otherwise. Further, let \mathcal{F}_n denote the function defined by (3.5) with G replaced by G_n . Then \mathcal{F}_n approximates \mathcal{F} in the sense of the following lemma, whose proof we omit.

Lemma 8.1. *Let $(p_n, A_n) \rightarrow (p, A) \in \mathbb{R}^2 \times S$ as $n \rightarrow \infty$. Then*

$$\limsup_{n \rightarrow \infty} (\mathcal{F}_n)^*(p_n, A_n) \leq \mathcal{F}^*(p, A),$$

$$\liminf_{n \rightarrow \infty} (\mathcal{F}_n)_*(p_n, A_n) \geq \mathcal{F}_*(p, A).$$

Since G_n is continuous, we may use [CGG, Theorem 6.8] to conclude that there is a unique, continuous viscosity solution $\Phi_n \in \hat{M}([0, \infty) \times \mathbb{R}^2)$ — of (3.4) with \mathcal{F} replaced by \mathcal{F}_n — satisfying $\Phi_n(\mathbf{x}, 0) = \Phi_0(\mathbf{x})$, and we define Φ^+ and Φ^- as in the proof of Theorem 5.2. Moreover, Lemma 8.1 together with classical stability results for viscosity solutions [FS, §2.6, §7.4] imply that Φ^+ and Φ^- are, respectively, a viscosity subsolution and a viscosity supersolution of (3.4) on $(0, \infty) \times \mathbb{R}^2$. We now follow the steps 4°–8° in the proof of Theorem 5.2 to conclude that $\Phi^+ = \Phi^- = \Phi$. Hence Φ is a level-set solution of (1.1) corresponding to Φ_0 .

We complete the proof by establishing (5.7). Let Φ be a level-set solution of (1.1) corresponding to an auxiliary function Φ_0 .

For $\delta > 0$, let $\eta_\delta : \mathbb{R} \rightarrow \mathbb{R}$ be smooth and satisfy: (i) $\eta_\delta' \geq 0$; (ii) $\eta_\delta(r) = 0$ for $r \leq c$; (iii) $\eta_\delta(r) = 1$ for $r \geq c + \delta$. Then the geometric property (3.12) implies that

$$\Phi_\delta(t, \mathbf{x}) = \eta_\delta(\Phi(t, \mathbf{x}))$$

is a level-set solution of (1.1).²⁴

Next,

$$u^+(t, \mathbf{x}) = \limsup_{\substack{\delta \downarrow 0 \\ (s, \mathbf{y}) \rightarrow (t, \mathbf{x})}} \Phi_\delta(s, \mathbf{y})$$

is a viscosity subsolution of (3.4) (cf. [FS, §2.6, §7.4]). If we let $\hat{u}(t, \mathbf{x})$ be the characteristic function of

$$\hat{U}(t) = \{ \mathbf{x} : \Phi(t, \mathbf{x}) \geq c \},$$

then the continuity of Φ and the properties of η_δ yield

$$u^+(t, \mathbf{x}) = \hat{u}^*(t, \mathbf{x}) = \hat{u}(t, \mathbf{x}),$$

so that $\hat{U}(t)$ is a χ -subsolution of (1.1). In fact, since

$$\text{cl } \Omega_0 = \{ \mathbf{x} : \Phi_0(\mathbf{x}) \geq c \} = \{ \mathbf{x} : \limsup_{\substack{\delta \downarrow 0, \mathbf{y} \rightarrow \mathbf{x}}} \hat{u}(s, \mathbf{y}) = 1 \},$$

$\hat{U}(t)$ is a χ -subsolution of (1.1) compatible with Ω_0 .

Similarly,

$$u^-(t, \mathbf{x}) = \liminf_{\substack{\delta \downarrow 0 \\ (s, \mathbf{y}) \rightarrow (t, \mathbf{x})}} \Phi_\delta(s, \mathbf{y})$$

is a viscosity supersolution of (3.4), and, further, $u^- = \hat{u}_*$; hence $\hat{U}(t)$ is also χ -supersolution of (1.1) compatible with Ω_0 . Thus $\hat{U}(t)$ is a χ -solution of (1.1) compatible with Ω_0 .

Next, in view of the definition $U(t)$, $\hat{U}(t) \subseteq U(t)$. In fact, they are equal. To verify this, let $\Omega(t)$ be a χ -subsolution compatible with Ω_0 , and let $u(t, \mathbf{x})$ be the characteristic function of $\Omega(t)$. For any $d < c$, let

²⁴Cf. [CGG, Theorem 5.6] for the proof of this fact when G is continuous.

$u(t, \mathbf{x}; d)$ be the characteristic function of

$$\mathcal{L}(t; d) = \{ \mathbf{x} : \Phi(t, \mathbf{x}) > d \}.$$

Then $u^*(0, \mathbf{x}) \leq u_*(0, \mathbf{x}; d) = u(0, \mathbf{x}; d)$, since $\Omega(t)$ is compatible with Ω_0 and $\Phi_0(\mathbf{x}) = \Phi(0, \mathbf{x})$ is an auxiliary function for Ω_0 . Then, by Corollary 3.1, $\Omega(t) \subset \mathcal{L}(t; d)$ for any χ -subsolution $\Omega(t)$ of (1.1) compatible with Ω_0 . Hence $u(t) \subseteq \mathcal{L}(t; d)$ for all $d < c$, and, since the intersection over all such d of $\mathcal{L}(t; d)$ is $\hat{U}(t)$, $u(t) \subseteq \hat{U}(t)$. Thus $u(t) = \hat{U}(t)$.

The analogous assertion for $\mathcal{L}(t)$ is proved in the same manner. \square

Proof of Theorem 3.1. Parts (a) and (b) follow from Theorem 3.2. To prove (c) note first that, since $\partial\Omega_0$ is C^3 , there is a C^3 parametrization $\alpha \mapsto Y(\alpha)$ ($\mathbb{R} \rightarrow \mathbb{R}^2$), periodic with period 1, such that

$$\partial\Omega_0 = \{ Y(\alpha) : \alpha \in [0, 1] \}$$

and

$$Y'(\alpha) / |Y'(\alpha)| = T(\theta_0(\alpha)),$$

where $\theta_0(\alpha)$ the normal-angle at $Y(\alpha)$ and $T(\theta)$ is defined in (1.10).

Proceeding formally, let $\Omega(t)$ be a solution of (1.1) such that

$$X(t, \alpha) = Y(\alpha) + h(t, \alpha)N(\theta_0(\alpha))$$

is a parametrization²⁵ of $\partial\Omega(t)$ for some real-valued function $h(t, \alpha)$. Then

(8.1) $h(t, \alpha)$ is periodic in α with period 1.

Assume that h is C^2 . Let $K_0(\alpha)$ denote the curvature of $\partial\Omega_0$ at $Y(\alpha)$, and let $\theta(t, \alpha)$, $V(t, \alpha)$, and $K(t, \alpha)$ denote the normal angle, normal

²⁵A similar parametrization was used by Chen and Reitich [CR] in their proof of local existence for a modified Stefan problem.

velocity, and curvature of $\partial\Omega(t)$ at $\mathbf{X}(t,\alpha)$. Then

$$\mathbf{T}(\theta(t,\alpha)) = \mathbf{X}_\alpha(t,\alpha) / |\mathbf{X}(t,\alpha)|$$

(where the subscript denotes differentiation with respect to that variable), and defining

$$F_1(\alpha,h,h_\alpha) = [F_2(\alpha,h)^2 + h_\alpha^2]^{1/2},$$

$$F_2(\alpha,h) = |\mathbf{Y}'(\alpha)| - hK_0(\alpha),$$

a tedious computation yields

$$\theta(t,\alpha) = \theta(\alpha,h(t,\alpha),h_\alpha(t,\alpha)),$$

with $\theta(\alpha,h,h_\alpha)$ the solution of

$$(8.2) \quad \mathbf{T}(\theta(\alpha,h,h_\alpha)) = [F_2(\alpha,h)\mathbf{T}(\theta_0(\alpha)) + h_\alpha\mathbf{N}(\theta_0(\alpha))]F_1(\alpha,h,h_\alpha)^{-1},$$

and

$$(8.3a) \quad \mathbf{V}(t,\alpha) = \mathbf{X}_t(t,\alpha) \cdot \mathbf{N}(\theta(t,\alpha)) = h_t F_2(\alpha,h) / F_1(\alpha,h,h_\alpha),$$

$$(8.3b) \quad \begin{aligned} K(t,\alpha) &= \theta_\alpha(t,\alpha) / |\mathbf{X}_\alpha(t,\alpha)| = F_3(\alpha,h,h_\alpha,h_{\alpha\alpha}) \\ &= \{ F_2(\alpha,h)[h_{\alpha\alpha} + hK_0(\alpha)^2] + K_0(\alpha)|\mathbf{Y}'(\alpha)|^3 + \\ &\quad h_\alpha[2K_0(\alpha)h_\alpha + K_0'(\alpha)h] - \mathbf{T}(\theta_0(\alpha)) \cdot \mathbf{Y}''(\alpha)h_\alpha - \\ &\quad \mathbf{N}(\theta_0(\alpha)) \cdot \mathbf{Y}''(\alpha)hK_0(\alpha) \} F_1(\alpha,h,h_\alpha)^{-3}, \end{aligned}$$

(Note that $\theta(t,\alpha)$ is well defined provided the right side of (8.2) is nonzero.) Thus, since

$$\mathbf{B}(\theta(t,\alpha))\mathbf{V}(t,\alpha) = \mathbf{G}(\theta(t,\alpha))K(t,\alpha) - \mathbf{U},$$

$h(t,\alpha)$ satisfies

$$(8.4) \quad \tilde{B}(\alpha, h, h_\alpha) h_t = \tilde{G}(\alpha, h, h_\alpha) h_{\alpha\alpha} - \tilde{F}(\alpha, h, h_\alpha),$$

with

$$\begin{aligned} \tilde{B}(\alpha, h, h_\alpha) &= B(\theta(\alpha, h, h_\alpha)), \\ \tilde{G}(\alpha, h, h_\alpha) &= G(\theta(\alpha, h, h_\alpha)) F_1(\alpha, h, h_\alpha)^{-2}, \\ \tilde{F}(\alpha, h, h_\alpha) &= \tilde{G}(\alpha, h, h_\alpha) \{ h K_0(\alpha)^2 + F_2(\alpha, h)^{-1} (K_0(\alpha) |Y'(\alpha)|^3 + \\ &\quad h_\alpha [2K_0(\alpha) h_\alpha + K_0'(\alpha) h] - T(\theta_0(\alpha)) \cdot Y''(\alpha) h_\alpha - \\ &\quad N(\theta_0(\alpha)) \cdot Y''(\alpha) h K_0(\alpha)) \} - U F_1(\alpha, h, h_\alpha) F_2(\alpha, h)^{-1}. \end{aligned}$$

We will complete the proof by solving (8.4) subject to $h(\alpha, 0) = 0$. Let

$$G = \{ (\alpha, h) : |h| \leq |Y'(\alpha)|/2 |K_0(\alpha)| \}.$$

Then

$$|Y'(\alpha)| - h K_0(\alpha) \geq |Y'(\alpha)|/2 > 0.$$

Hence the right side of (8.2) is nonzero and $\theta(\alpha, h, h_\alpha)$ is continuous on $G \times \mathbb{R}$. Moreover, $\tilde{F} : G \times \mathbb{R} \rightarrow \mathbb{R}$ is continuous; $\tilde{B} : G \times \mathbb{R} \rightarrow [0, \infty)$ is continuous and strictly positive; $\tilde{G} : G \times \mathbb{R} \rightarrow [0, \infty)$ is continuous except at finitely two-dimensional manifolds in $G \times \mathbb{R}$, and suffers at most jump discontinuities across such manifolds.

Although \tilde{G} has discontinuities, one can prove a comparison result for viscosity sub and supersolutions of (8.4) using a modification of the analysis given in Section 7. Indeed, requisite modifications of all arguments except Proposition 7.1 are either straightforward or minor, and Proposition 7.1 should be replaced by

Proposition 8.1. *Let $v(t, \alpha)$ be semiconvex on $[0, \infty) \times \mathbb{R}$ and differentiable at (t_0, α_0) with $v_\alpha(t_0, \alpha_0) \neq 0$. Then there are $(t_n, \alpha_n) \rightarrow (t_0, \alpha_0)$ and $(q_n, p_n, a_n) \in cD^+v(t_n, \alpha_n) \subset \mathbb{R}^3$ such that*

$$\lim_{n \rightarrow \infty} (q_n, p_n) = (v_t, v_\alpha)(t_0, \alpha_0) = (q_0, p_0),$$

$$\liminf_{n \rightarrow \infty} \min \{ \theta(\alpha_n, h_n, p_n) - \theta(\alpha_0, h_0, p_0), F_3(\alpha_n, h_n, p_n, a_n) \} \leq 0,$$

where $h_n = v(t_n, \alpha_n)$, $h_0 = v(t_0, \alpha_0)$, and F_3 is defined in (8.3).

Once a comparison result has been obtained, the existence of a unique viscosity solution h of (8.4), satisfying (8.1) and an initial condition for $h(t, 0)$, can be established utilizing an approximation argument of the type used in the proof of Theorem 3.1. This solution is defined on $[0, \hat{T}_{\max}] \times \mathbb{R}$, where \hat{T}_{\max} is the largest time satisfying $(\alpha, h(t, \alpha)) \in G$ for all $(t, \alpha) \in [0, \hat{T}_{\max}] \times \mathbb{R}$.

Let

$$\varepsilon_0 = \inf_{\alpha \in \mathbb{R}} \{ |Y'(\alpha)| / 4 |K_0(\alpha)| \},$$

and for $|\varepsilon| \leq \varepsilon_0$, let $h(t, \alpha; \varepsilon)$, $(t, \alpha) \in [0, \hat{T}_{\max}(\varepsilon)] \times \mathbb{R}$, be the unique viscosity solution of (8.4) satisfying (8.1) and $h(0, \alpha; \varepsilon) = \varepsilon$. The uniqueness associated with such solutions ensures that $h(t, \alpha; \varepsilon)$ depends continuously on ε . Our next step will be to show that

$$T_* := \inf \{ \hat{T}_{\max}(\varepsilon) : |\varepsilon| \leq \varepsilon_0 \}$$

satisfies

$$0 < T_* \leq T_{\text{uniq}}.$$

To verify this assertion, define, for $(t, \mathbf{x}) \in [0, T_*] \times \mathbb{R}^2$,

$$\varphi(t, \mathbf{x}) = \begin{cases} \varepsilon & \text{if } \mathbf{x} \in \partial\Omega(t; \varepsilon), |\varepsilon| \leq \varepsilon_0, \\ \varepsilon_0 & \text{if } \mathbf{x} \in \partial\Omega(t; \varepsilon_0), \\ -\varepsilon_0 & \text{if } \mathbf{x} \notin \partial\Omega(t; -\varepsilon_0), \end{cases}$$

where $\Omega(t; \varepsilon)$ is the closed region enclosed by

$$(8.5) \quad \{ Y(\alpha) + h(t, \alpha; \varepsilon)N(\theta_0(\alpha)) : \alpha \in [0, 1] \}.$$

(Since $t \leq T_*$, $(\alpha, h(t, \alpha; \varepsilon)) \in G$ and the curve (8.5) encloses a region.) Further, a tedious calculation shows that φ is a level-set solution of (1.1) corresponding to an auxiliary function compatible with Ω_0 . By Theorem 3.2 (cf. (5.7)),

$$U(t) = \{ \mathbf{x} : \Phi(t, \mathbf{x}) \geq 0 \}, \quad L(t) = \{ \mathbf{x} : \Phi(t, \mathbf{x}) > 0 \}.$$

Since h depends continuously on ε ,

$$T_{\text{uniq}} \geq T^*.$$

To establish the positivity of T^* , observe that, by the maximum principle (or comparison result for (8.4)),

$$|h(t, \alpha; \varepsilon) - \varepsilon| \leq \kappa t$$

for all $|\varepsilon| \leq \varepsilon_0$, $t \in [0, \hat{T}_{\max}(\varepsilon))$, where κ is a suitable constant depending on the C^3 norm of $\partial\Omega_0$. Hence

$$|h(t, \alpha; \varepsilon)| \leq 2\varepsilon_0$$

for all $t \leq \varepsilon_0/\kappa$. Finally, the definitions of G and ε_0 imply that $T^* \geq \varepsilon_0/\kappa$.

□

Acknowledgment. This work was supported by the Army Research Office and the National Science Foundation.

REFERENCES

- [AL] Abresch U. and J. Langer, The normalized curve shortening flow and homothetic solutions, *J. Diff. Geom.* **23**, 175-196 (1986)
- [AC] Allen, S. M., and J. W. Cahn, A macroscopic theory for antiphase boundary motion and its application to antiphase domain coarsening, *Act. Metall.* **27**, 1085-1098 (1979)
- [ATW] Almgren, F., J. E. Taylor, and L. Wang, Curvature driven flows: a variational approach, *SIAM J. Control Opt.* (issue dedicated to W. Fleming), Forthcoming
- [Ag] Angenent, S., Parabolic equations for curves on surfaces. I. Curves with p -integrable curvature, *Ann. Math.* **132**, 451-483 (1990); II. Intersections, blowup, and generalized solutions, *Ann. Math.* **133**, 171-217 (1991)
- [AG1] Angenent, S. and M. E. Gurtin, Multiphase thermomechanics with interfacial structure. 2. Evolution of an isothermal interface, *Arch. Rational. Mech. Anal.* **108**, 323-391 (1989)
- [AG2] Angenent, S. and M. E. Gurtin, Anisotropic motion of a phase interface. Well-posedness of the initial-value problem and qualitative properties of the interface, *J. reine angew. Math.* Forthcoming
- [Ba] Barles, G., Remark on a flame propagation model, *Rapport INRIA* #464 (1985)
- [Br] Brakke, K. A., *The Motion of a Surface by its Mean Curvature*, Princeton University Press (1978)
- [BSS] Barles, G., H. M. Soner, and P. E. Souganidis, Front propagation and phase field theory, *SIAM J. Cont. Optim.* (issue dedicated to W. Fleming), Forthcoming
- [Ch] Chen, X. Generation and propagation of the interface for reaction diffusion equations, *J. Diff. Eqts.* **96**, 116-141 (1992)
- [CR] Chen, X. and F. Reitich, Local existence and uniqueness of solutions of the Stefan problem with surface tension and kinetic undercooling, *J. Math. Anal. Appl.* **164**, 352-362 (1992)
- [CGG] Chen, Y.-G., Y. Giga, and S. Goto, Uniqueness and existence of viscosity solutions of generalized mean curvature flow equations, *J. Diff. Geom.* **33**, 749-786 (1991)
- [CH] Cahn, J. W. and D. W. Hoffman, A vector thermodynamics for anisotropic surfaces 2: curved and faceted surfaces, *Act. Metall.* **22**, 1205-1214 (1974).
- [Cr] Crandall, M. G., Quadratic forms, semidifferentials and viscosity solutions of fully nonlinear elliptic equations, *Ann. IHP Anal. Nonlin.*, **6**, 419-435 (1989)
- [CEL] Crandall, M. G., Evans, L. C., and P.-L. Lions, Some properties of viscosity solutions of Hamilton-Jacoby equations, *Trans. AMS*, **282**, 487-502 (1984)

- [CIL] Crandall, M. G., Ishii, I., and P.-L. Lions, User's guide to viscosity solutions of second-order partial differential equations, *Bul. AMS*, 27/1, 1-67 (1992)
- [CL] Crandall, M. G. and P.-L. Lions, Viscosity solutions of Hamilton-Jacoby equations, *Trans. AMS*, 277, 1-43 (1983)
- [DS] DeMottoni, P. and M. Schatzman, Development of surfaces in \mathbb{R}^N , *Proc. Roy. Soc. Edinburgh*, 116A, 207-220 (1990)
- [DGP] DiCarlo, A., Gurtin M. E, and P. Podio-Guidugli, A regularized equation for anisotropic motion-by-curvature, *SIAM J. Appl. Math.* 52, 1111-1119 (1992)
- [ESS] Evans, L. C., Soner, H. M., and P. E. Souganidis, Phase transitions and generalized motion by mean curvature, *Comm Pure Appl. Math.* 45, 1097-1123 (1992)
- [ES1] Evans, L. C. and J. Spruck, Motion of level sets by mean curvature I, *J. Diff. Geom.*, 33, 635-681 (1991)
- [ES2] Evans, L. C. and J. Spruck, Motion of level sets by mean curvature II, *Trans. AMS*, Forthcoming
- [ES3] Evans, L. C. and J. Spruck, Motion of level sets by mean curvature III, *J. Geom. Anal.* Forthcoming
- [FS] Fleming, W. and H. M. Soner, *Controlled Markov Processes and Viscosity Solutions*, Springer-Verlag, Berlin (1993)
- [GH] Gage, M. and R. S. Hamilton, The heat equation shrinking convex plane curves, *J. Diff. Geom.* 23, 69-95 (1986)
- [GS] Giga, Y. and M. H. Sato, Generalized interface condition with the Neumann boundary condition, *Proc. Japan Acad., Ser. A Math.* 67, 263-266 (1991)
- [Gj] Gjostein, N. A., Adsorption and surface energy 2: thermal faceting from minimization of surface energy, *Act. Metall.* 11, 969-977 (1963).
- [Gr] Grayson, M. A., The heat equation shrinks embedded plane curves to round points, *J. Diff. Geom.* 26, 285-314 (1987)
- [G1] Gurtin, M. E., Multiphase thermomechanics with interfacial structure. 1. Heat conduction and the capillary balance law, *Arch. Rational Mech. Anal.* 104, 185-221 (1988).
- [G2] Gurtin, M. E., *Thermomechanics of Evolving Phase Boundaries in the Plane*, Oxford Press, Forthcoming.
- [Je] Jensen, R. The maximum principle for viscosity solutions of second-order fully nonlinear partial differential equations, *Arch. Rational Mech. Anal.* 101, 1-27 (1988)
- [LL] Lasry, J. M., and P.-L. Lions, A remark on regularization in Hilbert spaces, *Israel J. Math.* 55, 257-266 (1986)
- [OS] Osher, S. and J. A. Sethian, Fronts propagating with curvature dependent speed, *J. Comp. Phys.* 79, 12-49 (1988)
- [ORS] Owen, N. C., Rubinstein, J., and P. Sternberg, Minimizers and gradient flows for singularly perturbed bi-stable potentials with a

- Dirichlet condition, *Proc. Roy. Soc. London* 429, 505-532 (1989)
- [OwS] Owen, N. C. and P. Sternberg, Gradient flow and front propagation with boundary contact energy, Forthcoming
- [RS] Rubinstein, J. and P. Sternberg, Nonlocal reaction-diffusion equations and nucleation, *IMA J. Appl. Math.*, Forthcoming
- [Se] Sethian, J. A., Curvature and evolution of fronts, *Comm. Math. Phys.* 101, 487-499 (1985)
- [So] Soner, H. M., Motion of a set by the mean curvature of its boundary, *J. Diff. Eqts.*, Forthcoming
- [SZ] Sternberg, P. and W. Ziemer, Generalized motion by curvature with a Dirichlet boundary condition, Forthcoming
- [TCH] Taylor, J., J. W. Cahn, and C. A. Handwerker, Geometric models of crystal growth 1, *Acta Met.* 40, 1443-1474 (1992)

References, continued

[BK] L. Bronsard and R. Kohn, Motion by mean curvature as the singular limit of Ginzburg-Landau model, *Jour. Diff. Equations*, 90 (1991), 211-237.

[JLS] R. Jensen, P.-L. Lions and P.E. Souganidis, A uniqueness result for viscosity solutions of second order fully nonlinear partial differential equations, *Proc. AMS.*, 102 (1988), 975-978.

[OS] M. Ohnuma and M. Sato, Singular degenerate parabolic equations with applications to geometric evolutions, (1992) preprint.

[RSK] J. Rubinstein, P. Sternberg and J.B. Keller, Fast reaction, slow diffusion and curve shortening, *SIAM J. Appl. Math.*, 49 (1989), 116-133.

- Dirichlet condition, *Proc. Roy. Soc. London* 429, 505-532 (1989)
- [OwS] Owen, N. C. and P. Sternberg, Gradient flow and front propagation with boundary contact energy, Forthcoming
- [RS] Rubinstein, J. and P. Sternberg, Nonlocal reaction-diffusion equations and nucleation, *IMA J. Appl. Math.*, Forthcoming
- [Se] Sethian, J. A., Curvature and evolution of fronts, *Comm. Math. Phys.* 101, 487-499 (1985)
- [So] Soner, H. M., Motion of a set by the mean curvature of its boundary, *J. Diff. Eqts.*, Forthcoming
- [SZ] Sternberg, P. and W. Ziemer, Generalized motion by curvature with a Dirichlet boundary condition, Forthcoming
- [TCH] Taylor, J., J. W. Cahn, and C. A. Handwerker, Geometric models of crystal growth 1, *Acta Met.* 40, 1443-1474 (1992)

**Center for Nonlinear Analysis
Report Series • Complete List**

Nonlinear Analysis Series

No.

- 91-NA-001 [] Lions, P.L., **Jacobians and Hardy spaces**, June 1991
- 91-NA-002 [] Giga, Y. and Soto, M.-H., **Generalized interface evolution with the Neumann boundary condition**, July 1991
- 91-NA-003 [] Soner, H.M. and Souganidis, P.E., **Uniqueness and singularities of cylindrically symmetric surfaces moving by mean curvature**, July 1991
- 91-NA-004 [] Coleman, B.D., Marcus, M. and Mizel, V.J., **On the Thermodynamics of periodic phases**, August 1991
- 91-NA-005 [] Gurtin, M.E. and Podio-Guidugli, P., **On the formulation of mechanical balance laws for structured continua**, August 1991
- 91-NA-006 [] Gurtin, M.E. and Voorhees, P., **Two-phase continuum mechanics with mass transport and stress**, August 1991
- 91-NA-007 [] Fried, E., **Non-monotonic transformation kinetics and the morphological stability of phase boundaries in thermoelastic materials**, September 1991
- 91-NA-008 [] Gurtin, M.E., **Evolving phase boundaries in deformable continua**, September 1991
- 91-NA-009 [] Di Carlo, A., Gurtin, M.E., and Podio-Guidugli, P., **A regularized equation for anisotropic motion-by-curvature**, September 1991
- 91-NA-010 [] Kinderlehrer, D. and Ou, B., **Second variation of liquid crystal energy at $x/|x|$** , August 1991
- 91-NA-011 [] Baughman, L.A. and Walkington, N., **Co-volume methods for degenerate parabolic problems**, August 1991
- 91-NA-012 [] James, R.D. and Kinderlehrer, D., **Frustration and microstructure: An example in magnetostriction**, November 1991
- 91-NA-013 [] Angenent, S.B. and Gurtin, M.E., **Anisotropic motion of a phase interface**, November 1991

- 92-NA-001 [] Nicolaidis, R.A. and Walkington, N.J., **Computation of microstructure utilizing Young measure representations**, January 1992
- 92-NA-002 [] Tartar, L., **On mathematical tools for studying partial differential equations of continuum physics: H-measures and Young measures**, January 1992
- 92-NA-003 [] Bronsard, L. and Hilhorst, D., **On the slow dynamics for the Cahn-Hilliard equation in one space dimension**, February 1992
- 92-NA-004 [] Gurtin, M.E., **Thermodynamics and the supercritical Stefan equations with nucleations**, March 1992
- 92-NA-005 [] Antonic, N., **Memory effects in homogenisation linear second order equation**, February 1992
- 92-NA-006 [] Gurtin, M.E. and Voorhees, P.W., **The continuum mechanics of coherent two-phase elastic solids with mass transport**, March 1992
- 92-NA-007 [] Kinderlehrer, D. and Pedregal, P., **Remarks about gradient Young measures generated by sequences in Sobolev spaces**, March 1992
- 92-NA-008 [] **Workshop on Shear Bands**, March 23-25, 1992 (Abstracts), March 1992
- 92-NA-009 [] Armstrong, R. W., **Microstructural/Dislocation Mechanics Aspects of Shear Banding in Polycrystals**, March 1992
- 92-NA-010 [] Soner, H. M. and Souganidis, P. E., **Singularities and Uniqueness of Cylindrically Symmetric Surfaces Moving by Mean Curvature**, April 1992
- 92-NA-011 [] Owen, David R., Schaeffer, Jack, and Wang, Keming, **A Gronwall Inequality for Weakly Lipschitzian Mappings**, April 1992
- 92-NA-012 [] Alama, Stanley and Li, Yan Yan, **On "Multibump" Bound States for Certain Semilinear Elliptic Equations**, April 1992
- 92-NA-013 [] Olmstead, W. E., Nemat-Nasser, S., and Li, L., **Shear Bands as Discontinuities**, April 1992
- 92-NA-014 [] Antonic, N., **H-Measures Applied to Symmetric Systems**, April 1992
- 92-NA-015 [] Barroso, Ana Cristina and Fonseca, Irene, **Anisotropic Singular Perturbations - The Vectorial Case**, April 1992
- 92-NA-016 [] Pedregal, Pablo, **Jensen's Inequality in the Calculus of Variations**, May 1992
- 92-NA-017 [] Fonseca, Irene and Muller, Stefan, **Relaxation of Quasiconvex Functionals in $BV(\Omega, \mathbb{R}^p)$ for Integrands $f(x, u, \nabla u)$** , May 1992

- 92-NA-018 [] **Alama, Stanley and Tarantello, Gabriella, On Semilinear Elliptic Equations with Indefinite Nonlinearities, May 1992**
- 92-NA-019 [] **Owen, David R., Deformations and Stresses With and Without Microslip, June 1992**
- 92-NA-020 [] **Barles, G., Soner, H. M., Souganidis, P. E., Front Propagation and Phase Field Theory, June 1992**
- 92-NA-021 [] **Bruno, Oscar P. and Reitich, Fernando, Approximation of Analytic Functions: A Method of Enhanced Convergence, July 1992**
- 92-NA-022 [] **Bronsard, Lia and Reitich, Fernando, On Three-Phase Boundary Motion and the Singular Limit of a Vector-Valued Ginzburg-Landau Equation, July 1992**
- 92-NA-023 [] **Cannarsa, Piermarco, Gozzi, Fausto and Soner, H.M., A Dynamic Programming Approach to Nonlinear Boundary Control Problems of Parabolic Type, July 1992**
- 92-NA-024 [] **Fried, Eliot and Gurtin, Morton, Continuum Phase Transitions With An Order Parameter; Accretion and Heat Conduction, August 1992**
- 92-NA-025 [] **Swart, Pieter J. and Homes, Philip J., Energy Minimization and the Formation of Microstructure in Dynamic Anti-Plane Shear, August 1992**
- 92-NA-026 [] **Ambrosio, I., Cannarsa, P. and Soner, H.M., On the Propagation of Singularities of Semi-Convex Functions, August 1992**
- 92-NA-027 [] **Nicolaidis, R.A. and Walkington, Noel J., Strong Convergence of Numerical Solutions to Degenerate Variational Problems, August 1992**
- 92-NA-028 [] **Tarantello, Gabriella, Multiplicity Results for an Inhomogenous Neumann Problem with Critical Exponent, August 1992**
- 92-NA-029 [] **Noll, Walter, The Geometry of Contact, Separation, and Reformation of Continuous Bodies, August 1992**
- 92-NA-030 [] **Brandon, Deborah and Rogers, Robert C., Nonlocal Superconductivity, July 1992**
- 92-NA-031 [] **Yang, Yisong, An Equivalence Theorem for String Solutions of the Einstein-Matter-Gauge Equations, September 1992**
- 92-NA-032 [] **Spruck, Joel and Yang, Yisong, Cosmic String Solutions of the Einstein-Matter-Gauge Equations, September 1992**
- 92-NA-033 [] **Workshop on Computational Methods in Materials Science (Abstracts), September 16-18, 1992.**

- 92-NA-034 [] **Leo, Perry H. and Heng-Jeng Jou, Shape evolution of an initially circular precipitate growing by diffusion in an applied stress field, October 1992**
- 92-NA-035 [] **Gangbo, Wilfrid, On the weak lower semicontinuity of energies with polyconvex integrands, October 1992**
- 92-NA-036 [] **Katsoulakis, Markos, Kossioris, Georgios T. and Retich, Fernando, Generalized motion by mean curvature with Neumann conditions and the Allen-Cahn model for phase transitions, October 1992**
- 92-NA-037 [] **Kinderlehrer, David, Some methods of analysis in the study of microstructure, October 1992**
- 92-NA-038 [] **Yang, Yisong, Self duality of the Gauge Field equations and the Cosmological Constant, November 1992**
- 92-NA-039 [] **Brandon, Deborah and Rogers, Robert, Constitutive Laws for Pseudo-Elastic Materials, November 1992**
- 92-NA-040 [] **Leo, P. H., Shield, T. W., and Bruno, O. P., Transient Heat Transfer Effects on the Pseudoelastic Behavior of Shape-Memory Wires, November 1992**
- 92-NA-041 [] **Gurtin, Morton E., The Dynamics of Solid-Solid Phase Transitions 1. Coherent Interfaces, November 1992**
- 92-NA-042 [] **Gurtin, Morton E., Soner, H. M., and Souganidis, P. E., Anisotropic Motion of an Interface Relaxed by the Formation of Infinitesimal Wrinkles, December 1992**

Stochastic Analysis Series

- 91-SA-001 [] **Soner, H.M., Singular perturbations in manufacturing, November 1991**
- 91-SA-002 [] **Bridge, D.S. and Shreve, S.E., Multi-dimensional finite-fuel singular stochastic control, November 1991**
- 92-SA-001 [] **Shreve, S. E. and Soner, H. M., Optimal Investment and Consumption with Transaction Costs, September 1992**

DEC 12 2003

Carnegie Mellon University Libraries



3 8482 01360 3580