NOTICE WARNING CONCERNING COPYRIGHT RESTRICTIONS:

The copyright law of the United States (title 17, U.S. Code) governs the making of photocopies or other reproductions of copyrighted material. Any copying of this document without permission of its author may be prohibited by law.

NAMT 94.015

Three-phase boundary motions under constant velocities.

I: The vanishing surface tension limit.

Fernando Reitich Carnegie Mellon University

H. Mete Soner Carnegie Mellon University

Research Report No. 94-NA-015

May 1994

Sponsors

U.S. Army Research Office Research Triangle Park NC 27709

National Science Foundation 1800 G Street, N.W. Washington, DC 20550 University Libration Carnegic Melian Charactery Pittsburgh PA 15213-3897

NAMT 94-015

1.

'n

μ

University Libraries Carpagie Metica University Pharmana (Action 2008)

.

•

ri

Three-phase boundary motions under constant velocities. I: The vanishing surface tension limit

Fernando Reitich

H. Mete Soner

Department of Mathematics Carnegie Mellon University Pittsburgh, PA 15213-3890

May 1993

÷

TI

Abstract

In this paper we deal with the dynamics of material interfaces such as solid-liquid, grain or antiphase boundaries. We concentrate on the situation in which these internal surfaces separate three regions in the material with different physical attributes (e.g. grain boundaries in a polycrystal). The basic two-dimensional model proposes that the motion of an interface Γ_{ij} between regions *i* and *j* $(i, j = 1, 2, 3, i \neq j)$ is governed by the equation

$$V_{ij} = \mu_{ij} \left(f_{ij} \kappa_{ij} + F_{ij} \right) \,. \tag{0.1}$$

Here V_{ij} , κ_{ij} , μ_{ij} and f_{ij} denote, respectively, the normal velocity, the curvature, the mobility and the surface tension of the interface and the numbers F_{ij} stand for the (constant) difference in bulk energies. At the point where the three phases coexist, local equilibrium requires that

the curves meet at prescribed angles.
$$(0.2)$$

In case the material constants f_{ij} are small, $f_{ij} = \epsilon \widehat{f_{ij}}$ and $\epsilon \ll 1$, previous analyses based on the parabolic nature of the equations (0.1) do not provide good qualitative information on the behavior of solutions. In this case it is more appropriate to consider the singular case with $f_{ij} = 0$. It turns out that this problem, (0.1) with $f_{ij} = 0$, admits infinitely many solutions. Here, we show that a unique solution, "the vanishing surface tension (VST) solution", is selected by letting $\epsilon \to 0$. Furthermore, we introduce the concept of *weak viscosity solution* for the problem with $\epsilon = 0$ and show that the VST solution coincides with the unique weak solution. Finally, we give examples showing that, in several cases of physical relevance, the VST solution *differs* from results proposed previously.

1991 Mathematics Subject Classifications: 65B10, 41A21, 41A25.

Keywords: Mean curvature flow, triple junction, viscosity solutions

.

Γ

1 Introduction

A variety of mathematical models have been devised to investigate the dynamics of interfaces such as solid-liquid, grain or antiphase boundaries. These internal surfaces are, in general, non-equilibrium features of a material: they have a positive excess free energy. Thus, for example, grain boundaries (i.e. boundaries between single crystals in a polycrystal) migrate to reduce the total amount of grain boundary area. Understanding the evolution of these surfaces is of fundamental importance, not only for its intrinsic interest, but also for its technological significance: they constitute a key factor in determining a wide range of material properties, from mechanical strength to electrical conductivity (see e.g. [3]).

The simplest models for interface dynamics can be obtained by neglecting the changes in the bulk and concentrating solely on the evolution of the internal boundaries. In many instances these kinds of models provide a good representation of the physics. In this paper we study a two-dimensional model for the motion of interfaces in multi-phase continua. This model, which applies in particular to grain growth, corresponds to the "small surface tension" limit of the one derived by Mullins [10] and investigated by Bronsard and Reitich [5]. We show here that, even though the "vanishing surface tension" problem admits infinitely many solutions, a unique solution, which we shall call "the vanishing surface tension (VST) solution" is selected by the limiting process (see §2). Furthermore, we introduce the concept of *weak viscosity solution* for the problem and show that the VST solution coincides with the unique weak solution.

Mullins' model describes the evolution of three curves in the plane (the twodimensional boundaries) which move according to

Normal Velocity =
$$V = \kappa$$
 = Curvature. (1.1)

The three curves are assumed to meet at a point, a "triple junction" (see Figure 1), with prescribed angles θ_i ($\theta_i = 120^\circ$ in the case of grain boundaries). More generally, (1.1) should be replaced by the equation

$$V = \mu f \kappa - \mu F \,, \tag{1.2}$$

which describes the evolution of an isotropic interface that is driven by an energy difference F between the bulk phases [2]. The material constants f and μ denote, respectively, the energy and mobility of the interface; they may be different for different curves. The existence and uniqueness of solutions for a given initial state was established in [5] by means of a parametric formulation of the problem and the analysis of the resulting quasi-linear parabolic system of equations with fully nonlinear boundary conditions. When $\mu f \ll 1$, the system is near degenerate and the analysis in [5] does not provide good qualitative information on the behavior of



Figure 1: The interfaces Γ_{ij} move with normal velocities $V_{ij} = \mu_{ij} (\epsilon f_{ij} \kappa_{ij} + e_i - e_j)$ and prescribed angles θ_i .

solutions. For this, it is more appropriate to look into the problem with $\mu f = 0$, that is a model in which the curves evolve under constant velocities.

The problem of three curves moving under constant velocities and meeting at a triple junction has been previously considered by Taylor [12] and Merriman, Bence and Osher [9]. It is readily checked that, if no further conditions on the motion are imposed, the solution is not unique (see Figure 2 below). In [12] a Huygens' Principle is applied in order to select a particular solution, and a catalog of solutions depending on the initial configuration is provided. The implementation of the level set approach developed in [9] yields solutions which coincide with those in [12]. As we shall see, these solutions do not always coincide with the VST or weak solutions we present here.

Perhaps the simplest way to motivate the model is to associate to the system an energy functional of the form

$$\mathcal{E} = \epsilon f_{12} \operatorname{length}(\Gamma_{12}) + \epsilon f_{23} \operatorname{length}(\Gamma_{23}) + \epsilon f_{31} \operatorname{length}(\Gamma_{31}) + e_1 \operatorname{vol}(\Omega_1) + e_2 \operatorname{vol}(\Omega_2) + e_3 \operatorname{vol}(\Omega_3).$$
(1.3)

Then, it easy to see that the corresponding gradient flow is

$$V_{ij} = \mu_{ij} \left(\epsilon f_{ij} \kappa_{ij} + e_i - e_j \right), \quad \epsilon \ll 1$$
(1.4)

T

where V_{ij} , κ_{ij} and μ_{ij} denote, respectively, the normal velocity, the curvature and the mobility of the interface between phase *i* and phase *j*. Furthermore, a straightforward calculation shows that, in the absence of dissipation at the triple junction,

the requirement that

$$\frac{d\mathcal{E}}{dt} \leq 0$$

imposes the condition

$$\frac{\sin(\theta_1)}{f_{23}} = \frac{\sin(\theta_2)}{f_{31}} = \frac{\sin(\theta_3)}{f_{12}} \quad (\text{see Figure 1}). \tag{1.5}$$

This condition coincides with the classical condition at triple points that can be found in the materials science literature (see e.g. [11]). It was recently shown in [5] that (1.5) is also recovered in the small interface-thickness limit of a (diffuseinterface) Allen-Cahn type model for three-phase boundary motion (see also [1]). Notice that in the particular case of grain boundaries we have

$$f_{12} = f_{23} = f_{31}$$

and (1.5) implies that $\theta_i = 120^{\circ}$.

The remainder of the paper is devoted to understanding the qualitative behavior of (1.4),(1.5) in the limit $\epsilon \to 0$. First, in §2, we discuss the non-uniqueness of solutions in the case $\epsilon = 0$. We present a perturbation analysis (§2.1) and a class of self-similar solutions (§2.2), which suggest that a unique solution (the VST solution) is selected by letting $\epsilon \to 0$ does not. In §3 we introduce the concept of weak viscosity solution for the constant-velocities problem ((1.4) with $\epsilon = 0$). Since the motion is governed by a system of Hamilton-Jacobi equations, the classical notion of viscosity solution can be used away from the triple points. At the triple points, however, and in accordance with (1.5), we need to introduce the idea of weak angle conditions. As we shall show, this concept singles out a unique solution which coincides with the one obtained in the limit $\epsilon \to 0$ (§4). Finally, in §5, we present a preliminary result of convergence of solutions as $\epsilon \to 0$ in the case where two interfaces are symmetric relative to the third (which is given by a semi-infinite stationary line). A simple example of a non-physical geometric problem is also discussed at the end of §5.

2 Non-uniqueness and the VST solution

As discussed in §1, for $\epsilon > 0$ the equations (1.4) subject to the angle conditions (1.5) admit (in a finite time interval) unique classical solutions for smooth initial data. Of course, when $\epsilon = 0$ the resulting (first order) equations

$$V_{ij} = \mu_{ij} \left(e_i - e_j \right) \tag{2.1}$$

cannot be constrained by the angle conditions. On the other hand, the equations (2.1) subject only to the requirement that the interfaces meet at a triple point, admit infinitely many solutions. The simplest case in which this is apparent corresponds to that of Figure 2, where

$$\mu_{ij} = 1$$

and

$$e_2 - e_1 = \alpha$$
 and $e_2 - e_3 = \alpha$ $(\alpha > 0)$.



Figure 2: (a) The solution proposed in [12]. (b) The VST solution.

Figure 2 contains two different solutions to the problem. The lighter curves correspond to the initial configuration and the heavier ones to the the solution at time t = 1. Since the interface velocities are constant, the solution at time t = T can be obtained by simply dilating these graphs. The solution in Figure 2(a) corresponds

T

to the construction in [12]. However, it is easily checked that the solution that is singled out by taking the limit as $\epsilon \to 0$ in (1.4) is the one depicted in Figure 2(b). Indeed, if the initial configuration satisfies the angle conditions, it is clear that the solution in Figure 2(b) is an exact solution of (1.4), (1.5) for $\epsilon > 0$ (since the interfaces have zero curvature) and therefore it coincides with the limit solution.



Figure 3: (a) The solution proposed in [12]. (b) The VST solution; one of the angle conditions is preserved in this solution.

More generally, consider the situation of Figure 3, where

$$e_2 - e_1 = \alpha$$
 and $e_2 - e_3 = \beta$ $(\alpha \ge \beta > 0)$.

Again, Figure 3(a) shows the solution proposed in [12] (at t = 1). The construction is as follows: first draw a parallel line to the initial interface $\Gamma_{12}(0)$ at a distance α , up to the point of tangency with the circle of radius α . Then, follow this circle up to the new position of the triple junction on the x-axis: this defines Γ_{12} at time t = 1. To construct Γ_{23} at time t = 1, first draw a parallel line to the initial interface $\Gamma_{23}(0)$ at a distance β up to the point of tangency with the circle of radius β and then a line segment from the triple junction which is tangent to this circle (notice that this introduces a corner in Γ_{23}). The construction in Figure 3(b) is similar but in building Γ_{12} we do not go all the way down to the x-axis on the circle of radius α . Rather, we propose a solution which, at t = 1, contains only part of this arc of circle and then joins Γ_{12} with Γ_{13} with a straight line segment which is tangent to the circle of radius α ; the ensuing construction of Γ_{23} is as in Figure 3(a). Notice that the length of the arc of circle in the solution is a free parameter, giving rise to infinitely many solutions. The solution we propose, in case

$$\theta_i = 120^o \tag{2.2}$$

and

$$0 < \beta < \alpha < 2\beta, \tag{2.3}$$

is the one where this length is exactly the one for which the construction results in an angle of 120° between Γ_{12} and Γ_{23} . In §2.1 and §2.2 we shall show, through a perturbation analysis and the construction of self-similar solutions, that this is indeed the solution that is selected if we let $\epsilon \to 0$ in (1.4), (2.2). Further evidence of this fact will be provided in §3 after we motivate and define the notion of viscosity solution to (2.1), (2.2). As we shall see, the solution depicted in Figure 3(b) is the unique viscosity solution of (2.1), (2.2).

2.1 Perturbation analysis

As we discussed above, the solution in Figure 2(b) is an exact solution of the problem

$$V_{21} = \epsilon f_{12} \kappa_{21} + \alpha ,$$

$$V_{23} = \epsilon f_{23} \kappa_{23} + \alpha ,$$

$$V_{13} = \epsilon f_{13} \kappa_{13}$$
(2.4)

subject to the angle conditions (2.2). In order to gain some insight into the structure of solutions, it is natural then to study small perturbations of this problem (see Figure 4):

$$V_{21} = \epsilon f_{12} \kappa_{21} + \alpha ,$$

$$V_{23} = \epsilon f_{23} \kappa_{23} + (\alpha - \delta) , \quad \delta \ll 1 ,$$

$$V_{13} = \epsilon f_{13} \kappa_{13} .$$
(2.5)

T

In the coordinates of Figure 4, the equations (2.5) can be written in the form

$$u_{t} = \epsilon f_{12} \frac{u_{xx}}{1 + u_{x}^{2}} + \alpha \sqrt{1 + u_{x}^{2}}, \ x < s(t)$$

$$v_{t} = \epsilon f_{23} \frac{v_{xx}}{1 + v_{x}^{2}} - (\alpha - \delta) \sqrt{1 + v_{x}^{2}}, \ x < s(t) \quad (\delta \ll 1), \qquad (2.6)$$

$$w_{t} = \epsilon f_{13} \frac{w_{xx}}{1 + w_{x}^{2}}, \ x > s(t)$$

. 7



Figure 4: Perturbation analysis: (a) Exact solution. (b) A perturbation.

where s(t) denotes the (unknown) x-coordinate of the triple junction at time t. The requirement that the solutions meet at the triple junction translates into

$$u(s(t),t) = w(s(t),t), v(s(t),t) = w(s(t),t),$$
(2.7)

and the angle conditions (2.2) become

$$1 + u_x(s(t), t)v_x(s(t), t) = -\frac{1}{2}\sqrt{1 + u_x(s(t), t)^2}\sqrt{1 + v_x(s(t), t)^2},$$

$$-1 - u_x(s(t), t)w_x(s(t), t) = -\frac{1}{2}\sqrt{1 + u_x(s(t), t)^2}\sqrt{1 + w_x(s(t), t)^2}.$$
 (2.8)

The exact solution for $\delta = 0$ is

$$u = u_0(x,t) = -\sqrt{3}x + 2\alpha t$$
,

$$v = v_0(x,t) = \sqrt{3}x - 2\alpha t, \qquad (2.9)$$
$$w = w_0(x,t) \equiv 0,$$
$$s = s_0(t) = \frac{2\alpha}{\sqrt{3}}t.$$

To first order in δ

$$u = u_0 + \delta u_1$$
, $v = v_0 + \delta v_1$, $w = w_0 + \delta w_1$, $s = s_0 + \delta s_1$

and the linearized version of the free boundary problem (2.6), (2.7), (2.8) is

$$\begin{aligned} u_{1t}(x,t) &= \epsilon \frac{f_{12}}{4} u_{1xx}(x,t) - \alpha \frac{\sqrt{3}}{2} u_{1x}(x,t), \ x < s_0(t), \\ v_{1t}(x,t) &= \epsilon \frac{f_{23}}{4} v_{1xx}(x,t) - \alpha \frac{\sqrt{3}}{2} v_{1x}(x,t) + 2, \ x < s_0(t), \\ w_{1t}(x,t) &= \epsilon f_{13} w_{1xx}(x,t), \ x > s_0(t), \\ u_{1x}(s_0(t),t) &= v_{1x}(s_0(t),t) \\ u_{1x}(s_0(t),t) &= 4 w_{1x}(s_0(t),t) \\ u_{1}(s_0(t),t) &= v_{1}(s_0(t),t) + 2\sqrt{3}s_1(t) \\ u_{1}(s_0(t),t) &= w_{1}(s_0(t),t) + \sqrt{3}s_1(t). \end{aligned}$$

$$(2.10)$$

At this point, the equation (2.10) becomes the key to understand the behavior of solutions as $\epsilon \to 0$. Indeed, the condition

$$u_{1x}(s_0(t),t) = v_{1x}(s_0(t),t)$$

implies that, near the triple junction, the solutions u and v for $\delta \ll 1$ differ from the solution for $\delta = 0$ by a rotation of angle

$$\theta_R \approx -\frac{u_{1x}}{4}\delta \quad (\delta \ll 1).$$
(2.11)

In particular, the angle between u and v is preserved in the limit, i.e.

$$Angle(u, v) = 120^{\circ}.$$
 (2.12)

T

Figure 5: A self-similar solution.

2.2 Self-similar solutions

In what follows we shall construct self-similar solutions $(u^{\epsilon}, v^{\epsilon}, w^{\epsilon})$ to (cf. (2.6) with $\alpha - \delta = \beta$)

$$\begin{aligned} u_t^{\epsilon} &= \epsilon f_{12} \frac{u_{xx}^{\epsilon}}{1 + (u_x^{\epsilon})^2} + \alpha \sqrt{1 + (u_x^{\epsilon})^2}, \ x < s^{\epsilon}(t) \\ v_t^{\epsilon} &= \epsilon f_{23} \frac{v_{xx}^{\epsilon}}{1 + (v_x^{\epsilon})^2} - \beta \sqrt{1 + (v_x^{\epsilon})^2}, \ x < s^{\epsilon}(t) \end{aligned}$$

$$\begin{aligned} w_t^{\epsilon} &= \epsilon f_{13} \frac{w_{xx}^{\epsilon}}{1 + (w_x^{\epsilon})^2}, \ x > s^{\epsilon}(t) \end{aligned}$$
(2.13)

subject to the free-boundary conditions (2.7), (2.8). As we shall see, these solutions will converge, as $\epsilon \to 0$, to a solution of

$$u_{t} = \alpha \sqrt{1 + u_{x}^{2}}, x < s(t)$$

$$v_{t} = -\beta \sqrt{1 + v_{x}^{2}}, x < s(t),$$

$$w_{t} = 0, x > s(t)$$
(2.14)

satisfying (2.12).

10

The first step in the construction is to assume

$$u^{\epsilon}(x,t) = u(x,t) = -a(x - s^{\epsilon}(t)), \quad a > 0,$$

$$v^{\epsilon}(x,t) = v(x,t) = b(x - s^{\epsilon}(t)), \quad b > 0,$$

$$s^{\epsilon}(t) = \sigma t, \quad \sigma > 0.$$
(2.15)

Then, equations (2.13) take on the form

$$\sigma a = \alpha \sqrt{1 + a^2},$$

$$\sigma b = \beta \sqrt{1 + b^2},$$
(2.16)

and the angle condition for $(u^{\epsilon}, v^{\epsilon})$ becomes (cf. (2.8))

$$1 - ab = -\frac{1}{2}\sqrt{1 + a^2}\sqrt{1 + b^2}.$$
(2.17)

Solving (2.16), (2.17) for (a, b, σ) we obtain

$$a = \frac{\sqrt{3}\alpha}{2\beta - \alpha}, \qquad b = \frac{\sqrt{3}\beta}{2\alpha - \beta},$$
$$\sigma = \frac{2}{\sqrt{3}}\sqrt{\alpha^2 + \beta^2 - \alpha\beta}.$$
(2.18)

Next, we want to solve for w^{ϵ} by proposing a self-similar form

$$w^{\epsilon}(x,t) = \epsilon \psi(\frac{x-\sigma t}{\epsilon})$$

Then, from (0.1)

$$-\sigma\psi'(z) = f_{13}\frac{\psi''(z)}{1+(\psi'(z))^2}, \quad z > 0, \qquad (2.19)$$

 \mathbf{and}

$$\psi(0) = 0. \tag{2.20}$$

.

T

The requirements that

$$Angle(u^{\epsilon}, w^{\epsilon}) = 120^{\circ}, \quad Angle(v^{\epsilon}, w^{\epsilon}) = 120^{\circ}$$

translate into

$$\psi'(0) = -\eta \quad (\eta > 0)$$
 (2.21)

where

$$\eta = \tan(\theta_R) = \frac{a - \sqrt{3}}{1 + a\sqrt{3}} = \frac{\sqrt{3} - b}{1 + b\sqrt{3}} = \sqrt{3} \frac{(\alpha - \beta)}{(\alpha + \beta)}.$$
 (2.22)

It is easily checked that the (unique) solution to (2.19) subject to the initial conditions (2.20) and (2.21) is given by

$$\psi(z) = \frac{f_{13}}{\sigma} \left[\arcsin\left(\frac{\eta}{\sqrt{1+\eta^2}} e^{-\sigma z/f_{13}}\right) - \arcsin\left(\frac{\eta}{\sqrt{1+\eta^2}}\right) \right].$$
(2.23)

With this definition, $(u^{\epsilon} = u, v^{\epsilon} = v, w^{\epsilon})$ is a self-similar solution of (0.1), (2.7) and (2.8) (see Figure 5). Since $\psi(z)$ is uniformly bounded, $(u^{\epsilon}, v^{\epsilon}, w^{\epsilon}) \rightarrow (u, v, 0)$ uniformly as $\epsilon \rightarrow 0$. In particular, the angle between u and v is again preserved in the limit, cf. (2.12), see Figure 6. Notice that the other angles *are not* preserved.

Figure 6: The limit $(\epsilon \rightarrow 0)$ of self-similar solutions (see Figure 5).

3 The weak solution

In §2.2 we have shown that, for $0 < \epsilon \ll 1$, the solutions of (1.4), (1.5) develop boundary layers at the triple junction and that in the limit as $\epsilon \to 0$ the angle conditions are not necessarily satisfied. In the theory of viscosity solutions this phenomenon is well understood and weak formulations of different types of boundary conditions have been developed [6, 7]. In this section we shall derive a weak formulation of (2.1), (1.5) in the spirit of those obtained in the theory of viscosity solutions. However, it should be noted here that this system of equations does not have a comparison principle –a property which is essential in the theory of viscosity solutions.

3.1 The definition

For each $t \ge 0$, consider three unbounded, closed and connected regions $\Omega_i(t)$, i = 1, 2, 3, satisfying (see Figure 1)

$$\Omega_1(t) \cap \Omega_2(t) \cap \Omega_3(t) = \{T(t)\}, \quad t \ge 0,$$
(3.1)

$$\Omega_1(t) \cap \Omega_2(t) = \Gamma_{ij} = \{ p_{ij}(s,t) : s \ge 0 \} \quad (i \ne j), \quad t \ge 0,$$
(3.2)

for some continuous parametrization $p_{ij}: [0,\infty)^2 \to \mathbb{R}^2$ with

$$p_{ij}(0,t) = T(t), \quad t \ge 0, p_{ij}(s,t) \ne p_{ij}(s',t), \quad \text{if } s \ne s'.$$
(3.3)

We assume that near the triple junction T(t) the domains Ω_i satisfy the local epigraph property: for each $i \in \{1, 2, 3\}$ and $t_0 \geq 0$ there exists a neighborhood $\mathcal{O} \times (t_0 - \delta, t_0 + \delta)$ of $(T(t_0), t_0)$ such that, for $|t - t_0| < \delta$ and in an appropriate coordinate system,

$$T(t) = (s(t), d(t))$$
 (3.4)

and there exist continuous functions u(x,t), w(x,t) satisfying

$$\Gamma_{ij} \cap \mathcal{O} = \{(x, y) \in \mathcal{O} : y = u(x, t), x \leq s(t)\},$$

$$\Gamma_{ik} \cap \mathcal{O} = \{(x, y) \in \mathcal{O} : y = w(x, t), x \geq s(t)\} \quad (j \neq i \neq k), \qquad (3.5)$$

$$\Omega_i \cap \mathcal{O} = \{(x, y) \in \mathcal{O} : y \geq G(x, t)\}$$

where

$$G(x,t) = \begin{cases} u(x,t), & x \le s(t), \\ w(x,t), & x \ge s(t). \end{cases}$$
(3.6)

T

As for regularity, we only assume that the domains $\Omega_i(\cdot)$'s are continuous in the Hausdorff metric, i.e.

$$\lim_{\tau \to t} d_H(\Omega_i(t), \Omega_i(\tau)) = 0, \quad i = 1, 2, 3, \quad t \ge 0,$$
(3.7)

where, for two closed sets $A, B \subset \mathbb{R}^2$,

$$d_{H}(A,B) = \max\left\{\sup_{x \in A} \inf_{y \in B} |x - y|, \sup_{y \in B} \inf_{x \in A} |x - y|\right\}.$$
 (3.8)

If $p_{ij}(\cdot,t)$ is differentiable at s = 0 and $|p_{ijs}(0,t)| \neq 0$ then the angles $\theta_i(t)$ between the curves Γ_{ij} are well defined (see Figure 1). Our goal is to obtain a weak formulation of the angle conditions

$$\theta_i(t) = \theta_i^*, \quad \forall t \ge 0. \tag{3.9}$$

For this, assume that there exist smooth test functions $\phi(x,t)$ and $\psi(x,t)$ such that $(s(t_0), t_0)$ is the strict minimizer of G - F where s(t) is defined by (3.4), G is the defining function for Ω_i in the local epigraph condition (3.5) and

$$F(x,t) = \begin{cases} \phi(x,t), & x \le s(t), \\ \psi(x,t), & x \ge s(t), \end{cases}$$
(3.10)

(see Figure 7).

Figure 7: The functions ϕ and ψ are test functions for the definition of Angle $(u(\cdot,t), w(\cdot,t)) \ge \theta^*$ weakly.

Since ϕ and ψ are smooth, the angle between their graphs at time t, $Angle(\phi(\cdot, t), \psi(\cdot, t))$, is well-defined. If $\theta_i(t_0)$ were defined and satisfied $\theta_i(t_0) \geq \theta_i^*$, then we would have

$$Angle(\phi(\cdot,t),\psi(\cdot,t)) \geq \theta_i(t_0) \geq \theta_i^*.$$

Hence we expect to have further restrictions on ϕ and ψ only when

$$Angle(\phi(\cdot, t), \psi(\cdot, t)) < \theta_i^*.$$
(3.11)

Suppose then that (3.11) holds. Let $\{\Omega_i^{\epsilon} : t \ge 0, i = 1, 2, 3\}$ denote the solution to the curvature perturbed problem (1.4), (1.5). We assume that Ω_i^{ϵ} satisfy (3.1)-(3.3) and denote the corresponding triple junction by $T^{\epsilon}(t) = (s^{\epsilon}(t), d^{\epsilon}(t))$. We also assume that $\Omega_i^{\epsilon}(t)$ has a form similar to that of $\Omega_i(t)$, i.e.

$$\Gamma_{ij}^{\epsilon} \cap \mathcal{O} = \{(x, y) \in \mathcal{O} : y = u^{\epsilon}(x, t), x \leq s^{\epsilon}(t)\},$$

$$\Gamma_{ik}^{\epsilon} \cap \mathcal{O} = \{(x, y) \in \mathcal{O} : y = w^{\epsilon}(x, t), x \geq s^{\epsilon}(t)\},$$
(3.12)

for smooth functions u^{ϵ} , w^{ϵ} satisfying

$$u_{t}^{\epsilon} = \epsilon \alpha_{ij} \frac{u_{xx}^{\epsilon}}{1 + (u_{x}^{\epsilon})^{2}} + v_{ji} \sqrt{1 + (u_{x}^{\epsilon})^{2}}, x < s^{\epsilon}(t)$$
$$w_{t}^{\epsilon} = \epsilon \alpha_{ik} \frac{w_{xx}^{\epsilon}}{1 + (w_{x}^{\epsilon})^{2}} + v_{ki} \sqrt{1 + (w_{x}^{\epsilon})^{2}}, x > s^{\epsilon}(t)$$
(3.13)

$$Angle(u^{\epsilon}(t), w^{\epsilon}(t)) = \theta_{i}^{*}$$

where

$$\alpha_{ij} = \mu_{ij} f_{ij}, \quad v_{ij} = \mu_{ij} (e_i - e_j).$$

Finally, we assume that

$$\lim_{\epsilon \to 0} \sup_{|t-t_0| < \delta} d_H(\Omega_i^{\epsilon}(t), \Omega_i(t)) = 0$$

$$\lim_{\epsilon \to 0} \inf_{|t-t_0| < \delta} |s^{\epsilon}(t) - s(t)| + |s^{\epsilon'}(t) - s'(t)| = 0.$$

 \mathbf{Set}

$$F^{\epsilon}(x,t) = \begin{cases} \phi(x+s(t)-s^{\epsilon}(t),t), & x \leq s^{\epsilon}(t), \\ \psi(x+s(t)-s^{\epsilon}(t),t), & x \geq s^{\epsilon}(t), \end{cases}$$
$$G^{\epsilon}(x,t) = \begin{cases} u^{\epsilon}(x,t), & x \leq s^{\epsilon}(t), \\ w^{\epsilon}(x,t), & x \geq s^{\epsilon}(t). \end{cases}$$

TT

Then, the local minimizers $(x^{\epsilon}, t^{\epsilon})$ of $G^{\epsilon} - F^{\epsilon}$ converge, as $\epsilon \to 0$, to $(s(t_0), t_0)$. If $x^{\epsilon} = s^{\epsilon}(t^{\epsilon})$, we would have

$$\theta^*_i = Angle(u^\epsilon(t^\epsilon), w^\epsilon(t^\epsilon)) \leq Angle(\phi(t^\epsilon), \psi(t^\epsilon)) < \theta^*_i.$$

Hence, $x^{\epsilon} \neq s^{\epsilon}(t^{\epsilon})$. From (3.13), we then obtain either

$$\phi_t + (s'(t^{\epsilon}) - s^{\epsilon'}(t^{\epsilon}))\phi_x \ge \epsilon \alpha_{ij} \frac{\phi_{xx}}{1 + (\phi_x)^2} + v_{ji}\sqrt{1 + (\phi_x)^2},$$

if $x^{\epsilon} < s^{\epsilon}(t^{\epsilon})$, or

$$\psi_t + (s'(t^{\epsilon}) - s^{\epsilon'}(t^{\epsilon}))\psi_x \ge \epsilon \alpha_{ik} \frac{\psi_{xx}}{1 + (\psi_x)^2} + v_{ki} \sqrt{1 + (\psi_x)^2} \,,$$

if $x^{\epsilon} > s^{\epsilon}(t^{\epsilon})$. Here, ϕ , ψ and their derivatives are evaluated at $(x^{\epsilon} + s(t^{\epsilon}) - s^{\epsilon}(t^{\epsilon}), t^{\epsilon})$. Now, let $\epsilon \to 0$ and use the assumption that $s^{\epsilon} \to s$ in C^1 , to conclude

$$\max\left\{\phi_t - v_{ji}\sqrt{1 + (\phi_x)^2}, \, \psi_t - v_{ki}\sqrt{1 + (\psi_x)^2}\right\} \ge 0 \quad \text{at } (s(t_0), t_0).$$

Since this inequality was derived under the assumption (3.11) we have

$$\max \left\{ Angle(\phi(\cdot, t), \psi(\cdot, t)) - \theta_i^*, \\ \phi_t - v_{ji}\sqrt{1 + (\phi_x)^2}, \ \psi_t - v_{ki}\sqrt{1 + (\psi_x)^2} \right\} \ge 0$$
(3.14)

at $(s(t_0), t_0)$. We are thus led to the following definition.

Definition 3.1 Let $\{\Omega_i(t) : t \ge 0, i = 1, 2, 3\}$ be unbounded, connected, closed regions satisfying (3.1)-(3.3), (3.7) and the local epigraph condition at the triple junction. We say that

$$\theta_i(t) \geq \theta_i^* \quad weakly \ \forall t > 0,$$

if for any smooth functions ϕ and ψ such that G - F(cf. (3.6), (3.10)) has a local minimum at $(s(t_0), t_0)$ for some t_0 , the inequality (3.14) holds at $(s(t_0), t_0)$.

Similarly we define the concept of $\theta_i(t) \leq \theta_i^*$ weakly.

Definition 3.2 Let $\{\Omega_i(t) : t \ge 0, i = 1, 2, 3\}$ be as in Definition 3.1. We say that

 $\theta_i(t) \leq \theta_i^* \quad weakly \ \forall t > 0$

if for any smooth functions ϕ and ψ , such that G - F (cf. (3.6), (3.10)) has a local maximum at $(s(t_0), t_0)$ for some t_0 , the inequality

$$\min \left\{ Angle(\phi(\cdot, t), \psi(\cdot, t)) - \theta_i^*, \\ \phi_t - v_{ji}\sqrt{1 + (\phi_x)^2}, \ \psi_t - v_{ki}\sqrt{1 + (\psi_x)^2} \right\} \le 0$$
(3.15)

holds at $(s(t_0), t_0)$.

Definition 3.3 We say that

$$\theta_i(t) = \theta_i^* \quad weakly \ \forall t > 0, \tag{3.16}$$

if $\theta_i(t) \leq \theta_i^*$ weakly for all t > 0 and $\theta_i(t) \geq \theta_i^*$ weakly for all t > 0.

Finally we give the definition of weak solution of

Normal Velocity of
$$\Gamma_{ij} = V_{ij} = v_{ij} = \mu_{ij}(e_i - e_j).$$
 (3.17)

subject to the angle conditions. In this formulation we interpret (3.17) as in the theory of viscosity solutions [6, 7], and the angle conditions as in Definition 3.3. Observe that if the parametrization p_{ij} is differentiable at some (s,t) with $|p_{ijs}(s,t)| \neq 0$, then (3.17) is equivalent to

$$p_{ij_t} \cdot n_{ij} = v_{ij} , \qquad (3.18)$$

Π

where $n_{ij}(s,t)$ denotes the unit vector normal to Γ_{ij} pointing into $\Omega_j(t)$.

Definition 3.4 Let the domains Ω_i be as in Definition 3.1. We shall say that $\{\Omega_i : i = 1, 2, 3\}$ is a weak (viscosity) solution of (3.17) and the angle conditions, if they satisfy (3.16) and they solve (3.17) in the sense of viscosity solutions (see [4, 8]).

In §4 we shall show that, there exists a unique viscosity solution in the sense of the above definition, see Theorem 4.3 and 4.4. In the case of Figure 3, it can be easily checked that the weak solution is the one depicted in Figure 3(b); in particular, the

Figure 8: (a) The solution proposed in [11] (see also Figure 3) and (b) comparison functions for the weak formulation. The $Angle(u(\cdot,t),w(\cdot,t)) > 120^{\circ}$ but these comparison functions show that $Angle(u(\cdot,t),w(\cdot,t))$ is not $\leq 120^{\circ}$ weakly.

solution depicted in Figure 3(a) is not a weak solution in the sense of Definition 3.4. To see this, let us rotate (90° clockwise) the graph in Figure 3(a), and define u(x,t) and w(x,t) as in Figure 8(a).

If α and β are as in (2.3), then $Angle(u(\cdot,t), w(\cdot,t)) > 120^{\circ}$. We want to show that this angle *is not* less than or equal to 120° weakly, i.e. that the condition (3.15) in Definition 3.2, with i = 2, j = 3, k = 1, $\theta_2^* = 120^{\circ}$, $v_{12} = -\alpha$ and $v_{32} = -\beta$, *is not* satisfied for some test functions ϕ and ψ . Let $\phi_1(x,t)$ and $\psi_1(x,t)$ denote the solution corresponding to Figure 3(b), that is the solution satisfying $Angle(\phi_1(\cdot,t),\psi_1(\cdot,t)) = 120^{\circ}$, see Figure 8(b). Then if $(0, -\alpha t)$ and $(0, -\sigma t)$ denote the positions of the triple junction in Figures 8(a) and (b) respectively, the functions

$$\phi(x,t) = \phi_1(x,t) + (\sigma - \alpha)t$$
 and $\psi(x,t) = \psi_1(x,t) + (\sigma - \alpha)t$

are admissible test functions for the proposed solution u(x,t), w(x,t). For these test functions we have, at x = 0,

$$\phi_t - v_{32}\sqrt{1 + (\phi_x)^2} = \phi_{1t} + \sigma - \alpha + \beta\sqrt{1 + (\phi_{1x})^2} = \sigma - \alpha > 0$$
(3.19)

 \mathbf{and}

$$\psi_t - v_{12}\sqrt{1 + (\psi_x)^2} = \psi_{1t} + \sigma - \alpha + \alpha\sqrt{1 + (\psi_{1x})^2} = \sigma - \alpha > 0.$$
 (3.20)

Thus, we can slightly change the slopes of ϕ and ψ and still maintain the above inequalities, while at the same time satisfying

$$Angle(\phi(\cdot, t), \psi(\cdot, t)) - 120^{\circ} > 0.$$
(3.21)

Since inequalities (3.19)-(3.21) contradict (3.15) we conclude that this is not a viscosity solution.

Finally, we want to show that the solution in Figure 3(b) is a viscosity solution.

Figure 9: The VST solution *does* satisfy the weak formulation: $\theta(t) \ge 120^{\circ}$ weakly.

Since strongly $Angle(\Gamma_{12}(t), \Gamma_{23}(t)) = 120^{\circ}$, $Angle(\Gamma_{12}(t), \Gamma_{13}(t)) < 120^{\circ}$, and $Angle(\Gamma_{23}(t), \Gamma_{13}(t)) > 120^{\circ}$, we need only verify that

 $Angle(\Gamma_{12}(t),\Gamma_{13}(t)) \geq 120^{o} \quad \text{and} \quad Angle(\Gamma_{23}(t),\Gamma_{13}(t)) \leq 120^{o} \quad weakly.$

We shall only show that the first inequality holds (in the weak sense) since the second inequality can be checked in an analogous manner. For this, consider smooth test functions $\phi(x,t)$ (x < s(t)) and $\psi(x,t)$ (x > s(t)) such that G - F (cf. (3.6), (3.10)) has a local minimum at $(s(t_0), t_0)$ for some $t_0 > 0$ (see Figure 9). We want to show that the inequality (3.14) holds at $(s(t_0), t_0)$ with $i = 1, j = 2, k = 3, v_{31} = 0$ and $\theta_1^* = 120^\circ$. If we assume that

$$Angle(\phi(\cdot, t_0), \psi(\cdot, t_0)) < 120^{\circ}$$

then we need to show that

$$\max\left\{\phi_t - v_{21}\sqrt{1 + (\phi_x)^2}, \psi_t\right\} \ge 0 \quad \text{at } (s(t_0), t_0). \tag{3.22}$$

T

Now, since $w(s(t), t) - \psi(s(t), t)$ has a local minimum at $t = t_0$, we have

$$0 = \frac{d}{dt} \left(w(s(t), t) - \psi(s(t), t) \right) \Big|_{t=t_0} = -\psi_t(s(t_0), t_0) - s'(t_0)\psi_x(s(t_0), t_0)$$

that is,

$$\psi_t(s(t_0), t_0) = -s'(t_0)\psi_x(s(t_0), t_0).$$

But since $0 = \psi(s(t_0), t_0) \ge \psi(x, t_0)$, it follows that $\psi_x(s(t_0), t_0) \le 0$, so that (since s'(t) > 0)

$$\psi_t(s(t_0),t_0)\geq 0$$

and (3.22) holds.

3.2 An equivalent formulation

For solutions that are differentiable at the triple junction, like the ones discussed at the end of §3.1, the definition of weak solution can be formulated in terms of *algebraic* conditions. In order to derive these conditions, consider a situation like the one in Figure 7 where (with i = 1, j = 2, k = 3)

$$\Gamma_{12}(t) \cap \mathcal{O} = \{ (x, y) \in \mathcal{O} : y = u(x, t), x \le s(t) \}$$

and

$$\Gamma_{13}(t) \cap \mathcal{O} = \{(x, y) \in \mathcal{O} : y = w(x, t), x \ge s(t)\}$$

(cf. (3.5)). Let $\phi(x,t)$ and $\psi(x,t)$ be smooth test functions and let F and G be defined by (3.10) and (3.6). Assume that G - F has a minimum at $(s(t_0), t_0)$ and that

$$Angle(\phi(\cdot, t_0), \psi(\cdot, t_0)) \equiv \theta_1 < \theta_1^*.$$

Then, using (3.14), we have that $Angle(u(\cdot,t),w(\cdot,t)) \ge \theta_1^*$ weakly at t_0 if and only if

$$\max\left\{\phi_t - v_{21}\sqrt{1 + (\phi_x)^2}, \, \psi_t - v_{31}\sqrt{1 + (\psi_x)^2}\right\} \ge 0 \quad \text{at } (s(t_0), t_0). \tag{3.23}$$

Let T(t) = (s(t), d(t)) denote the position of the triple junction and n_u, n_w, n_{ϕ} and n_{ψ} the unit normal vectors to the graphs of u, w, ϕ and ψ at $(s(t_0), d(t_0))$ (pointing towards Ω_1). Since (u, w) is a differentiable solution, we have

$$T_t(t_0) \cdot n_u = v_{21}$$
 and $T_t(t_0) \cdot n_w = v_{31}$.

Thus,

$$T_t(t_0) = An_u + Bn_w \tag{3.24}$$

where

$$A = \frac{(v_{21} - v_{31}n_u \cdot n_w)}{1 - (n_u \cdot n_w)^2} = \frac{(v_{21} + v_{31}\cos(\theta_1))}{\sin^2(\theta_1)}$$
$$B = \frac{(-v_{21}n_u \cdot n_w + v_{31})}{1 - (n_u \cdot n_w)^2} = \frac{(v_{21}\cos(\theta_1) + v_{31})}{\sin^2(\theta_1)}.$$
(3.25)

On the other hand, since $\phi(s(t), t) = d(t)$,

$$T_t \cdot n_{\phi} = (s', d') \cdot \frac{(-\phi_x, 1)}{\sqrt{1 + (\phi_x)^2}} = \frac{-s'\phi_x + d'}{\sqrt{1 + (\phi_x)^2}} = \frac{\phi_t}{\sqrt{1 + (\phi_x)^2}}$$

and similarly for $T_t \cdot n_{\psi}$, so that (3.23) is equivalent to

 $\max \{T_t \cdot n_{\phi} - v_{21}, T_t \cdot n_{\psi} - v_{31}\} \ge 0 \quad \text{at } (s(t_0), t_0).$

Then, using (3.24), the above condition becomes

 $\max \{An_{u} \cdot n_{\phi} + Bn_{w} \cdot n_{\phi} - v_{21}, An_{u} \cdot n_{\psi} + Bn_{w} \cdot n_{\psi} - v_{31}\} \ge 0, \quad (3.26)$ at $(s(t_{0}), t_{0})$. Now, let

$$\begin{array}{lll} Angle(u(\cdot,t_0),\psi(\cdot,t_0)) &\equiv & \eta \ge \theta_1 \\ Angle(w(\cdot,t_0),\phi(\cdot,t_0)) &\equiv & \alpha \ge \theta_1 \end{array}$$

so that

$$n_{u} \cdot n_{\phi} = \cos(\alpha - \theta_{1})$$

$$n_{w} \cdot n_{\phi} = -\cos(\alpha)$$

$$n_{u} \cdot n_{\psi} = -\cos(\eta)$$

$$n_{w} \cdot n_{\psi} = \cos(\eta - \theta_{1}).$$

Then, substituting (3.25) and (3.27) into (3.26) we obtain

$$\max\left\{\left(\sin(\alpha)\sin(\theta_1) - \sin^2(\theta_1)\right)v_{21} + \left(-\cos(\alpha)\sin^2(\theta_1) + \sin(\alpha)\cos(\theta_1)\sin(\theta_1)\right)v_{31}, \\ \left(-\cos(\eta)\sin^2(\theta_1) + \sin(\eta)\cos(\theta_1)\sin(\theta_1)\right)v_{21} + \left(\sin(\eta)\sin(\theta_1) - \sin^2(\theta_1)\right)v_{31}\right\} \ge 0$$

T

۵

1

or, equivalently,

$$\max\left\{\frac{1}{\sin(\theta_{1})}v_{21} + \frac{\cos(\theta_{1})}{\sin(\theta_{1})}v_{31} - \frac{1}{\sin(\alpha)}v_{21} - \frac{\cos(\alpha)}{\sin(\alpha)}v_{31}, \frac{1}{\sin(\theta_{1})}v_{31} + \frac{\cos(\theta_{1})}{\sin(\theta_{1})}v_{21} - \frac{1}{\sin(\eta)}v_{31} - \frac{\cos(\eta)}{\sin(\eta)}v_{21}\right\} \ge 0.$$
(3.27)

Thus, since $Angle(\phi(\cdot, t), \psi(\cdot, t)) = \alpha + \eta - \theta_1$, the condition (3.23) holds if and only if (3.27) is satisfied for all α and η with

$$heta_1 \leq lpha \ , \ \eta < \pi \quad ext{and} \quad lpha + \eta - heta_1 < heta_1^*.$$

Analogous equivalent conditions can be derived for the definition of $\theta_1 \leq \theta_1^*$ weakly as well as for θ_2 and θ_3 . Indeed, if we let

$$H_{ijk}(\lambda) = \frac{1}{\sin(\lambda)} v_{ji} + \frac{\cos(\lambda)}{\sin(\lambda)} v_{ki} \quad (i \neq j \neq k = 1, 2, 3)$$

then we have the following theorem.

Theorem 3.5 Assume the domains Ω_i are differentiable at the triple junction. Then $\theta_i(t) \geq \theta_i^*$ (resp. $\theta_i(t) \leq \theta_i^*$) weakly if and only if either

$$\theta_i(t) \ge \theta_i^* \quad (resp.\theta_i(t) \le \theta_i^*) \quad strongly,$$

or

$$\max (resp. \min) \{H_{i,i+1,i+2}(\theta_i(t)) - H_{i,i+1,i+2}(\alpha), \\ H_{i,i+2,i+1}(\theta_i(t)) - H_{i,i+2,i+1}(\eta)\} \ge 0 \quad (resp. \le 0)$$
(3.28)

for all α and η satisfying

$$\theta_{i}(t) \leq \alpha, \eta < \pi \quad and \quad \alpha + \eta - \theta_{i}(t) < \theta_{i}^{*}$$

$$(resp. \ 0 < \alpha, \eta \leq \theta_{i}(t) \quad and \quad \alpha + \eta - \theta_{i}(t) > \theta_{i}^{*}).$$

$$(3.29)$$

Here the indices i + 1 and i + 2 should be interpreted modulo 3.

3.3 A catalog of weak solutions

In the previous sections we have presented the general definition of weak solutions as well as the equivalent formulation for solutions which are differentiable at the triple junction. We can now attempt to catalog them according to the interface velocities and initial conditions. Since the behavior at the triple junction is a local property we shall restrict our attention to solutions with linear initial data. Furthermore, for simplicity we will assume that the smallest (in magnitude) interface velocity is equal to zero and that $\theta_i^* = 120^\circ$. Under these assumptions, the five possible solutions for *compatible* initial data are presented in Figure 10. We note here that the only solutions in Figure 10(a) do not always coincide with the ones derived in [12].

Finally, Figure 11 shows solutions for *incompatible* initial data. Notice here that if the initial angles are not too far from 120° then the solution will produce a corner in order to keep an angle at 120° for t > 0 (Figure 11(a)). On the other hand, if the initial angles differ from 120° by more than a critical amount (which can be explicitly calculated) then it becomes impossible to keep the angle condition in the strong sense (Figure 11(b)).

4 Existence and uniqueness of weak solutions

In this section we prove a local existence and uniqueness result for differentiable solutions of the system of Hamilton-Jacobi equations (3.18) subject to the weak angle conditions (3.16).

4.1 Existence

For the sake of clarity we shall restrict ourselves to the case in which

$$v_{21} = \alpha \ge v_{23} = \beta > v_{13} = 0$$

with

$$\alpha \leq 2\beta$$

and uniform angle conditions

$$\theta_i(t) = \theta_i^* = \frac{2\pi}{3}, \quad i = 1, 2, 3 \quad t > 0.$$
 (4.1)

T

In this case the weak angle conditions (3.28) can be simplified, as we show in the following lemma.

Lemma 4.1 Suppose that $\theta_2(t) < \pi$. Then the angle conditions (4.1) hold weakly if, and only if,

$$\theta_2(t) \in \left\{\frac{2\pi}{3}\right\} \cup [\theta_0, \pi) \tag{4.2}$$

where $\theta_0 \in [\frac{2\pi}{3}, \pi)$ satisfies

$$\cos(\theta_0) = -\frac{(4\alpha\beta - \alpha^2 - \beta^2)}{2(\alpha^2 + \beta^2 - \alpha\beta)}.$$
(4.3)

Proof. First we show that $\theta_1(t) \ge \frac{2\pi}{3}$ weakly regardless of the value of $\theta_1(t)$. In view of Theorem 3.5, we need to show that, whenever $\theta_1(t) < \frac{2\pi}{3}$, we have

$$\max\left\{H_{1,2,3}(\theta_1(t)) - H_{1,2,3}(\xi), H_{1,3,2}(\theta_1(t)) - H_{1,3,2}(\eta)\right\} \ge 0 \tag{4.4}$$

for all ξ and η satisfying

$$\theta_1(t) \leq \xi, \eta < \pi \text{ and } \xi + \eta - \theta_1(t) < \theta_1^*.$$

Here,

$$H_{1,2,3}(\xi) = \frac{\alpha}{\sin(\xi)}$$
 and $H_{1,3,2}(\eta) = \frac{\alpha \cos(\eta)}{\sin(\eta)}$.

Then, since $H'_{1,3,2}(\eta) < 0$, we have

$$H_{1,3,2}(\theta_1(t_0)) - H_{1,3,2}(\eta) \ge 0$$

for $\eta \in [\theta_1(t_0), \frac{2\pi}{3}]$ and (4.4) holds. The opposite inequality, $\theta_1(t) \leq \frac{2\pi}{3}$, as well as the weak equality $\theta_3(t) = \frac{2\pi}{3}$ can be proved in a similar manner. Thus, in this case, the weak angle conditions impose no restrictions on $\theta_1(t)$ and $\theta_3(t)$.

Finally, we want to show that $\theta_2(t) = \frac{2\pi}{3}$ weakly if and only if (4.2) holds. Since

$$H_{2,1,3}(\xi) = -\frac{(\alpha + \beta \cos(\xi))}{\sin(\xi)} \text{ and } H_{2,3,1}(\eta) = -\frac{(\alpha \cos(\eta) + \beta)}{\sin(\eta)}$$

we have $H'_{2,1,3}(\xi) > 0$ for $\xi < \arccos(-\beta/\alpha)$ and $H'_{2,3,1}(\eta) > 0$ for all η . It follows that $\theta_2(t) \geq \frac{2\pi}{3}$ weakly if and only if $\theta_2(t) \geq \frac{2\pi}{3}$ strongly. On the other hand,

 $\theta_2(t) \leq \frac{2\pi}{3}$ weakly if and only if $\theta_2(t) \leq \frac{2\pi}{3}$ strongly or $H_{2,1,3}(\theta_2(t)) \leq H_{2,1,3}(\xi)$ for all ξ with $\frac{2\pi}{3} \leq \xi \leq \theta_2(t)$. In the latter case, since $H_{2,1,3}(\xi)$ has a maximum at $\xi = \arccos(-\beta/\alpha) > \frac{2\pi}{3}$, we must have

$$\pi > \theta_2(t) \ge \theta_0$$

where $\theta_0 > \arccos(-\beta/\alpha)$ satisfies

$$H_{2,1,3}(\theta_0) = H_{2,1,3}(\frac{2\pi}{3})$$

or , equivalently, θ_0 is given by (4.3).

Remark 4.2 Notice that the condition (4.2) is in agreement with the construction of weak solutions for incompatible initial data shown in Figure 11. Indeed, Figure 11(a) corresponds to solutions for which $\theta_2(t) = \frac{2\pi}{3}$ while the solution in Figure 11(b) satisfies $\theta_2(t) \in [\theta_0, \pi)$.

With the help of Lemma 4.1 we will now establish the existence of weak solutions for smooth initial data. First notice that if a solution satisfies

$$\theta_2(t) = \frac{2\pi}{3} \text{ strongly}$$
(4.5)

then

$$\theta_1(t) = \frac{\pi}{2} + \overline{\eta}, \quad \theta_3(t) = \frac{\pi}{2} + \widehat{\eta}$$
(4.6)

where $\overline{\eta}, \, \widehat{\eta} \in [0, \frac{\pi}{3}]$ are given by

$$\cos(\overline{\eta}) = \frac{\alpha}{\sigma}, \quad \cos(\widehat{\eta}) = \frac{\beta}{\sigma}, \quad \sigma = \frac{2}{\sqrt{3}}\sqrt{\alpha^2 + \beta^2 - \alpha\beta}.$$

Moreover, the triple junction satisfies

$$\left|\frac{d}{dt}T(t)\right| = \sigma. \tag{4.7}$$

Π

We consider initial data of the following form

.

.

$$\begin{array}{rcl} T(0) &=& (0,0)\,, \\ \Gamma_{13} &=& \{(x,w_0(x))\,:\, x\geq 0\}\,, \\ \Gamma_{12} &=& \{(u_0(y),y)\,:\, y\geq 0\}\,, \\ \Gamma_{23} &=& \{(v_0(y),y)\,:\, y\leq 0\}\,, \end{array}$$

where u_0, v_0, w_0 are smooth functions satisfying

$$\begin{array}{ll} (i) & u_0(0) = w_0(0) = v_0(0) \,, \\ (ii) & u_0, w_0, v_0 \in C^2(\mathbb{R}) \text{ and their limits as } |x| \to \infty \text{ exist}, \\ (iii) & w_0'(0) = 0 \\ (iv) & u_0'(0) < \tan(\overline{\eta}) \,, \\ (v) & v_0'(0) > -\tan(\widehat{\eta}). \end{array}$$

$$(4.8)$$

We look for a solution of the form

$$\begin{array}{rcl} T(t) &=& (s(t), w_0(s(t))) \,, \\ \Gamma_{13} &=& \{(x, w_0(x)) \,:\, x \geq s(t)\} \,\,, \\ \Gamma_{12} &=& \{(u(y, t), y) \,:\, y \geq w_0(s(t))\} \,\,, \\ \Gamma_{23} &=& \{(v(y, t), y) \,:\, y \leq w_0(s(t))\} \end{array}$$

satisfying (4.5) and (4.6). Here s(t) is an increasing function that, in view of (4.7), solves

$$s'(t) = \frac{\sigma}{\sqrt{1 + (w'_0(s(t)))^2}}$$
(4.9)

while the functions u and v solve (in the viscosity sense)

$$u_{t} = \alpha \sqrt{1 + u_{y}^{2}}, \quad y > w_{0}(s(t)), \quad t > 0,$$

$$v_{t} = \beta \sqrt{1 + v_{y}^{2}}, \quad y < w_{0}(s(t)), \quad t > 0, \quad (4.10)$$

$$u(w_0(s(t)),t) = v(w_0(s(t)),t) = s(t), t > 0.$$

Theorem 4.3 (Existence) Let u_0 , v_0 and w_0 satisfy (4.8) and let s(t) denote the unique solution of (4.9) with s(0) = 0. Then there exist T > 0 and functions u(y,t) and v(y,t) such that (u,v) is a viscosity solution of (4.10) for $t \in (0,T)$ with initial data (u_0,v_0) . Moreover, (u,v) is differentiable at $(w_0(s(t)),t)$ for all $t \in (0,T)$ and

$$u_{y}(w_{0}(s(t)), t) = -\tan(\overline{\eta} + \theta(t)),$$

$$v_{y}(w_{0}(s(t)), t) = \tan(\widehat{\eta} - \theta(t)),$$
(4.11)

where $|\theta(t)| < \frac{\pi}{2}$ satisfies $\tan(\theta(t)) = w'_0(s(t))$.

Proof. We shall construct the solution v(y,t) in the form of the value function for a deterministic control problem. A construction using the method of characteristics is also possible and we will indicate the connection between these two approaches below. The construction of u(y,t) is similar.

First we write the equation for v in (4.10) in the form

$$v_t = H(v_y) \tag{4.12}$$

where

$$H(p) = \beta \sqrt{1+p^2} = \sup_{|b| \le \beta} \left(-bp + \sqrt{\beta^2 - b^2} \right).$$

The above representation for H suggests that (4.12) is related to the following control problem [7]: fix (y,t) with $y \leq w_0(s(t)), t > 0$. Let $\mathcal{A}(y,t)$ denote the collection of all pairs $(\xi(\cdot), \tau)$ satisfying

$$\begin{array}{rcl} \xi & : & [0,t] \rightarrow {\rm I\!R} \,, & \tau \in [0,t] \,, \\ \xi(r) & \leq & w_0(s(r)) \,, & \forall r \in [\tau,t] \,, \\ \xi(\tau) & = & w_0(s(\tau)) \,, {\rm if} \, \tau > 0 \,, \\ \xi(t) & = & y \\ |\xi'(r)| & \leq & \beta \,, \, \forall r \in [0,t] \,. \end{array}$$

For $(\xi(\cdot), \tau) \in \mathcal{A}(y, t)$ set

$$J(y,t;\xi(\cdot),\tau) = \int_{\tau}^{t} \sqrt{\beta^2 - (\xi'(s))^2} \, ds + s(\tau)\chi_{\tau>0} + v_0(\xi(0))\chi_{\tau=0}$$

1

and

$$v(y,t) = \sup_{(\xi(\cdot),\tau)\in\mathcal{A}(y,t)} J(y,t;\xi(\cdot),\tau)$$

÷

Then v(y,t) is the viscosity solution of (4.12), see e.g. [7]. Furthermore, the maximizers in the definition of v above are straight lines. Hence,

$$v(y,t) = \max_{b \in \mathbf{R}} \left\{ (t-\tau) \sqrt{\beta^2 - b^2} + v_0 (y-bt) \chi_{\tau=0} + s(\tau) \chi_{\tau>0} \right\}$$

where for any $b \in \mathbb{R}$, $\tau \in [0, t]$ is defined by

$$\tau = \inf \{ \rho \in [0, t] : y + b(r - t) \le w_0(s(r)), \forall r \in [\rho, t] \}.$$

Now, observe that whenever $v'_0(0) \leq 0$, the assumption

$$v_0'(0) > -\tan(\widehat{\eta})$$

yields

$$\frac{\beta}{\sigma} < \frac{1}{\sqrt{1 + \left(v_0'(0)\right)^2}}.$$

Then an elementary analysis using $s'(0) = \sigma$ and the above fact implies that v(w(s(t)), t) = s(t) for all sufficiently small $t \ge 0$. Indeed, there exist T > 0, $\delta(t) > 0$ such that for $t \in (0, T]$ and $y \in (w_0(s(t)) - \delta(t), w_0(s(t))]$ we have

$$v(t,y) = V(\tau,t) \equiv s(\tau) + \beta(t-\tau)\cos(\theta(\tau) - \hat{\eta})$$
(4.13)

where $\theta(\cdot)$, $\hat{\eta}$ are as the statement of the Theorem and $\tau = \tau(y, t)$ is the solution of

$$y = w_0(s(\tau)) + \beta(t-\tau)\sin(\theta(\tau) - \hat{\eta}). \tag{4.14}$$

Thus, differentiating (4.13) and (4.14) with respect to y and evaluating at $(w_0(s(t)), t)$ we obtain

$$v_{y} = (s'(t) - \beta \cos(\theta(t) - \hat{\eta})) \tau_{y}$$

and

$$1 = \left(w_0'(s(t))s'(t) - \beta\sin(\theta(t) - \widehat{\eta})\right)\tau_y.$$

Since

$$s'(t) = \sigma \cos(\theta(t)), \quad w'_0(s(t))s'(t) = \sigma \sin(\theta(t))$$

we have

$$v_{y} = \frac{\sigma \cos(\theta(t)) - \beta \cos(\theta(t) - \hat{\eta})}{\sigma \sin(\theta(t)) - \beta \sin(\theta(t) - \hat{\eta})}$$

and since $\cos(\hat{\eta}) = \sigma/\beta$,

$$v_{y} = \frac{\cos(\theta(t)) - \cos(\hat{\eta})\cos(\theta(t) - \hat{\eta})}{\sin(\theta(t)) - \cos(\hat{\eta})\sin(\theta(t) - \hat{\eta})}$$
$$= \frac{\cos(\hat{\eta} + (\theta(t) - \hat{\eta})) - \cos(\hat{\eta})\cos(\theta(t) - \hat{\eta})}{\sin(\hat{\eta} + (\theta(t) - \hat{\eta})) - \cos(\hat{\eta})\sin(\theta(t) - \hat{\eta})}$$
$$= \tan(\hat{\eta} - \theta(t))$$
(4.15)

ž

T

and the proof of the Theorem is complete.

Finally let us mention that one could also use the method of characteristics to derive (4.13). Indeed,

$$Y(t;\tau) = w_0(s(\tau)) + \beta(t-\tau)\sin(\theta(\tau) - \hat{\eta})$$

is the characteristic curve for (4.12) emanating from the boundary point $(w_0(s(\tau)), \tau)$. Characteristics emanating from the initial data are

$$\overline{Y}(t;\hat{y}) = \hat{y} + \beta t \frac{v_0'(\hat{y})}{\sqrt{1 + (v_0'(\hat{y}))^2}}$$

and, for $t \leq t_0(\hat{y})$,

$$v(\overline{Y}(t;\hat{y}),t) = \overline{V}(\hat{y},t) \equiv v_0(\hat{y}) + \beta t \frac{1}{\sqrt{1 + (v_0'(\hat{y}))^2}}.$$

Now assume that $Y(t;\tau) = \overline{Y}(t;\hat{y})$ for some $0 \le \tau \le t$ and $\hat{y} \le 0$. Then, for sufficiently small t > 0, we have $V(t,\tau) > \overline{V}(\hat{y},t)$ and (4.13) follows.

4.2 Uniqueness

In this section we prove that the solution $(u(y,t), v(y,t), w_0(x))$ constructed in the previous subsection is the *unique* viscosity solution of (3.16) and (3.18). Suppose that there is another solution $\{\hat{\Gamma}_{ij} : t \in [0,T]\}$ of (3.16) and (3.18), satisfying (3.1), (3.2), (3.7). We further assume that there is a neighborhood \mathcal{N} of $\{(t,0) : t \in (0,T]\}$ such that for $i \neq j \in \{1,2,3\}$

$$\hat{p}_{ij} \in C^1(\mathcal{N}) \quad |\hat{p}_{ij,s}| > 0 \text{ on } \mathcal{N} > 0,$$
(4.16)

where \hat{p}_{ij} is a parametrization of the $\hat{\Gamma}_{ij}$'s. Let $\hat{T}(t)$ be the triple point of $\hat{\Gamma}$. Since \hat{p}_{ij} is differentiable on \mathcal{N} , for $t \in (0,T]$

$$\hat{T}'(t) \cdot n_{13}(0,t) = 0, \quad \hat{T}'(t) \cdot n_{21}(0,t) = \alpha, \quad \hat{T}'(t) \cdot n_{23}(0,t) = \beta.$$
 (4.17)

Moreover the angles between the arcs $\hat{\Gamma}_{ij}$ are well defined and by (4.1) we have,

if
$$\theta_2(t) \neq \frac{2\pi}{3}$$
, then $\theta_2(t) \ge \theta_0$, (4.18)

where $\theta_0 \in (\frac{2\pi}{3}, \pi)$ is as in (4.3). Since $\theta_2 \geq \frac{2\pi}{3}$, and $\hat{T}'(t) \cdot n_{13}(0, t) = 0$, it is elementary to show that

$$\hat{V}(t) := \hat{T}'(t) \cdot \frac{p_{13,s}(0,t)}{|p_{13,s}(0,t)|} \ge \sigma > \alpha, \tag{4.19}$$

where σ is as in (4.7).

Theorem 4.4 (Uniqueness) $\hat{\Gamma}_{ij}(t) = \Gamma_{ij}(t)$ for all $t \leq T$ and i, j = 1, 2, 3.

Proof. First we will show that

$$\hat{\Gamma}_{13}(t) \subset \Gamma_{13}(t), \quad \forall t \in [0, T].$$
(4.20)

Fix $t_0 > 0$. In view of (4.16), there is $\delta = \delta(t_0) > 0$ such that, the arc

$$S_{\delta}(t) = \hat{\Gamma}_{13}(t) \cap \{(x, y) : |(x, y) - \hat{T}(t_0)| < \delta\},\$$

is continuously differentiable for all $t_0 \leq t \leq t_0 + \delta$. Suppose that $x \in S_{\delta}(t)$ for some $t_0 \leq t \leq t_0 + \delta$ and $x \neq \hat{T}(t)$. Then $x \neq \hat{T}(s)$ for all s near t. Set

$$s(x,t) = \begin{cases} t_0, & \text{if } x \neq \hat{T}(s), \forall s \in [t_0,t], \\ \sup\{s < t : x = \hat{T}(s)\}, & \text{otherwise.} \end{cases}$$

Since the normal velocity at x is zero, we conclude that $x \in \hat{\Gamma}_{13}(s(x,t))$. Hence for all $t_0 \leq t \leq t_0 + \delta$ we have,

$$S_{\delta}(t) \subset S_{\delta}(t_0) \cup \{T(s) : t_0 \leq s \leq t_0 + \delta\}.$$

If for some $t_0 \leq t \leq t_0 + \delta$, $\hat{T}(t) \notin \hat{\Gamma}_{13}(t_0)$ then,

$$\frac{p_{13,s}(0,t)}{|p_{13,s}(0,t)|} = -\frac{\hat{T}'}{|\hat{T}'|},$$

which contradicts (4.19). Hence for all $t \in [t_0, t_0 + \delta]$,

 $T(t)\in \widehat{\Gamma}_{13}(t_0),$

and consequently,

$$T(t) = \hat{p}_{13}(s(t), t_0),$$

for some s(t).

Now extend $\hat{\Gamma}_{13}(t)$ in the following way,

$$\bar{\Gamma}_{13}(t) = \hat{\Gamma}_{13}(t) \cup \{p_{13}(s, t_0) : s \in [0, s(t)]\} \cup \{T(t_0) - \tau \hat{p}_{13,s}(s(t_0), t_0) : \tau < 0\}.$$

Then for $t \in [t_0, t_0 + \delta]$, $\overline{\Gamma}_{13}$ solves a two-phase geometric problem with normal velocity zero. By the uniqueness result for the two-phase problem (see e.g. [4]), we conclude that $\overline{\Gamma}_{13}(t) = \overline{\Gamma}_{13}(t_0)$, for all $t \in [t_0, t_0 + \delta]$. (4.20) follows from the continuity of $\widehat{\Gamma}$.

By (4.20) we conclude that,

$$\hat{T}(t) = (x(t), w_0(x(t))).$$

We claim that x(t) is equal to the unique solution of (4.9) and that $\theta_2(t) = \frac{2\pi}{3}$ for all t > 0. Indeed if $\theta_2(t) \neq \frac{2\pi}{3}$ for some t > 0. By (4.18) $\theta_2(t) \neq \frac{2\pi}{3}$ for all t > 0. Moreover by (4.17) we conclude that $\theta_1 < \frac{\pi}{2}$.

TT

 \mathbf{Set}

$$d(t) = \sup\{y : (x, y) \in \overline{\Gamma}_{12}(t) \text{ for some } x\}.$$

Since $\theta_1 < \pi/2$, and $\hat{\Gamma}$ is continuous in the Hausdorff topology, we conclude that the maximizer in the above expression is attained. Since $\hat{\Gamma}_{12}$ is viscosity solution of the equation $V = \alpha$, we conclude that

$$\frac{d}{dt}d(t) \le \alpha$$

in the sense of viscosity solutions. Therefore, $d(t) \leq \alpha t$. But this contradicts (4.19) and we conclude that $\theta_2(t) = \frac{2\pi}{3}$ for all t > 0. Then from (4.17) we obtain that x(t) = s(t). Hence

$$T(t) = \hat{T}(t), \quad \forall t \in [0, T].$$

Finally we define,

$$D = \{(x,y) : x \ge 0, y \ge w_0(x)\} \cup \{(x,y) : x \le 0, y \ge 0\}.$$

Then $\hat{\Gamma}_{12}$ and Γ_{12} both solve a two-phase geometric problem with normal velocity α in D satisfying the Dirichlet data

$$\hat{\Gamma}_{12} \cap \partial D = T(t), \quad \forall t \in [0, T].$$

Hence by standard comparison results on viscosity solutions for two-phase problems, [4], we conclude that the two solutions coincide.

A similar argument shows that $\hat{\Gamma}_{23} = \Gamma_{23}$.

5 A simple convergence result

In this section, we prove the convergence of solutions in the symmetric case: in the notation of §4, we take $\alpha = \beta$ and $w_0(x) \equiv 0$, $u_0(y) = v_0(-y)$. Without loss of generality we set $\alpha = 1$. Then the solutions to the ϵ -problem (2.5) have the form,

$$\begin{split} \Gamma_{13}^{\epsilon}(t) &= \{(x,0) \, : \, x \geq s^{\epsilon}(t)\}, \ t \geq 0\\ \Gamma_{21}^{\epsilon}(t) &= \{(u^{\epsilon}(y,t),y) \, : \, y \geq 0\}, \ t \geq 0\\ \Gamma_{23}^{\epsilon}(t) &= \{(u^{\epsilon}(-y,t),y) \, : \, y \leq 0\}, \ t \geq 0, \end{split}$$

where $s^{\epsilon}(t) = u^{\epsilon}(0, t)$ and u^{ϵ} solves,

$$u_t^{\epsilon} = \epsilon \frac{u_{yy}^{\epsilon}}{1 + (u_y^{\epsilon})^2} + \sqrt{1 + (u_y^{\epsilon})^2}, \qquad (5.1)$$

with Neumann data

$$u_{y}^{\epsilon}(0,t) = -\frac{1}{\sqrt{3}}.$$
(5.2)

We assume that there are constants $0 < k \leq \frac{1}{\sqrt{3}} \leq K$, satisfying

$$-K \leq u_{y}^{\epsilon}(y,0) \leq -k, \ \forall y \geq 0.$$

Then by the maximum principle we have

$$-K \le u_{y}^{\epsilon}(y,t) \le -k, \ \forall y,t \ge 0.$$

Hence for every $\epsilon > 0$ the solution exists for all time.

Theorem 5.1 (Convergence) As $\epsilon \to 0$, u^{ϵ} converges locally uniformly to the unique viscosity solution of

$$u_t = \sqrt{1 + (u_y)^2},\tag{5.3}$$

2

T

with Neumann boundary data (5.2). Moreover for all $t \ge 0$, $u_y(0,t)$ exists and satisfies (5.2) pointwise and

$$\lim_{\epsilon \to 0} s^{\epsilon}(t) = \frac{2}{\sqrt{3}}t.$$

Proof. Let $\Phi(y)$ be a smooth, convex function satisfying

$$\Phi(0) = 0, \quad \Phi_y(0) = -\frac{1}{\sqrt{3}}, \quad \Phi_y(y) = -k, \quad \forall y \ge Y(\Phi), \tag{5.4}$$

with some constant $Y(\Phi) > 0$ depending on Φ . For any Φ satisfying (5.4) define

$$\begin{split} \beta(\Phi) &= \sup\{\frac{\Phi_{yy}}{1+(\Phi_y)^2} + \sqrt{1+(\Phi_y)^2}\}\\ c^{\epsilon}(\Phi) &= \sup_{y\geq 0}\{-\epsilon\Phi(\frac{y}{\epsilon}) - ky\}. \end{split}$$

(Observe that the above maximum is achieved at $Y(\Phi)$ and that $c^{\epsilon}(\Phi)$ converges to zero as ϵ tends to zero.) Then for any fixed $t_0 \geq 0$, the function

$$\bar{u}^{\epsilon}(y,t) = s^{\epsilon}(t_0) + \epsilon \Phi(\frac{y}{\epsilon}) + \beta(\Phi)(t-t_0) + c^{\epsilon}(\Phi), \ t \ge t_0.$$

is a supersolution of (5.1) and (5.2). Since $u_y^{\epsilon}(y,t) \ge k$,

$$u^{\epsilon}(y,t_0) \leq ky - s^{\epsilon}(t_0) \leq \bar{u}^{\epsilon}(y,t_0), \quad \forall y \geq 0.$$

Hence by the maximum principle, $u^{\epsilon} \leq \bar{u}^{\epsilon}$ and, in particular, for any $h \geq 0$,

$$s^{\epsilon}(t_0+h) = u^{\epsilon}(0,t_0+h) \leq \overline{u}^{\epsilon}(0,t_0+h) = s^{\epsilon}(t_0) + \beta(\Phi)h + c^{\epsilon}(\Phi).$$

Therefore

$$s^{\epsilon}(t_0+h) - s^{\epsilon}(t_0) \le \beta(\Phi)h + c^{\epsilon}(\Phi), \quad \forall h \ge 0,$$
(5.5)

for any function Φ satisfying (5.4).

Similarly let $\hat{\Phi}$ be a smooth, concave function satisfying (5.4) with k replaced by K, i.e., $\hat{\Phi}$ satisfies

$$\hat{\Phi}(0) = 0, \quad \hat{\Phi}_{y}(0) = -\frac{1}{\sqrt{3}}, \quad \hat{\Phi}_{y}(y) = -K, \quad \forall y \ge Y(\hat{\Phi}),$$
 (5.6)

and set

$$\begin{split} \hat{\beta}(\hat{\Phi}) &= \inf\{\frac{\hat{\Phi}_{yy}}{1+(\hat{\Phi}_y)^2} + \sqrt{1+(\hat{\Phi}_y)^2}\}\\ \hat{c}^{\epsilon}(\hat{\Phi}) &= \sup_{y\geq 0}\{\epsilon\hat{\Phi}(\frac{y}{\epsilon}) + Ky\}. \end{split}$$

Then, arguing as above,

$$s^{\epsilon}(t_0+h) - s^{\epsilon}(t_0) \ge \hat{\beta}(\hat{\Phi})h - \hat{c}^{\epsilon}(\hat{\Phi}), \quad \forall h \ge 0.$$
(5.7)

Now let,

$$s^*(t) = \limsup_{\epsilon \to 0, r \to t} s^{\epsilon}(r), \qquad s_*(t) = \liminf_{\epsilon \to 0, r \to t} s^{\epsilon}(r).$$

Since $c^{\epsilon}(\Phi)$ converges to zero as $\epsilon \to 0$, (5.5) implies that for any $t, h \ge 0$ we have,

 $s^*(t+h) - s_*(t) \le \beta(\Phi)h,$

for all Φ satisfying (5.4), and similarly by (5.7),

$$s_*(t+h) - s^*(t) \ge \hat{\beta}(\hat{\Phi})h,$$

for all $\hat{\Phi}$ satisfying (5.6). It is easy to show that the infimum of $\beta(\Phi)$ over all Φ satisfying (5.4) is equal to $2/\sqrt{3}$. Also the supremum of $\hat{\beta}(\hat{\Phi})$ over all $\hat{\Phi}$ is equal to $2/\sqrt{3}$. Hence we have

$$\frac{2}{\sqrt{3}}h \le s_*(t+h) - s^*(t) \le s^*(t+h) - s_*(t) \le -\frac{2}{\sqrt{3}}h$$

for every $t, h \ge 0$. Therefore

$$s_*(t) = s^*(t) = \frac{2}{\sqrt{3}}t,$$

and we conclude that s^{ϵ} converges uniformly.

Finally set

$$u^*(y,t) = \limsup_{\epsilon \to 0, \ (z,r) \to (y,t)} u^{\epsilon}(z,r), \qquad u_*(y,t) = \liminf_{\epsilon \to 0, \ (z,r) \to (y,t)} u^{\epsilon}(z,r).$$

TT

1

:

Then from the theory of viscosity solutions (see e.g. [6, 7]) it follows that u^* is a viscosity subsolution of (5.3), and u_* is a viscosity supersolution of (5.3). Moreover

$$u^*(0,t) = u_*(0,t) = \frac{2}{\sqrt{3}}t.$$

Since there is a unique viscosity solution u of (5.3) satisfying the above (Dirichlet) boundary condition, a standard comparison theorem yields $u^* = u_* = u$.

In this paper, we have developed a weak theory for geometric equations of the type

$$V_{ij} = \mu_{ij}[e_i - e_j],$$

where V_{ij} is the normal velocity of an interface and $\mu_{ij} > 0$, e_1, e_2, e_3 are given constants. One may also consider more general geometric equations

$$V_{ij} = v_{ij}$$

with given constants v_{ij} . In general for any given v_{ij} 's there may not be any μ_{ij} and e_i satisfying $v_{ij} = \mu_{ij}[e_i - e_j]$ for every $i \neq j = 1, 2, 3$. In this non-physical situation, there is no convergence and our theory does not apply. We illustrate this lack of convergence and hence the lack of solutions with a simple example. Consider the problem

$$V_{ij} = 1, \quad i \neq j = 1, 2, 3,$$

with uniform angle conditions and initial data of three half lines meeting at the origin with equal angles. Let $\Gamma_{ij}^{\epsilon}(t)$ be the solution of the curvature perturbed problem. By symmetry, each one is obtained from another one by a rotation of 120° and the triple junction remains at the origin. Thus it suffices to consider the evolution of only one of the arcs, Γ_{13} . Suppose that $\Gamma_{13}(0)$ coincides with the x-axis. Then by an elementary argument shows that as, $\epsilon \to 0$, $\Gamma_{13}(t)$ spirals around the origin infinitely many times "converging" to

$$\{(x,y): x^2 + y^2 \le t\} \cup \{(x,t): x \ge 0\}.$$

Hence there is no limit of the vanishing surface tension problem. A numerical study of this example can be found in [9, Figure 28].

Acknowledgments.

This work was partially supported by the Army Research Office and the National Science Foundation through the Center for Nonlinear Analysis. The second author gratefully acknowledges support from the National Science Foundation through grant DMS-9200801.

References

- S. M. Allen and J. W. Cahn, A microscopic theory for antiphase boundary motion and its application to antiphase domain coarsening, Acta Metall. 27 (1979), 1085-1095.
- [2] S. Angenent and M. E. Gurtin, Multiphase thermomechanics with interfacial structure 2. Evolution of an isothermal interface, Arch. Rational Mech. Anal. 108 (1989), 323-391.
- [3] H. V. Atkinson, Theories of normal grain growth in pure single phase systems, Acta Metall. 36 (1988), 469-491.
- [4] G. Barles, H. M. Soner and P. Souganidis, Front propagation and phase filed theory, SIAM Control & Opt., 31/2 (1993),439-469.
- [5] L. Bronsard and F. Reitich, On three-phase boundary motion and the singular limit of a vector-valued Ginzburg-Landau equation, Arch. Rational Mech. Anal. 124 (1993), 355-379.
- [6] M. Crandall, H. Ishii and P.-L. Lions, User's guide to viscosity solutions of second order partial differential equations, Bulletin AMS, 27/1 (1992), 1-68.
- [7] W. Fleming and H. M. Soner, Controlled Markov Processes and Viscosity Solutions, Springer Verlag, New York, (1993).
- [8] M. Gurtin, P. Souganidis and H.M. Soner, Anisotropic planar motion of an interface relaxed by the formation of infinitesimal wrinkles, Journal of Differential Equations, forthcoming.
- [9] B. Merriman, J. K. Bence and S. J. Osher, Motion of multiple junctions: a level set approach, preprint.
- [10] W. Mullins, Two dimensional motion of idealized grain boundaries, J. Appl. Phys. 27 (1956), 900-904.
- [11] C. S. Smith, Grain shapes and other metallurgical applications of topology, in Metal Interfaces, American Society for Metals, Cleveland (1952), 65-108.
- [12] J. E. Taylor, The motion of multi-phase junctions under prescribed phaseboundary velocities, preprint.

37

1

2

Figure 10: A catalog of weak solutions for compatible initial data: (a) $\alpha > \beta > \alpha/2 > 0$; (b) $\alpha/2 > \beta > 0$; (c) $\alpha > -\beta > 0$; (d) $-\beta > \alpha > 0$; (e) $-\alpha > -\beta > 0$.

Figure 11: The weak solution for incompatible initial data. (a) One of the angle conditions is preserved; (b) No angle condition is preserved is the strong sense.

1

Б. Н

