

**NOTICE WARNING CONCERNING COPYRIGHT RESTRICTIONS:**

The copyright law of the United States (title 17, U.S. Code) governs the making of photocopies or other reproductions of copyrighted material. Any copying of this document without permission of its author may be prohibited by law.

**NAMT**

**96-003**

**Regularity Results for Equilibria in a  
Variational Model for Fracture**

**Emilio Acerbi  
Universita di Parma**

**Irene Fonseca  
Carnegie Mellon University**

**Nicola Fusco  
Universita di Firenze  
Research Report No. 96-NA-003**

**February 1996**

**Sponsors**

**U.S. Army Research Office  
Research Triangle Park  
NC 27709**

**National Science Foundation  
1800 G Street, N.W.  
Washington, DC 20550**

University Libraries  
Carnegie Mellon University  
Pittsburgh PA 15213-3890

TMAU  
DR-003

## REGULARITY RESULTS FOR EQUILIBRIA IN A VARIATIONAL MODEL FOR FRACTURE

Emilio Acerbi <sup>1</sup> Irene Fonseca <sup>2</sup> Nicola Fusco <sup>3</sup>

### Abstract

In recent years models describing interactions between fracture and damage have been proposed in which the relaxed energy of the material is given by a functional involving bulk and interfacial terms, of the form

$$\mathcal{G}(K, u) := \int_{\Omega \setminus K} F(\nabla u) dx + \lambda \int_{\Omega \setminus K} |u - g|^q dx + \beta H^{N-1}(K \cap \Omega),$$

where  $\Omega$  is an open, bounded subset of  $\mathbb{R}^N$ ,  $q \geq 1$ ,  $g \in L^\infty(\Omega; \mathbb{R}^N)$ ,  $\lambda, \beta > 0$ , the bulk energy density  $F$  is quasiconvex,  $K \subset \mathbb{R}^N$  is closed, and the admissible deformation  $u : \Omega \rightarrow \mathbb{R}^N$  is  $C^1$  in  $\Omega \setminus K$ . One of the main issues has to do with regularity properties of the “crack site”  $K$  for a minimizing pair  $(K, u)$ . In the scalar case, i.e. when  $u : \Omega \rightarrow \mathbb{R}$ , similar models were adopted to image segmentation problems, and the regularity of the “edge” set  $K$  has been successfully resolved for a quite broad class of convex functions  $F$  with growth  $p > 1$  at infinity. In turn, this regularity entails the existence of classical solutions. The methods thus used cannot be carried out to the vectorial case, except for a very restrictive class of integrands. In this paper we deal with a vector-valued case on the plane, obtaining regularity for minimizers of  $\mathcal{G}$  corresponding to polyconvex bulk energy densities of the form

$$F(\xi) = \frac{1}{2} |\xi|^2 + h(\det \xi),$$

where the convex function  $h$  grows linearly at infinity.

*1991 Mathematics subject classification (Amer. Math. Soc.):* 35J20, 49Q20, 49J45, 49N60

*Key Words :* quasiconvexity, recession function, regularity, bounded variation

<sup>1</sup> Dipartimento di Matematica, Via Massimo D’Azeglio 85/A, 43100 Parma (Italy). Research supported by MURST, Gruppo Nazionale 40%.

<sup>2</sup> Department of Mathematics, Carnegie Mellon University, Pittsburgh, PA 15213. Research partially supported by the Army Research Office and the National Science Foundation through the Center for Nonlinear Analysis, and by the National Science Foundation under Grant No. DMS-9500531.

<sup>3</sup> Dipartimento di Matematica “U. Dini”, Università di Firenze, Viale Morgagni 67/a, 50131 Firenze (Italy). Research supported by MURST, Gruppo Nazionale 40%.

## 1. Introduction

In recent years models involving bulk and interfacial energies have been used in the contexts of fracture mechanics, phase transitions, and image segmentation in computer vision (see [BZ], [DGCL], [FFr], [MS]). The underlying quasistatic problems deal with minimization of an energy functional of the form

$$\mathcal{G}(K, u) := \int_{\Omega \setminus K} F(\nabla u) dx + \lambda \int_{\Omega \setminus K} |u - g|^q dx + \beta H^{N-1}(K \cap \Omega)$$

where  $\Omega$  is an open, bounded subset of  $\mathbb{R}^N$ ,  $q > 1$ ,  $g \in L^\infty(\Omega; \mathbb{R}^k)$ ,  $\lambda, \beta > 0$ , and  $H^{N-1}$  stands for the  $(N - 1)$ -dimensional Hausdorff measure, among all pairs  $(K, u)$  with  $K$  closed in  $\mathbb{R}^N$  and  $u : \Omega \rightarrow \mathbb{R}^k$  is smooth; the first two terms in  $\mathcal{G}$  represent the bulk energy and the last one accounts for interfacial energy. In order to attack this problem De Giorgi and Ambrosio (see [A1], [DGA]) introduced the space *SBV* of functions of *special bounded variation*, i.e. *BV* functions  $u$  whose distributional derivative  $Du$ , which is a finite, Radon measure, may be decomposed into an absolutely continuous part  $\nabla u \mathcal{L}^N$  with respect to the  $N$ -dimensional Lebesgue measure  $\mathcal{L}^N$ , and a singular part whose support is an  $(N - 1)$ -dimensional rectifiable set  $S_u$ . This is the "jump set" of  $u$ , in the sense that  $u$  has traces  $u^+ \neq u^-$   $H^{N-1}$ -a.e. on the two sides of  $S_u$  (for example, the Cantor-Vitali function is *BV* but not *SBV*). Then to the functional  $\mathcal{G}$  one may associate the functional  $\mathcal{F}$  defined on *SBV* as

$$\mathcal{F}(u) := \int_{\Omega} F(\nabla u) dx + \lambda \int_{\Omega} |u - g|^q dx + \beta H^{N-1}(S_u \cap \Omega).$$

Due to the relaxation result of Fonseca and Francfort (see [FFr]), and under suitable growth conditions for the bulk energy density  $F$ , it is not restrictive to take  $F$  quasiconvex. We recall that  $F$  is said to be *quasiconvex* if

$$F(\xi) \leq \int_Q F(\xi + \nabla \varphi(x)) dx$$

for all  $k \times N$  matrix  $\xi$ ,  $\varphi \in C_0^\infty(Q; \mathbb{R}^k)$ , and where  $Q = (0, 1)^N$ . A particular class of quasiconvex functions which plays an important role in elasticity is the class of *polyconvex* functions, i.e. convex functions of all minors of the matrix  $\xi$ .

Under the quasiconvexity assumption, and if  $F$  is superlinear and  $g$  is bounded, the lower semicontinuity and compactness results of Ambrosio (see [A1], [A2], [A3]) yield the existence of a *SBV* minimizer of  $\mathcal{F}$ . Note that, in general, if  $u \in \text{SBV}$  then the set  $S_u$  is far from being closed, i.e.  $H^{N-1}((\overline{S_u} \setminus S_u) \cap \Omega) > 0$ .

In the scalar-valued case where  $u : \Omega \rightarrow \mathbb{R}$ , quasiconvexity reduces to convexity and De Giorgi, Carriero and Leaci [DGCL] proved that if

$$F(\xi) = |\xi|^2$$

and if  $g$  is bounded, then

$$H^{N-1}((\overline{S_u} \setminus S_u) \cap \Omega) = 0$$

for any minimizer  $u \in SBV(\Omega; \mathbb{R})$  of  $\mathcal{F}$ . From this property one immediately concludes that the pair  $(\overline{S_u}, u)$  is a minimizer of the original functional  $\mathcal{G}$ . The result of [DGCL] has been extended to a quite broad class of convex functions  $F$  with growth  $p > 1$  at infinity, and further regularity on the set  $S_u$  has been obtained (see [AFP], [AP], [B], [DG], [DS], [FFu]).

In this paper we deal with a two-dimensional, vector-valued, polyconvex case, considering a class of functionals  $\mathcal{G}$  which includes in particular the model

$$\int_{\Omega \setminus K} \left[ \frac{1}{2} |\nabla u|^2 + |\det \nabla u| \right] dx + \lambda \int_{\Omega \setminus K} |u - g|^q dx + \beta H^1(K \cap \Omega),$$

where  $u \in W^{1,2}(\Omega \setminus K; \mathbb{R}^2)$  and  $g$  is bounded. In Theorem 2.1 we prove that this functional admits a minimizing pair of the form  $(\overline{S_u}, u)$ , where  $u \in SBV(\Omega; \mathbb{R}^2)$  and, moreover,  $u \in C^{0,\alpha}(\Omega \setminus \overline{S_u}; \mathbb{R}^2)$  for every  $0 < \alpha < 1$ .

## 2. Statements and Auxiliary Results

If  $\Omega \subset \mathbb{R}^N$  is open, we say that a function of *bounded variation*  $u \in BV(\Omega; \mathbb{R}^k)$  is a function of *special bounded variation*,  $u \in SBV(\Omega; \mathbb{R}^k)$ , if, denoting by  $S_u$  the complement of the set of Lebesgue points of  $u$ , the distributional derivative  $Du$  is represented by

$$Du = \nabla u \mathcal{L}^N + (u^+ - u^-) \otimes \nu H^{N-1} \llcorner S_u,$$

where  $\nabla u$  is the Radon-Nikodym derivative of the finite, Radon measure  $Du$  with respect to the  $N$ -dimensional Lebesgue measure  $\mathcal{L}^N$ ,  $\nu$  is the normal to the rectifiable set  $S_u$ ,  $u^+$  and  $u^-$  are the traces of  $u$  on  $S_u$ , and  $H^{N-1}$  denotes the  $(N-1)$ -dimensional Hausdorff measure.

We recall that the *recession function*  $h^\infty$  of a convex function  $h : \mathbb{R} \rightarrow [0, +\infty)$  is defined by

$$h^\infty(t) := \lim_{s \rightarrow +\infty} \frac{h(st)}{s},$$

and it is convex and positively homogeneous of degree one.

In this paper we prove the following theorem.

**Theorem 2.1.** *Let  $\Omega$  be a bounded, open subset of  $\mathbb{R}^2$ . Let  $h : \mathbb{R} \rightarrow [0, \infty)$  be a convex function such that  $h(0) \leq h(t) \leq C(1 + |t|)$ , and*

$$\left| h^\infty(t) - \frac{h(st)}{s} \right| \leq \frac{C}{s^m}$$

for some  $0 < m < 1$ , and for all  $t \in \mathbb{R}$ ,  $s \geq s_0 > 0$ . Let  $\lambda, \beta > 0$  and define

$$\mathcal{G}(K, u) := \int_{\Omega \setminus K} \left[ \frac{1}{2} |\nabla u|^2 + h(\det \nabla u) \right] dx + \lambda \int_{\Omega \setminus K} |u - g|^q dx + \beta H^1(\Omega \cap K).$$

There exists a minimizer of  $\mathcal{G}(\cdot, \cdot)$  of the form  $(\overline{S_u}, u)$ , with  $u \in SBV(\Omega, \mathbb{R}^2)$ , among all pairs  $(K, v)$  with  $K \subset \Omega$  closed and  $v \in W^{1,2}(\Omega \setminus K; \mathbb{R}^2)$ . Moreover,

$$H^1((\overline{S_u} \setminus S_u) \cap \Omega) = 0.$$

In order to prove Theorem 2.1, we invite the reader to follow the same steps taken in the proof of Theorem 4.1 of [FFu]. It is easy to verify that all the auxiliary lemmas hold in this case, with one exception: the Density Lower Bound ([FFu], Lemma 4.3) uses the Decay Lemma ([FFu], Lemma 4.4), which was proved by Carriero and Leaci [CL] only in the  $p$ -harmonic case, i.e.  $F(\xi) = |\xi|^p$ . Knowing that local minimizers are locally Lipschitz functions played a fundamental role in the analysis.

We introduce the notation

$$\mathcal{F}_0(u, c, A) := \int_A \left[ \frac{1}{2} |\nabla u|^2 + h^\infty(\det \nabla u) \right] dx + cH^1(S_u \cap \overline{A}),$$

$$\Phi_0(u, c, A) := \inf \{ \mathcal{F}_0(w, c, A) : w \in SBV(\Omega; \mathbb{R}^2), w = u \text{ in } \Omega \setminus \overline{A} \},$$

$$\Psi_0(u, c, A) := \mathcal{F}_0(u, c, A) - \Phi_0(u, c, A),$$

where  $A$  is an open subset of  $\Omega$ ,  $c > 0$ ,  $u \in SBV(\Omega; \mathbb{R}^2)$ . In the case where  $u \in W^{1,2}(A; \mathbb{R}^2)$  we simply write

$$\mathcal{F}_0(u, A) := \int_A \left[ \frac{1}{2} |\nabla u|^2 + h^\infty(\det \nabla u) \right] dx.$$

In the sequel, we use the notation  $B_R$ ,  $R > 0$ , to denote a generic open ball of radius  $R$ , centered at  $x \in \Omega$ , such that  $B_R \subset \Omega$ .

**Definition.** We say that  $u \in W^{1,2}(\Omega)$  is a  $W^{1,2}$ -local minimizer of

$$I(v; \Omega) := \int_\Omega F(\nabla v) dx, \quad v \in W^{1,2}(\Omega)$$

if

$$I(u; B_R(x_0)) = \min \left\{ I(v; B_R(x_0)) : v \in u + W_0^{1,2}(B_R(x_0)) \right\}$$

for all balls  $B_R(x_0) \subset \Omega$ .

We now state the version of the decay lemma which holds in our case. The density lower bound, and thus Theorem 2.1, follow from the decay lemma by the same argument used in [FFu].

**Lemma 2.2. [Decay Lemma]** For all  $\gamma \in (0, 1)$  there exists  $\tau_\gamma \in (0, 1)$  such that for every  $\tau \in (0, \tau_\gamma)$  and for every  $c > 0$  there exist  $\varepsilon = \varepsilon(c, \tau, \gamma)$ ,  $\theta = \theta(c, \tau, \gamma)$ ,  $R_0 = R_0(c, \tau, \gamma)$ , such that if  $0 < \rho < R_0$ , and if  $u \in SBV(\Omega; \mathbb{R}^2)$  is such that  $\mathcal{F}_0(u, c, B_\rho) \leq \varepsilon^2 \rho$  and  $\Psi_0(u, c, B_\rho) \leq \theta \mathcal{F}_0(u, c, B_\rho)$ , then

$$\mathcal{F}_0(u, c, B_{\tau\rho}) \leq \tau^{2-\gamma} \mathcal{F}_0(u, c, B_\rho).$$

In order to prove Lemma 2.2, we follow the proof of Lemma 3.12 in [FFu], and we suppose that the result is not true; then there exist  $\gamma \in (0, 1)$ , for every  $\tau_\gamma \in (0, 1)$  there exists  $\tau \in (0, \tau_\gamma)$ , there exist  $c > 0$ , two sequences  $\{\varepsilon_h\}$ ,  $\{\theta_h\}$ , with  $\lim_h \varepsilon_h = \lim_h \theta_h = 0$ , a sequence  $\{u_h\} \subset SBV(\Omega; \mathbb{R}^2)$ , and a sequence of balls  $B_{R_h}(x_h) \subset\subset \Omega$  with  $\lim_h R_h = 0$  such that

$$\mathcal{F}_0(u_h, c, B_{R_h}(x_h)) = \varepsilon_h R_h, \quad \Psi_0(u_h, c, B_{R_h}(x_h)) = \theta_h \mathcal{F}_0(u_h, c, B_{R_h}(x_h)),$$

and

$$\mathcal{F}_0(u_h, c, B_{\tau R_h}(x_h)) > \tau^{2-\gamma} \mathcal{F}_0(u_h, c, B_{R_h}(x_h)).$$

After rescaling, it is easily seen that the rest of the proof can be carried out in a similar way to Lemma 3.12 in [FFu], provided an estimate of the type

$$\int_{B_\tau} \left[ \frac{1}{2} |\nabla u|^2 + h^\infty(\det \nabla u) \right] dx \leq C \tau^{2-\alpha} \int_{B_1} \left[ \frac{1}{2} |\nabla u|^2 + h^\infty(\det \nabla u) \right] dx$$

holds for local minimizers of  $\mathcal{F}_0$  in  $W^{1,2}(B_1; \mathbb{R}^2)$ , for any  $0 < \alpha < 1$  or, equivalently,

**Theorem 2.3.** *If  $u \in W^{1,2}(B_1, \mathbb{R}^2)$  is a  $W^{1,2}$ -local minimizer of  $\mathcal{F}_0$  then*

$$\int_{B_\tau} |\nabla u|^2 dx \leq C \tau^{2-\alpha} \int_{B_1} |\nabla u|^2 dx \quad (2.1)$$

for all  $\alpha, \tau \in (0, 1)$  and some constant  $C > 0$ ,  $C \equiv C(\alpha, h^\infty)$ .

This is, therefore, the only result still needed to prove Theorem 2.1. As an immediate consequence of (2.1), it follows that  $u \in C^{0,\alpha}$  for all  $\alpha \in (0, 1)$  (see [G], Theorem 1.1, Chapter 3).

The rest of the paper will be dedicated to proving Theorem 2.3. Although we will rely strongly on the arguments used by Dougherty [D] to obtain higher integrability of local minimizers in the case where  $h \in C^1(\mathbb{R})$ , we could not find an easier way to adapt directly his proof to more general Lipschitz functions  $h$  by means of a simple approximation and density approach.

The following lemma may be found in [G] (Chap. 3, Lemma 2.1) in the case where with  $\gamma = \beta$ ; although stated in a slightly weaker form, its proof yields the result below.

**Lemma 2.4.** *Let  $\phi : [0, +\infty) \rightarrow [0, +\infty)$  be a nonnegative, nondecreasing function, such that*

$$\phi(\rho) \leq H \left( \frac{\rho}{R} \right)^\alpha \phi(R) + K R^\beta$$

for all  $0 < \rho < R \leq R_0$  and for some constants  $H, K \geq 0$  and  $0 < \beta < \alpha$ . Then for every  $\gamma \in [\beta, \alpha)$  there exists a constant  $C = C(H, \alpha, \beta, \gamma)$  such that

$$\phi(\rho) \leq C \left[ \left( \frac{\rho}{R} \right)^\gamma \phi(R) + K \rho^\beta \right]$$

for all  $0 < \rho < R \leq R_0$ .



Next, we prove a decay estimate for solutions of a well known elliptic equation.

**Lemma 2.5.** *Let  $\Omega \subset \mathbb{R}^N$  be an open, bounded domain. If  $f \in L^p(\Omega; \mathbb{R}^N)$ ,  $p \geq 2$ , and if  $v \in W^{1,2}(\Omega)$  satisfies*

$$\Delta v = \operatorname{div} f \quad \text{in } \Omega$$

*then for any  $\delta \in (0, 2N/p)$  there exists a constant  $C = C(p, N, \delta)$  such that if  $B_R \subset \Omega$  and  $0 < \rho < R$  then*

$$\int_{B_\rho} |Dv|^2 dx \leq C \left[ \left( \frac{\rho}{R} \right)^{N-\delta} \int_{B_R} |Dv|^2 dx + \rho^{N-2N/p} \left( \int_{B_R} |f|^p dx \right)^{2/p} \right]. \quad (2.2)$$

*Proof.* Fix a ball  $B_R \subset \Omega$ , and let  $w$  be the harmonic function which minimizes

$$z \mapsto \int_{B_R} |\nabla z|^2 dx, \quad z \in v + W_0^{1,2}(B_R).$$

Since  $|Dw|^2$  is subharmonic we get

$$\int_{B_\rho} |Dw|^2 dx \leq \left( \frac{\rho}{R} \right)^N \int_{B_R} |Dw|^2 dx \quad (2.3)$$

for all  $\rho < R$ .

We claim that if  $\rho < R$  then

$$\int_{B_\rho} |Dv|^2 dx \leq 2 \left( \frac{\rho}{R} \right)^N \int_{B_R} |Dv|^2 dx + C R^{N-2N/p} \left( \int_{B_R} |f|^p dx \right)^{2/p}. \quad (2.4)$$

If (2.4) holds, then (2.2) follows from Lemma 2.4, setting

$$\phi(\rho) := \int_{B_\rho} |\nabla v|^2 dx, \quad \alpha := N, \quad \beta := N - \frac{2N}{p}, \quad \gamma := N - \delta.$$

To prove the claim, we start by noting that

$$\begin{aligned} \int_{B_R} Dv \cdot (Dv - Dw) dx &= \int_{B_R} f \cdot (Dv - Dw) dx, \\ \int_{B_R} Dw \cdot (Dv - Dw) dx &= 0, \end{aligned}$$

so that

$$\int_{B_R} |Dv - Dw|^2 dx = \int_{B_R} f \cdot (Dv - Dw) dx.$$

Using Cauchy-Schwartz, Young's, and Hölder's inequality, we obtain

$$\int_{B_R} |Dv - Dw|^2 dx \leq \int_{B_R} |f|^2 dx \leq C R^{N-2N/p} \left( \int_{B_R} |f|^p dx \right)^{2/p},$$

which, together with (2.3), yields

$$\begin{aligned} \int_{B_\rho} |Dv|^2 dx &\leq 2 \left( \int_{B_\rho} |Dw|^2 dx + \int_{B_R} |Dv - Dw|^2 dx \right) \\ &\leq 2 \left( \frac{\rho}{R} \right)^N \int_{B_R} |Dw|^2 dx + C R^{N-2N/p} \left( \int_{B_R} |f|^p dx \right)^{2/p}. \end{aligned}$$

This inequality reduces to (2.4) since, by definition of  $w$ ,

$$\int_{B_R} |Dw|^2 dx \leq \int_{B_R} |Dv|^2 dx.$$

□

### 3. The Main Theorem

The main result of this section is the following.

**Theorem 3.1.** *Let  $\Omega$  be an open subset of  $\mathbb{R}^2$ , and let  $u \in W^{1,2}(\Omega, \mathbb{R}^2)$  be a  $W^{1,2}$ -local minimizer of*

$$\mathbb{F}(v; \Omega) := \int_{\Omega} \left[ \frac{1}{2} |Dv|^2 + M |\det Dv| \right] dx$$

with  $M > 0$ . If  $\alpha \in (0, 1)$ ,  $B_R \subset \Omega$ , and if  $\rho < R$ , then

$$\int_{B_\rho} |Du|^2 dx \leq C \left( \frac{\rho}{R} \right)^{2\alpha} \int_{B_R} |Du|^2 dx$$

for some constant  $C$ . In particular,  $u \in C^{0,\alpha}(\Omega; \mathbb{R}^2)$ .

Before proving this result, we show how Theorem 2.3 can be derived as a simple corollary.

**Proof of Theorem 2.3.** Let  $u$  be a  $W^{1,2}$ -local minimizer of

$$\mathcal{F}_0(v, A) := \int_A \left[ \frac{1}{2} |\nabla v|^2 + h^\infty(\det \nabla v) \right] dx, \quad v \in W^{1,2}(\Omega; \mathbb{R}^2),$$

where the recession function is given by

$$h^\infty(t) = \begin{cases} at & \text{if } t > 0 \\ -bt & \text{if } t < 0 \end{cases}$$

for some  $a, b \geq 0$ . Since  $v \mapsto \det \nabla v$  is a null-lagrangian, i.e.

$$\int_A \det \nabla v dx = \int_A \det \nabla w dx$$

whenever  $v, w \in W^{1,2}(\Omega; \mathbb{R}^2)$ ,  $A \subset \Omega$  has Lipschitz boundary and  $v = w$  on  $\partial A$ , we conclude that  $u$  is a  $W^{1,2}$ -local minimizer of

$$\begin{aligned} \mathcal{F}_1(v; A) &= \int_A \left[ \frac{1}{2} |\nabla v|^2 + a(\det \nabla v)^+ + b(\det \nabla v)^- + \frac{b-a}{2} \det \nabla v \right] dx \\ &= \int_A \left[ \frac{1}{2} |\nabla v|^2 + \frac{a+b}{2} |\det \nabla v| \right] dx. \end{aligned}$$

The result now follows from Theorem 3.1.  $\square$

Next we recall some algebraic inequalities used by Dougherty [D], and which will enter in the proof of Theorem 3.1. For completeness, we include the proofs.

Consider a  $2 \times 2$  matrix

$$D = \begin{pmatrix} a & b \\ c & d \end{pmatrix}.$$

Then

$$\det D = ad - bc, \quad \text{adj } D = \begin{pmatrix} d & -c \\ -b & a \end{pmatrix}, \quad \text{tr}(D^T D) = |D|^2 = a^2 + b^2 + c^2 + d^2,$$

where  $(\text{adj } D)^T D = \det D \mathbf{I}$ , and we define

$$A := \frac{1}{2}(a^2 + c^2 - b^2 - d^2), \quad B := ab + cd, \quad \sigma := \text{sign}(\det D) \text{ whenever } \det D \neq 0.$$

**Lemma 3.2.** *The following inequalities hold:*

- (i)  $|A| + |B| \leq |D|^2$ ;
- (ii)  $|D|^2 \leq 2(|A| + |B|)$  if  $\det D = 0$ ;
- (iii)  $|\sigma \text{adj } D - D| \leq 4\sqrt{|A| + |B|}$  if  $\det D \neq 0$ .

*Proof.* It is clear that

$$\begin{aligned} |A| + |B| &\leq \frac{1}{2}(|a| + |b|)^2 + \frac{1}{2}(|c| + |d|)^2 \\ &\leq a^2 + b^2 + c^2 + d^2, \end{aligned}$$

proving (i). Also, consider the right stretching tensor  $U := \sqrt{D^T D}$ ; then its eigenvalues  $\nu_1 \geq \nu_2 \geq 0$  are the principal stretches of  $D$ , and it can be seen easily that

$$|\det D| = \nu_1 \nu_2, \quad |D|^2 = \nu_1^2 + \nu_2^2.$$

By virtue of the polar decomposition, we may find rotations  $R, Q \in SO(2)$  such that

$$D = RU, \quad U = Q^T \begin{pmatrix} \nu_1 & 0 \\ 0 & \nu_2 \end{pmatrix} Q,$$

and so

$$\begin{aligned} \begin{pmatrix} A & B \\ B & -A \end{pmatrix} &= D^T D - \frac{1}{2}|D|^2 \mathbf{I} \\ &= \frac{1}{2} Q^T \begin{pmatrix} \nu_1^2 - \nu_2^2 & 0 \\ 0 & \nu_2^2 - \nu_1^2 \end{pmatrix} Q. \end{aligned}$$

Hence

$$\nu_1^2 - \nu_2^2 = 2\sqrt{|A|^2 + |B|^2}. \quad (3.1)$$

If  $\det D = 0$  then  $\nu_2 = 0$ , and

$$|D|^2 = \nu_1^2 = 2\sqrt{|A|^2 + |B|^2} \leq 2(|A| + |B|),$$

and we conclude (ii). If  $\det D > 0$  then

$$\begin{aligned} |\operatorname{adj} D - D| &= \left| \begin{pmatrix} d-a & -c-b \\ -b-c & a-d \end{pmatrix} \right| \\ &= \sqrt{2|a-d|^2 + 2|c+b|^2} \\ &\leq 2(\nu_1 - \nu_2), \end{aligned} \quad (3.2)$$

because (as a direct computation shows)

$$\begin{aligned} |a-d| &= \left| \operatorname{tr} \left[ \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} D \right] \right| \\ &= \operatorname{tr} \left[ \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} R Q^T \begin{pmatrix} \nu_1 & 0 \\ 0 & \nu_2 \end{pmatrix} Q \right] \\ &\leq \nu_1 - \nu_2, \end{aligned}$$

$$\begin{aligned} |b+c| &= |D_{12} + D_{21}| \\ &= \left| \left[ R Q^T \begin{pmatrix} \nu_1 & 0 \\ 0 & \nu_2 \end{pmatrix} Q \right]_{12} + \left[ R Q^T \begin{pmatrix} \nu_1 & 0 \\ 0 & \nu_2 \end{pmatrix} Q \right]_{21} \right| \\ &\leq \nu_1 - \nu_2. \end{aligned}$$

Finally, (3.1), (3.2), and (i) yield

$$\begin{aligned} |\operatorname{adj} D - D| &\leq 2(\nu_1 - \nu_2) = 4 \frac{\sqrt{|A|^2 + |B|^2}}{\nu_1 + \nu_2} \\ &\leq 4 \frac{\sqrt{|A|^2 + |B|^2}}{|D|} \leq 4 \frac{\sqrt{|A|^2 + |B|^2}}{\sqrt{|A| + |B|}} \leq 4\sqrt{|A| + |B|}. \end{aligned}$$

If  $\det D < 0$  then set

$$\tilde{D} := R_0 D, \quad \text{where } R_0 := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

With obvious notation we have

$$A = \tilde{A}, \quad B = \tilde{B}$$

while  $\det \tilde{D} > 0$ , and so

$$\begin{aligned}
|\sigma \operatorname{adj} D - D| &= |R_0^T \operatorname{adj} R_0 D - D| \\
&= |\operatorname{adj} \tilde{D} - \tilde{D}| \\
&\leq 4\sqrt{|A| + |B|}.
\end{aligned}$$

□

To avoid overburdening the reader with indices, we change the notation just for the length of the proof of Theorem 3.1, conforming to the one employed in [D]: we denote by  $\mathcal{U}$  the local minimizer of  $\mathbb{F}$ , replacing the former  $u$ . Then we are free to denote by  $(u, v)$  the components of  $\mathcal{U}$ , whereas the variables are  $X = (x, y)$ . We also use  $\Phi = (\phi, \psi)$  to designate any smooth function  $\Phi \in C_0^1(\Omega; \mathbb{R}^2)$  with support in a ball  $B_R \subset \Omega$ .

**Proof of Theorem 3.1. Step 1.** Following the argument of Dougherty [D], we consider a variation of the domain for the local minimizer  $\mathcal{U}$ , i.e. we study the variation  $\mathcal{U}(I + \varepsilon\Phi)$  near  $\varepsilon = 0$ , obtaining a first set of Euler-Lagrange equations.

Let  $B_R \subset\subset \Omega$ , fix  $\Phi \in C_0^1(B_R, \mathbb{R}^2)$ , and  $\varepsilon > 0$  small. Setting  $\mathcal{U}_\varepsilon(Y) := \mathcal{U}(Y + \varepsilon\Phi(Y))$ , due to the minimality of  $\mathcal{U}$  we have

$$\mathbb{F}(\mathcal{U}_\varepsilon; B_R) - \mathbb{F}(\mathcal{U}; B_R) \geq 0,$$

and since a simple change of variables yields

$$\int_{B_R} |\det D\mathcal{U}_\varepsilon| dY = \int_{B_R} |\det D\mathcal{U}| dX,$$

we deduce that

$$\int_{B_R} (|D\mathcal{U}_\varepsilon|^2 - |D\mathcal{U}|^2) dY \geq 0,$$

and so

$$\int_{B_R} A(\phi_x - \psi_y) + B(\phi_y + \psi_x) dX = 0, \quad (3.3)$$

where, using the notations introduced above,

$$D = \begin{pmatrix} u_x & u_y \\ v_x & v_y \end{pmatrix},$$

$$A := \frac{1}{2}(u_x^2 + v_x^2 - u_y^2 - v_y^2)$$

$$B := u_x u_y + v_x v_y.$$

Hence  $A, B \in L^1(\Omega)$ , and (3.3) can be written, in the sense of distributions, as

$$\begin{cases} A_x + B_y = 0 \\ B_x - A_y = 0, \end{cases}$$

and so  $A, B$  are harmonic in  $B_R$ . In particular,

$$\sup_{B_{R/2}} (|A| + |B|) \leq \frac{4}{\pi R^2} \int_{B_R} (|A| + |B|) dX. \quad (3.4)$$

Step 2 . For  $t \geq 0$  define

$$\begin{aligned}\Omega_t^+ &:= \{X \in \Omega : \det DU > t\}, \\ \Omega_t^- &:= \{X \in \Omega : \det DU < -t\}, \\ \Omega_t &:= \{X \in \Omega : |\det DU| \leq t\}.\end{aligned}$$

Let  $\sigma(X)$  denote the sign of  $\det DU(X)$  whenever the determinant is not equal to zero. Clearly, if  $t > 0$  and  $\varepsilon$  is sufficiently small then  $X \in \Omega_t^+$  implies that  $\det(DU + \varepsilon\Phi)(X) > 0$ , while  $\det(DU + \varepsilon\Phi)(X) < 0$  whenever  $X \in \Omega_t^-$ . Therefore

$$\begin{aligned}& \int_{\Omega} \frac{1}{2} \frac{|D(U + \varepsilon\Phi)|^2 - |DU|^2}{\varepsilon} dx + M \int_{\Omega_t^+} \frac{\det D(U + \varepsilon\Phi) - \det DU}{\varepsilon} dx \\ & - M \int_{\Omega_t^-} \frac{\det D(U + \varepsilon\Phi) - \det DU}{\varepsilon} dx + M \int_{\Omega_t} \frac{|\det D(U + \varepsilon\Phi)| - |\det DU|}{\varepsilon} dx \geq 0.\end{aligned}$$

Since

$$\det D(U + \varepsilon\Phi) = \det DU + \varepsilon \operatorname{adj} DU \cdot D\Phi + \varepsilon^2 \det D\Phi,$$

and

$$|\operatorname{adj} DU \cdot D\Phi| \geq \frac{|\det DU + \varepsilon \operatorname{adj} DU \cdot D\Phi| - |\det DU|}{\varepsilon},$$

we obtain, letting  $\varepsilon \rightarrow 0$ ,

$$\int_{\Omega} DU \cdot D\Phi dX + M \int_{\Omega \setminus \Omega_t} \sigma \operatorname{adj} DU \cdot D\Phi dX + M \int_{\Omega_t} |\operatorname{adj} DU \cdot D\Phi| dX \geq 0.$$

Replacing  $\Phi$  by  $-\Phi$ , this inequality reduces to

$$\int_{\Omega} DU \cdot D\Phi dX + M \int_{\Omega \setminus \Omega_t} \sigma \operatorname{adj} DU \cdot D\Phi dX - M \int_{\Omega_t} |\operatorname{adj} DU \cdot D\Phi| dX \leq 0,$$

or, equivalently,

$$\begin{aligned}(M+1) \int_{\Omega} DU \cdot D\Phi dX + M \int_{\Omega \setminus \Omega_t} (\sigma \operatorname{adj} DU \cdot D\Phi - DU \cdot D\Phi) dX \\ \leq M \int_{\Omega_t} (|\operatorname{adj} DU \cdot D\Phi| + DU \cdot D\Phi) dX.\end{aligned}$$

Letting  $t \rightarrow 0$ , and setting  $M' = 1 + 1/M$ , we conclude that

$$\begin{aligned}M' \int_{\Omega} DU \cdot D\Phi dX + \int_{\Omega \setminus \Omega_0} (\sigma \operatorname{adj} DU \cdot D\Phi - DU \cdot D\Phi) dX \\ \leq \int_{\Omega_0} (|\operatorname{adj} DU \cdot D\Phi| + DU \cdot D\Phi) dX.\end{aligned}\tag{3.5}$$

*Step 3* . We compare  $\mathcal{U}$  with the solution of a more regular problem: fix  $B_R \subset \Omega$ , and let  $\mathcal{V} \in W^{1,2}(B_{R/2}; \mathbb{R}^2)$  be the solution of

$$\begin{cases} M' \Delta \mathcal{V} = -\operatorname{div} [\chi_{\Omega \setminus \Omega_0} (\sigma \operatorname{adj} D\mathcal{U} - D\mathcal{U})] & \text{in } B_{R/2} \\ \mathcal{V} = \mathcal{U} & \text{on } \partial B_{R/2} . \end{cases} \quad (3.6)$$

By Lemma 3.2 (iii) it follows that the right hand side of the equation is the divergence of an  $L^2$  function, and so we have, in weak form,

$$\int_{B_{R/2}} M' D\mathcal{V} \cdot D\Phi \, dX + \int_{B_{R/2} \setminus \Omega_0} (\sigma \operatorname{adj} D\mathcal{U} - D\mathcal{U}) \cdot D\Phi \, dX = 0$$

for all  $\Phi \in W_0^{1,2}(B_{R/2}; \mathbb{R}^2)$ . Subtracting this equation from (3.5), and choosing  $\Phi = \mathcal{U} - \mathcal{V}$ , we obtain

$$\int_{B_{R/2}} M' |D\mathcal{U} - D\mathcal{V}|^2 \, dX \leq \int_{B_{R/2} \cap \Omega_0} (|\operatorname{adj} D\mathcal{U}| + |D\mathcal{U}|) |D\mathcal{U} - D\mathcal{V}| \, dX ,$$

and noticing that  $|\operatorname{adj} D\mathcal{U}| = |D\mathcal{U}|$ , by virtue of Cauchy-Schwartz inequality and Lemma 3.2 (ii), we have

$$\begin{aligned} \int_{B_{R/2}} |D\mathcal{U} - D\mathcal{V}|^2 \, dX &\leq 4 \int_{B_{R/2} \cap \Omega_0} |D\mathcal{U}|^2 \, dX \\ &\leq 8 \int_{B_{R/2} \cap \Omega_0} (|A| + |B|) \, dX . \end{aligned} \quad (3.7)$$

*Step 4* . Fix  $B_R \subset \Omega$ . Since  $A$  and  $B$  are harmonic functions, by Lemma 3.2 (iii) we have  $\chi_{\Omega \setminus \Omega_0} (\sigma \operatorname{adj} D\mathcal{U} - D\mathcal{U}) \in L_{\operatorname{loc}}^p(\Omega)$  for every  $p \geq 2$ , and so, applying Lemma 2.5 to (3.6) with  $\delta = 2/p$ , we obtain

$$\begin{aligned} &\int_{B_\rho} |D\mathcal{V}|^2 \, dX \\ &\leq C \left[ \left( \frac{\rho}{R} \right)^{2-2/p} \int_{B_{R/2}} |D\mathcal{V}|^2 \, dX + \rho^{2-4/p} \left( \int_{B_{R/2}} (|A|^{p/2} + |B|^{p/2}) \, dX \right)^{2/p} \right] \end{aligned}$$

for all  $\rho < R/2$ , where  $C = C(p)$ . This estimate, together with (3.7), yields

$$\begin{aligned}
& \int_{B_\rho} |DU|^2 dX \\
& \leq 2 \int_{B_\rho} |DV|^2 + 2 \int_{B_{R/2}} |DU - DV|^2 dX \\
& \leq C \left( \frac{\rho}{R} \right)^{2-2/p} \int_{B_{R/2}} |DV|^2 dX + C \rho^{2-4/p} \left( \int_{B_{R/2}} (|A|^{p/2} + |B|^{p/2}) dX \right)^{2/p} \\
& \quad + 2 \int_{B_{R/2}} |DU - DV|^2 dX \\
& \leq C \left[ \left( \frac{\rho}{R} \right)^{2-2/p} \int_{B_{R/2}} |DU|^2 dX + \rho^{2-4/p} \left( \int_{B_{R/2}} (|A|^{p/2} + |B|^{p/2}) dX \right)^{2/p} \right. \\
& \quad \left. + \int_{B_{R/2}} |DU - DV|^2 dX \right] \\
& \leq C \left[ \left( \frac{\rho}{R} \right)^{2-2/p} \int_{B_{R/2}} |DU|^2 dX + \rho^{2-4/p} \left( \int_{B_{R/2}} (|A|^{p/2} + |B|^{p/2}) dX \right)^{2/p} \right. \\
& \quad \left. + \int_{B_{R/2}} (|A| + |B|) dX \right] \\
& \leq C \left[ \left( \frac{\rho}{R} \right)^{2-2/p} \int_{B_{R/2}} |DU|^2 dX + R^{2-4/p} \left( \int_{B_{R/2}} (|A|^{p/2} + |B|^{p/2}) dX \right)^{2/p} \right].
\end{aligned}$$

Finally, by Lemma 2.4 with  $\gamma = \beta := 2 - 4/p$ ,  $\alpha = 2 - 2/p$ , we conclude that

$$\begin{aligned}
& \int_{B_\rho} |DU|^2 dX \\
& \leq C \left[ \left( \frac{\rho}{R} \right)^{2-4/p} \int_{B_{R/2}} |DU|^2 dX + \rho^{2-4/p} \left( \int_{B_{R/2}} (|A|^{p/2} + |B|^{p/2}) dX \right)^{2/p} \right],
\end{aligned}$$

for all  $\rho < R/2$ , where the constant  $C$  does not depend on  $R$ . From (3.4) and Lemma 3.2 (i), it follows that

$$\begin{aligned}
& \int_{B_\rho} |DU|^2 dX \\
& \leq C \left( \frac{\rho}{R} \right)^{2-4/p} \int_{B_{R/2}} |DU|^2 dX + C \frac{\rho^{2-4/p}}{R^{-4/p}} \sup_{B_{R/2}} (|A| + |B|) \\
& \leq C \left( \frac{\rho}{R} \right)^{2-4/p} \left[ \int_{B_R} |DU|^2 dX + \int_{B_R} (|A| + |B|) dX \right] \\
& \leq C \left( \frac{\rho}{R} \right)^{2-4/p} \int_{B_R} |DU|^2 dX
\end{aligned}$$

for all  $\rho < R/2$ , and the arbitrariness of  $p \geq 2$ , together with standard  $C^{0,\alpha}$  regularity results (see [G], Chapter 3, Theorem 1.1), concludes the proof.  $\square$





## References

- [A1] AMBROSIO, L., A compactness theorem for a new class of functions of bounded variation, *Boll. Un. Mat. Ital.* **3-B** (1989), 857–881.
- [A2] AMBROSIO, L., A new proof of the *SBV* compactness theorem, *Calc. Var.* **3** (1995), 127–137.
- [A3] AMBROSIO, L., On the lower semicontinuity of quasiconvex integrals in *SBV*( $\Omega, \mathbb{R}^k$ ), *Nonlinear Anal.* To appear.
- [AFP] AMBROSIO, L., N. FUSCO and D. PALLARA, Partial regularity of free discontinuity sets II, *Preprint Dip. Mat. e Appl. Napoli*, (1995).
- [AP] AMBROSIO, L. and D. PALLARA, Partial regularity of free discontinuity sets I, *Preprint Sc. Norm Sup. Pisa*, (1994).
- [BZ] BLAKE, A., and A. ZISSERMAN, Visual Reconstruction, *The MIT Press, Cambridge, Massachusetts, 1985*.
- [B] BONNET, A., On the regularity of edges in the Mumford-Shah model for image segmentation. To appear.
- [CL] CARRIERO, M. and A. LEACI,  $S^k$ -valued maps minimizing the  $L^p$  norm of the gradient with free discontinuities, *Ann. Sc. Norm. Sup. Pisa* **18** (1991), 321–352.
- [DS] DAVID, G. and S. SEMMES, On the singular set of minimizers of the Mumford-Shah functional, *J. Math. Pures et Appl.* To appear.
- [DG] DE GIORGI, E., *Free Discontinuity Problems in the Calculus of Variations*, a collection of papers dedicated to J. L. Lions on the occasion of his 60<sup>th</sup> birthday, North Holland (R. Dautray ed.), 1991.
- [DGA] DE GIORGI, E. and L. AMBROSIO, Un nuovo tipo di funzionale del calcolo delle variazioni, *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur.* **82** (1988), 199–210.
- [DGCL] DE GIORGI, E., M. CARRIERO and A. LEACI, Existence theorem for a minimum problem with free discontinuity set, *Arch. Rat. Mech. Anal.* **108** (1989), 195–218.
- [D] DOUGHERTY, M., Higher integrability of the gradient for minimizers of certain polyconvex functionals in the calculus of variations. Preprint.
- [FFr] FONSECA, I. and G. FRANCFORT, Relaxation in *BV* versus quasiconvexification in  $W^{1,p}$ ; a model for the interaction between fracture and damage, *Calc. Var.* **3** (1995), 407–446.
- [FFu] FONSECA, I. and N. FUSCO, Regularity results for anisotropic image segmentation models, *Ann. Sc. Norm. Sup. di Pisa*. To appear.
- [G] GIAQUINTA, M., *Multiple Integrals in the Calculus of Variations and Nonlinear Elliptic Systems*, Annals of Mathematics Studies, Princeton University Press, 1983.
- [MS] MUMFORD, D. and J. SHAH, Boundary detection by minimizing functionals, *Proc. IEEE Conf. on Computer Vision and Pattern Recognition (San Francisco, 1985)*.

This article was processed by the author  
using the Springer-Verlag TeX mamath macro package 1990.